

**DNA polymerase eta is involved in hypermutation
occurring during immunoglobulin class switch
recombination.**

Ahmad Faili, Said Aoufouchi, Sandra Weller, Françoise Vuillier, Anne Stary,
Alain Sarasin, Claude-Agnès Reynaud, Jean-Claude Weill

► **To cite this version:**

Ahmad Faili, Said Aoufouchi, Sandra Weller, Françoise Vuillier, Anne Stary, et al.. DNA polymerase eta is involved in hypermutation occurring during immunoglobulin class switch recombination.. The Journal of Experimental Medecine, The Rockefeller University Press, 2004, 199 (2), pp.265-70. 10.1084/jem.20031831 . pasteur-00177779

HAL Id: pasteur-00177779

<https://hal-pasteur.archives-ouvertes.fr/pasteur-00177779>

Submitted on 27 Feb 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

DNA Polymerase η Is Involved in Hypermutation Occurring during Immunoglobulin Class Switch Recombination

Ahmad Faili,¹ Said Aoufouchi,¹ Sandra Weller,¹ Françoise Vuillier,² Anne Stary,³ Alain Sarasin,³ Claude-Agnès Reynaud,¹ and Jean-Claude Weill¹

¹INSERM U373, Faculté de Médecine Necker-Enfants Malades, Université Paris V, 75730 Paris, France

²Unité d'Immuno-Hématologie et d'Immunopathologie, Institut Pasteur, 75724 Paris, France

³Etude des Relations Instabilité Génétique et Cancer, CNRS UPR 2169, Institut Gustave Roussy, 94805 Villejuif, France

Abstract

Base substitutions, deletions, and duplications are observed at the immunoglobulin locus in DNA sequences involved in class switch recombination (CSR). These mutations are dependent upon activation-induced cytidine deaminase (AID) and present all the characteristics of the ones observed during V gene somatic hypermutation, implying that they could be generated by the same mutational complex. It has been proposed, based on the V gene mutation pattern of patients with the cancer-prone xeroderma pigmentosum variant (XP-V) syndrome who are deficient in DNA polymerase η (pol η), that this enzyme could be responsible for a large part of the mutations occurring on A/T bases. Here we show, by analyzing switched memory B cells from two XP-V patients, that pol η is also an A/T mutator during CSR, in both the switch region of tandem repeats as well as upstream of it, thus suggesting that the same error-prone translesional polymerases are involved, together with AID, in both processes.

Key words: somatic mutation • pol η • translesional DNA polymerases • xeroderma pigmentosum variant syndrome

Introduction

Since the discovery of activation-induced cytidine deaminase (AID) and of its central role in the initiation of the three postrearrangement B cell diversification processes, somatic hypermutation (SHM), class switch recombination (CSR), and gene conversion, a major question is to unravel what are the common and the specific partners involved in these different pathways (1, 2). SHM implies the introduction mainly of base substitutions, but also of deletions and duplications into V genes to generate an antigen receptor with better affinity for the immunizing antigen. Translesional DNA polymerases, which are able to bypass specific DNA lesions in an error-free mode at the replication fork but are highly error-prone when copying undamaged DNA, have been suspected to be involved in an error-prone short patch DNA synthesis taking place during SHM (3). It has

since been reported, based on a study of SHM in patients with xeroderma pigmentosum variant (XP-V), a cancer-prone disease corresponding to a deficiency in DNA polymerase η (pol η ; 4), that this enzyme might be responsible for a large part of the mutations occurring at A/T basepairs (5).

In CSR, the heavy chain μ constant region is switched to downstream constant region genes that are responsible for various effector functions once the antibody has bound its cognate antigen. The heavy chain μ locus region that is targeted for CSR, encompasses a large DNA region starting downstream of the I μ exon up to a region of repetitive pentamer motifs constituting the core μ switch region (S μ ; 6). Base substitutions, deletions, and duplications have been observed around switch junctions (7). More recently, mutations that display many characteristics of SHM, that is, base substitutions on RGYW hotspot motifs (with R for purine, Y for pyrimidines, and W for A or T; 8), bias for transitions and AID dependence have been described downstream of

C.-A. Reynaud and J.-C. Weill are senior authors on this paper.

Address correspondence to Jean-Claude Weill, INSERM U373, Faculté de Médecine Necker-Enfants Malades, Université Paris V, 156 Rue de Vaugirard, 75730 Paris, France. Phone: 33-1-40-61-53-80; Fax: 33-1-40-61-55-90; email: weill@necker.fr; or Claude-Agnès Reynaud, INSERM U373, Faculté de Médecine Necker-Enfants Malades, Université Paris V, 156 Rue de Vaugirard, 75730 Paris, France. Phone: 33-1-40-61-56-84; Fax: 33-1-40-61-55-90; email: reynaud@necker.fr

Abbreviations used in this paper: AID, activation-induced cytidine deaminase; CSR, class switch recombination; pol η , polymerase η ; TLS, translesion DNA synthesis; S α , α switch region; SHM, somatic hypermutation; S μ , μ switch region; XP-V, xeroderma pigmentosum variant.

the I μ exon, outside the switch core region (9, 10). Here we show, by studying hypermutation at the heavy chain locus in memory B cells of two XP-V patients, that pol η is involved in hypermutation taking place during SHM and CSR as well, implying that the same mutational complex is probably recruited during these two DNA transactions.

Materials and Methods

Characteristics of XP-V Patients and Preparation of Cell Samples. Blood samples were obtained from four healthy donors (34, 37, 41, and 50 yr old) and two French XP-V patients (55 and 56 yr old) after informed consent. The two XP-V patients studied were diagnosed clinically and biologically. Patient 1 showed numerous skin tumors since the age of 26. Patient 2 had her first malignant melanoma at 30 and numerous melanomas, basal cell carcinomas, and squamous cell carcinomas since then. Their cultured fibroblasts exhibit normal UV-C-induced unscheduled DNA synthesis and are UV-C sensitive in the presence of caffeine, a hallmark of XP-V cells (11). Heterozygous mutations of the *POLH* gene were found for both of them: a stop codon at amino acid 303 and a 4-bp deletion at amino acid 408 for patient 1, and a Gly to Arg amino acid substitution at position 295 and a 2-bp deletion (1,727–1,728) at amino acid 576 for patient 2. CD19⁺ CD27⁺ IgD⁻ memory B cells were purified by cell sorting as previously described (12) and DNA was extracted using proteinase K digestion.

PCR and Sequence Analysis. The J_{H4} intronic sequence was amplified using an FR3 consensus primer CACGGCYGTG-TATTACTGTGC and a primer upstream from J_{H5}, AGGAC-CCCAGGCAAGAAC (at 94°C for 45 s, 55°C for 1 min, and 72°C for 2 min for 40 cycles with Pfu polymerase; Stratagene). The sequence upstream from the S μ core was amplified using the following primers: 5'-S μ , GAATGATTCCATGCCAAAGC and 3'-S μ , AGCTGGATGGAGTTGTCATGGC (at 94°C for 45 s, 56°C for 1 min, and 72°C for 3 min for 35 cycles with Pfu polymerase), and the S μ - α switch region (S α) switch junctions with S μ -ext, GGGGACCTGCTCATTTTTATC and S α -ext, CCCTCAGAACCCTAAGAAC (at 94°C for 45 s, 60°C for 1 min, and 72°C for 3 min for 40 cycles using Pfu Turbo polymerase). In this last case, multiple PCRs were performed on aliquots of 2,000–5,000 cells, and PCR products below 1 kb were selected for cloning. PCR products were cloned using the TOPO TA cloning kit (Invitrogen) and sequences were determined on an ABI prism 3100 Analyzer (Applied Biosystems). All amplifications were performed using Pfu polymerase with a number of cycles to keep background mutations at a minimal level. Mutations induced during amplification by polymerases devoid of proofreading show an A/T over G/C bias that may confuse analysis, notably in the case of switch junction mutations. Error rate of Pfu polymerase is in our assay around 5×10^{-5} for 40 cycles of amplification.

Results and Discussion

DNA pol η has been proposed as an A/T mutator of Ig V genes (5). This was based on the analysis of the mutation pattern of functional V_{H6} genes from three XP-V patients, with characterized mutations of pol η for two of them. However, the question of the exact contribution of pol η to hypermutation has been challenged by Dorner and Lipsky (13). By studying one of these three patients by a single cell PCR approach, designed at amplifying both functional and nonfunctional V_H genes, they failed to observe such a deficiency at A/T targeting and concluded that it might be a bias introduced by the study of selected V gene sequences (14).

To readdress this question, CD27⁺ IgD⁻ memory B cells from two XP-V patients and from four normal individuals were purified by cell sorting and mutations were analyzed in three noncoding regions of the heavy chain locus: the J_{H4} intron, the S μ core upstream region, and S μ -S α switch junctions (Fig. 1).

The J_{H4} intron downstream from rearranged V_H genes was amplified using a 5' FR3 primer consensus for all human V_H genes, and 283 bp starting at the border of J_{H4} intron were sequenced. Only sequences corresponding to different V-D-J junctions were considered. The J_{H4} segment was used in 50% of rearranged V_H genes; such an approach allows a large sampling of unselected sequences (Fig. 2 a; reference 15). The mutation frequency was 2.0/100 bp for the controls, and 1.5 for the XP-V patients (Table I). In both groups, mutations bore the hallmark of somatic mutations: prevalence of base substitutions (with 6–9% of total mutations representing deletions or duplications), bias for transitions (47–51% of all mutations), and targeting of RGYW motifs (Table II). This targeting was calculated taking into account mutations occurring at the G position of the hotspot motif only (considered for both strands of DNA), as compared with mutations at other G/C positions. Moreover, normal controls showed a marked preference for mutations at A over T positions. In accordance with Zeng et al. (5), we observed a drastic reduction of mutations on A/T bases in XP-V patients as compared with the controls (10.7 vs. 54.0%; Fig. 2 a), thus confirming the implication of pol η . The lower number of events on A/T bases also correlates with a slight reduction in the total amount of mutations, obviously only noticeable on such unselected targets.

To find out whether the mutations observed during CSR and SHM were generated by the same mutational




Figure 1. Schematic representation of the regions selected for mutation analysis at the Ig heavy chain locus. The configuration selected represents a V_H gene rearranged to J_{H4} and switched to the α -constant region (α 1 or α 2). Horizontal arrows above the locus represent transcription initiation sites. Dotted lines between

arrows mark the DNA regions amplified and the bold line represents the DNA regions sequenced. E μ , heavy chain enhancer; L, leader; V, variable; D, diversity; and J, joining coding elements; I μ , intronic leader exon; S μ and S α , core switch regions.




Figure 2. Mutation pattern of J_H4 intronic sequences and S_μ core upstream regions in normal controls and XP-V patients. (a) Distribution and nucleotide substitution preference of mutations in J_H4 intronic sequences. The distribution of mutations is represented along the 283 bp of intronic J_H4 sequence, with the proportion of mutated clones in ordinates. Mutations from normal controls are represented above the nucleotide position, and those from XP-V patients are below. Nucleotide substitution preferences are corrected for base composition. A, 18.0%; G, 31.1%; C, 32.2%; T, 18.7%. (b) Distribution and nucleotide substitution preference of mutations in S_μ core upstream sequences. Mutations obtained for normal controls are represented above the 295 nucleotides of the S_μ core upstream sequence, and those from XP-V patients are below. The G position of the RGYW motif (and the C position of the complementary WRCY motif on the other strand) is highlighted in gray. Nucleotide substitution preferences are corrected for base composition. A, 28.5%; G, 28.8%; C: 20.0%; T, 22.7%.

complex, we analyzed two regions of the heavy chain locus that undergo hypermutation during CSR in memory B cells of both groups: the region upstream of the S_μ core region and S_μ-S_α junctions. AID-dependent mutagenesis of the S_μ core upstream region has been shown to be induced after in vitro induction of switch in mouse splenic B cells (9, 10). Mutations in this region have also been detected in human chronic lymphocytic leukemia (16).

A region of 900 bp upstream from the core repetitive sequences of the S_μ region was amplified from controls and XP-V patients, and the sequence of the 3' 295 bp was determined (16). A mutation frequency of 0.31/100 and 0.21/100 bp was found for the controls and the XP-V in-

dividuals, respectively, i.e., a value six- to sevenfold lower than for the J_H4 intron (Table I). It should be noted that both alleles are scored in this assay and that the precise fraction of nonfunctional alleles that undergo isotype switch in memory B cells in vivo is not known. The pattern of mutation was very similar to the one observed at the VDJ locus considering the ratio of base substitutions over deletions and duplications, the bias for transitions and for A over T mutations (although somewhat reduced), and the targeting to G/C bases of the RGYW (WRCY) motifs (Table II and Fig. 2 b). This similarity extends to the selective deficiency of mutations at A and T positions observed for the two XP-V patients in this genomic se-

Table I. Somatic Mutations in J_H4 Intronic Sequences and S_μ Core Upstream Sequences from Normal Individuals and XP-V Patients

	J _H 4 intronic sequence (283 bp)		S _μ core upstream sequence (295 bp)	
	Control	XP-V	Control	XP-V
Number of sequences	73	41	199	101
Total length sequenced (bp)	20,659	11,603	58,705	29,795
Unmutated sequences (percent)	2.7	2.4	54	59
Total number of mutations	414	168	180	63
(Number of deletions and duplications)	(23)	(15)	(11)	(8)
Mutations range per mutated sequence	1–25	1–19	1–11	1–5
Mutation frequency (per 100 bp)	2.0	1.5	0.31	0.21

Table III. Somatic Mutations in Sequences Flanking S μ –S α Junctions from Normal Individuals and XP-V Patients

	Controls		XP-V	
	S μ	S α	S μ	S α
		Total		Total
Number of sequences		88		38
Total length sequenced (bp)	14,992		7,588	12,054
Unmutated sequences (percent)	47	60	34	29
Total number of mutations	84	61	57	43
(Number of deletions and duplications)	(9)	(4)	(6)	(1)
Mutation range per mutated sequence	1–8	1–4	1–13	1–4
Mutation frequency (per 100 bp)	0.56	0.33	0.75	0.36
		0.43		0.50

after correction for base composition, but that the absolute number of Cs and Gs mutated was the same. The I μ sterile transcript has been shown to form an R loop with its template strand (17), which would favor the attack of the single stranded nontranscribed strand of DNA by AID according to its in vitro substrate preference (18–22). Assuming that most of the G mutations scored are due to an AID-mediated attack of cytosines on the template strand, it is tempting to propose that the lower accessibility to AID of this strand that forms a DNA–RNA hybrid is compensated by its C richness, whereas the C poorness of the accessible strand is compensated by its preferential targeting. Both factors, AID substrate preference and evolution of target sequences, would ensure approximately the same amount of AID-mediated deamination on each strand, thus allowing for efficient generation of double stranded breaks, necessary for the switch recombination process (23).

In conclusion, we would like to propose that the mutations observed during CSR most probably involve the SHM mutasome. The pattern of mutation is clearly more G/C targeted in the S μ core region. This G/C targeting might be explained by the G/C richness within pentameric hotspots in this region and the unique accessibility provided by the R loop structure, so that less nuclease digestion, and/or less error-prone short patch DNA synthesis would be required to generate adjacent cleavage sites, i.e., double stranded breaks that can be engaged into a successful switch recombination event. A similar conclusion was drawn from a study on the role of MSH2 during CSR (24) showing that this molecule is more specifically required for switching occurring in the S μ upstream region. In this region, the lower density of hotspots, as compared with the S μ core, may require more processing of the DNA ends to render them suitable for recombination, a processing that would rely on the strand displacement activity of MSH2.

This work also brings the confirmation that the translesion DNA synthesis (TLS) polymerase pol η is an A/T mutator. Nevertheless, there are still mutations on A/T bases in both SHM and CSR. Therefore, it is probable that other

TLS polymerases, such as pol ι that interacts with pol η at replication foci, could be involved in both processes (25). Accordingly, in a human Burkitt's B cell line inducible for SHM and whose pattern of mutations is strongly G/C biased, induction of SHM was shown to be dependent on pol ι (26, 27). This result remains controversial because it was recently reported that 129/SvJ mice that lack a functional pol ι did not show any significant change in the frequency and pattern of Ig mutations (28). To what extent the human in vitro and mice in vivo systems differ and to what extent TLS polymerases may compensate for each other remains to be explored.

The rationale for using error-prone polymerases during CSR is not clearly perceptible. Here we propose that these enzymes participate in an AID-triggered, short patch DNA synthesis ensuring the efficient fill-in of staggered double strand breaks. However, we cannot exclude that the mutations observed may represent the bystander recruitment of AID partners involved in the mutagenesis of Ig V genes, and that the involvement of TLS polymerases as cofactors of the nonhomologous end joining process might be dispensable. An in vitro system that would enable to assess the impact of these polymerases on the strict efficiency of switching may help resolve this issue.

We thank Dr. M.F. Avril for providing blood samples from XP-V patients, C. Debacker for cell cultures, A. Van der Linden, and the laboratory of Alan Lehmann for the determination of pol η mutations in XP-V patients 1 and 2, respectively. We thank Jérôme Mégrét for performing cell sorting, Claire Soudais and Claire Fieschi for their contribution, and Guillaume Dighiero for stimulating discussions.

This work was supported by the Ligue contre le Cancer (Equipe Labelisée) and the Fondation Princesse Grace de Monaco.

Submitted: 21 October 2003

Accepted: 24 November 2003

References

1. Honjo, T., K. Kinoshita, and M. Muramatsu. 2002. Molecular mechanism of class switch recombination: linkage with

- somatic hypermutation. *Annu. Rev. Immunol.* 20:165–196.
2. Martin, A., and M.D. Scharff. 2002. AID and mismatch repair in antibody diversification. *Nat. Rev. Immunol.* 2:605–614.
 3. Goodman, M.F., and B. Tiffin. 2000. Sloppier copier DNA polymerases involved in genome repair. *Curr. Opin. Genet. Dev.* 10:162–168.
 4. Masutani, C., M. Araki, A. Yamada, N. Dohmae, M. Yokoi, M. Yuasa, M. Araki, S. Iwai, K. Takio, and F. Hanaoka. 1999. The XPV (xeroderma pigmentosum variant) gene encodes human DNA polymerase eta. *Nature.* 399:700–704.
 5. Zeng, X., D.B. Winter, C. Kasmer, K.H. Kraemer, A.R. Lehmann, and P.J. Gearhart. 2001. DNA polymerase eta is an A-T mutator in somatic hypermutation of immunoglobulin variable genes. *Nat. Immunol.* 2:537–541.
 6. Manis, J.P., M. Tian, and F.W. Alt. 2002. Mechanism and control of class-switch recombination. *Trends Immunol.* 23: 31–39.
 7. Dunnick, W., M. Wilson, and J. Stavnezer. 1989. Mutations, duplication, and deletion of recombined switch regions suggest a role for DNA replication in the immunoglobulin heavy-chain switch. *Mol. Cell. Biol.* 9:1850–1856.
 8. Rogozin, I.B., and N.A. Kolchanov, N.A. 1992. Somatic hypermutagenesis in immunoglobulin genes. II. Influence of neighbouring base sequences on mutagenesis. *Biochim. Biophys. Acta* 1171:11–18.
 9. Nagaoka, H., M. Muramatsu, N. Yamamura, K. Kinoshita, and T. Honjo. 2002. Activation-induced deaminase (AID)-directed hypermutation in the immunoglobulin Smu region: implication of AID involvement in a common step of class switch recombination and somatic hypermutation. *J. Exp. Med.* 195:529–534.
 10. Reina-San-Martin, B., S. Difilippantonio, L. Hanitsch, R.F. Masilamani, A. Nussenzweig, and M.C. Nussenzweig. 2003. H2AX is required for recombination between immunoglobulin switch regions but not for intra-switch region recombination or somatic hypermutation. *J. Exp. Med.* 197:1767–1778.
 11. Lehmann, A.R., S. Kirk-Bell, C.F. Arlett, M.C. Paterson, P.H. Lohman, E.A. de Weerd-Kastelein, and D. Bootsma. 1975. Xeroderma pigmentosum cells with normal levels of excision repair have a defect in DNA synthesis after UV-irradiation. *Proc. Natl. Acad. Sci. USA.* 172:219–223.
 12. Weller, S., A. Faili, C. Garcia, M.C. Braun, F. Le Deist, G. de Saint Basile, O. Hermine, A. Fischer, C.-A. Reynaud, and J.-C. Weill. 2001. CD40-CD40L independent Ig gene hypermutation suggests a second B cell diversification pathway in humans. *Proc. Natl. Acad. Sci. USA.* 98:1166–1170.
 13. Dorner, T., and P.E. Lipsky. 2001. Smaller role for pol eta? *Nat. Immunol.* 2:982–984.
 14. Yavuz, S., A.S. Yavuz, K.H. Kraemer, and P.E. Lipsky. 2002. The role of polymerase eta in somatic hypermutation determined by analysis of mutations in a patient with xeroderma pigmentosum variant. *J. Immunol.* 169:3825–3830.
 15. Levy, Y., N. Gupta, F. Le Deist, C. Garcia, A. Fischer, J.-C. Weill, and C.-A. Reynaud. 1998. Defect in IgV gene somatic hypermutation in common variable immuno-deficiency syndrome. *Proc. Natl. Acad. Sci. USA.* 95:13135–13140.
 16. Oppezco, P., F. Vuillier, Y. Vasconcelos, G. Dumas, C. Magnac, B. Payelle-Brogard, O. Pritsch, and G. Dighiero. 2003. Chronic lymphocytic leukemia B cells expressing AID display dissociation between class switch recombination and somatic hypermutation. *Blood.* 101:4029–4032.
 17. Yu, K., F. Chedin, C.L. Hsieh, T.E. Wilson, and M.R. Lieber. 2003. R-loops at immunoglobulin class switch regions in the chromosomes of stimulated B cells. *Nat. Immunol.* 4:442–451.
 18. Pham, P., R. Bransteitter, J. Petruska, and M.F. Goodman. 2003. Processive AID-catalysed cytosine deamination on single-stranded DNA simulates somatic hypermutation. *Nature.* 424:103–107.
 19. Chaudhuri, J., M. Tian, C. Khuong, K. Chua, E. Pinaud, and F.W. Alt. 2003. Transcription-targeted DNA deamination by the AID antibody diversification enzyme. *Nature.* 422:726–730.
 20. Ramiro, A.R., P. Stavropoulos, M. Jankovic, and M.C. Nussenzweig. 2003. Transcription enhances AID-mediated cytidine deamination by exposing single-stranded DNA on the nontemplate strand. *Nat. Immunol.* 4:452–456.
 21. Dickerson, S.K., E. Market, E. Besmer, and F.N. Papavasiliou. 2003. AID mediates hypermutation by deaminating single stranded DNA. *J. Exp. Med.* 197:1291–1296.
 22. Sohail, A., J. Klapacz, M. Samaranyake, A. Ullah, and A.S. Bhagwat. 2003. Human activation-induced cytidine deaminase causes transcription-dependent, strand-biased C to U deaminations. *Nucleic Acids Res.* 31:2990–2994.
 23. Reynaud, C.-A., S. Aoufouchi, A. Faili, and J.-C. Weill. 2003. What role for AID: mutator, or assembler of the immunoglobulin mutasome? *Nat. Immunol.* 4:631–638.
 24. Min, I.M., C.E. Schrader, J. Vardo, T.M. Luby, N. D’Avirro, J. Stavnezer, and E. Selsing. 2003. The S μ tandem repeat region is critical for Ig isotype switching in the absence of Msh2. *Immunity.* 19:515–529.
 25. Kannouche, P., A.R. Fernandez de Henestrosa, B. Coull, A.E. Vidal, C. Gray, D. Zicha, R. Woodgate, and A.R. Lehmann. 2002. Localization of DNA polymerases eta and iota to the replication machinery is tightly co-ordinated in human cells. *EMBO J.* 21:6246–6256.
 26. Faili, A., S. Aoufouchi, Q. Gueranger, C. Zober, A. Leon, B. Bertocci, J.-C. Weill, and C.-A. Reynaud. 2002. AID-dependent somatic hypermutation occurs as a DNA single-strand event in the BL2 cell line. *Nat. Immunol.* 3:815–821.
 27. Faili, A., S. Aoufouchi, E. Flatter, Q. Guéranger, C.-A. Reynaud, and J.-C. Weill. 2002. Induction of somatic hypermutation in immunoglobulin genes is dependent on DNA polymerase iota. *Nature.* 419:944–947.
 28. McDonald, J.P., E.G. Frank, B.S. Plosky, I.B. Rogozin, C. Masutani, F. Hanaoka, R. Woodgate, and P.J. Gearhart. 2003. 129-derived strains of mice are deficient in DNA polymerase iota and have normal immunoglobulin hypermutation. *J. Exp. Med.* 198:635–643.