



HAL
open science

Early phosphatidylinositol 3-kinase/Akt pathway activation limits poliovirus-induced JNK-mediated cell death.

Arnaud Autret, Sandra Martin-Latil, Cynthia Brisac, Laurence Mousson,
Florence Colbère-Garapin, Bruno Blondel

► **To cite this version:**

Arnaud Autret, Sandra Martin-Latil, Cynthia Brisac, Laurence Mousson, Florence Colbère-Garapin, et al.. Early phosphatidylinositol 3-kinase/Akt pathway activation limits poliovirus-induced JNK-mediated cell death.. *Journal of Virology*, 2008, 82 (7), pp.3796-802. 10.1128/JVI.02020-07 . pasteur-00316053

HAL Id: pasteur-00316053

<https://pasteur.hal.science/pasteur-00316053>

Submitted on 18 Sep 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Early phosphatidylinositol 3-kinase/Akt pathway activation limits**
2 **poliovirus-induced JNK-mediated cell death**

3
4
5 Arnaud Autret, Sandra Martin-Latil, Cynthia Brisac, Laurence Mousson,
6 Florence Colbère-Garapin and Bruno Blondel*

7
8
9 Biologie des Virus Entériques, Institut Pasteur, 75724 Paris cedex 15, France

10
11 *Corresponding author. Mailing address:

12 Biologie des Virus Entériques, Institut Pasteur, 28 rue du Docteur Roux, 75724 Paris cedex
13 15, France. Phone: (33) 1.40.61.35.90; Fax: (33) 1.40.61.34.21; E-mail: bblondel@pasteur.fr

14
15 Running title: PI3K/Akt pathway limits PV-induced apoptosis

16
17
18 Word count abstract: 104

19 Word count manuscript without references, acknowledgement, figure legends: 2402

20 Number of Figures: 5

21 **ABSTRACT**

22

23 PV-induced apoptosis seems to play a major role in tissue injury in the central nervous system
24 (CNS). We have previously shown that this process involves PV-induced Bax-dependent
25 mitochondrial dysfunction mediated by early JNK activation in IMR5 neuroblastoma cells.
26 We show here that PV simultaneously activates the phosphatidylinositol 3-kinase (PI3K)/Akt
27 survival signaling pathway in these cells, limiting the extent of JNK activation, and thereby
28 cell death. JNK inhibition is associated with PI3K-dependent negative regulation of the
29 apoptosis signal-regulating kinase 1 (ASK1), which acts upstream from JNK in PV-infected
30 IMR5 cells. In poliomyelitis, this survival pathway may limit the spread of PV-induced
31 damage in the CNS.

32 Poliovirus (PV), from the *Picornaviridae* family, causes paralytic poliomyelitis — a
33 disease in which the motor neurons are destroyed in association with PV replication. PV
34 consists of a single-stranded positive RNA genome surrounded by a nonenveloped
35 icosahedral protein capsid. The human PV receptor, CD155, and its simian counterparts
36 belong to the immunoglobulin superfamily (24, 25, 31) and are related to the nectin family of
37 adhesion molecules (28, 38).

38 PV is mostly transmitted via the fecal-oral route. It first infects the oropharynx and the
39 digestive tract, and then spreads to the central nervous system (CNS) in which it mostly
40 targets motor neurons. Studies in mouse models have shown that PV-infected motor neurons
41 in the spinal cord die by apoptosis (10, 19). PV-induced apoptosis therefore seems to play a
42 major role in the tissue injury occurring in the CNS.

43 PV triggers apoptosis *in vitro* in tissue cultures of human colon carcinoma cells
44 (CaCo-2) (4), promonocytic cells (U937) (29), dendritic cells (41), murine L cells expressing
45 CD155 (21, 36), HeLa cells (8, 39) and cultures of mixed mouse primary nerve cells (12)
46 from the cerebral cortex of mice transgenic for CD155. Analyses of the apoptotic pathways
47 induced following PV infection in several cell lines have demonstrated that mitochondria are
48 key actors of PV-induced apoptosis. In particular, mitochondrial outer membrane
49 permeabilization (MOMP) following PV infection leads to a loss of mitochondrial
50 transmembrane potential and the release of proapoptotic molecules, including cytochrome *c*,
51 from the mitochondria to the cytosol (8, 21). We recently demonstrated that MOMP in PV-
52 infected neuronal IMR5 cells was dependent on Bax, a proapoptotic member of the Bcl-2
53 family. Bax activation was mediated by c-Jun NH2-terminal kinase (JNK) phosphorylation
54 after PV infection (6). JNK activation occurred early after PV infection whereas apoptotic
55 features were observed later in PV-infected cells. These events may involve a balance
56 between pro- and antiapoptotic signals following PV infection. Pro- and antiapoptotic events

57 potentially acting in synergy or competing with each other during the reproduction cycle of
58 PV have been described by Agol's group (1, 39). However, the mechanisms involved in
59 maintaining this delicate balance remain unclear.

60 Cells become committed to undergoing apoptosis in response to a collection of
61 multiple survival and death signals. The phosphatidylinositol 3-kinase (PI3K) signaling
62 pathway plays a crucial role in the transmission of survival signals in various cell types (14,
63 26), including neurons (16). PI3K activates its downstream effector, the serine-threonine
64 kinase Akt (also known as protein kinase B, PKB) by promoting its phosphorylation at the
65 residues Thr308 and Ser473. Activated Akt then phosphorylates various substrates, activating
66 antiapoptotic factors and inactivating proapoptotic factors. The role of PI3K/Akt in the
67 regulation of cell survival and apoptosis in a number of viral infection models (11, 13, 17, 27,
68 30), including infection with coxsackievirus B3 (18), rhinovirus (32), foot-and-mouth disease
69 virus (35) and enterovirus 71 (40, 43) — all members of the *Picornaviridae* family — has
70 recently been investigated.

71

72 **PV activates the PI3K/Akt survival signaling pathway in IMR5 cells**

73 We began by determining whether PV infection of IMR5 neuroblastoma cells resulted
74 in Akt activation. IMR5 cells were infected with PV as previously described (6). Briefly, the
75 growth medium (DMEM supplemented with 10% FBS) was discarded. The virus was then
76 added to monolayers at a multiplicity of infection (MOI) of ten 50% tissue culture infective
77 dose units (TCID₅₀) per cell (this MOI was used for all assays performed in this study).
78 Adsorption was allowed to proceed for 30 min at 37°C in humidified air containing 5% CO₂.
79 Cells were then washed twice with serum-free medium to remove unbound particles and
80 incubated with fresh DMEM supplemented with 10% FBS at 37°C. The virus was allowed to
81 grow for the indicated times. Time zero postinfection (p.i) corresponds to the inoculation time

82 point. Mock-infected cells were used as a negative control. As previously described (6), both
83 adherent and detached cells were taken into account in all experiments. Kinetics of Akt
84 phosphorylation at serine 473 (Ser473), which is required for full Akt activation (3), was
85 investigated in mock- and PV-infected cells. Whole-cell lysates were analyzed at the indicated
86 times p.i. by Western blotting with a specific anti-phospho (Ser473)-Akt antibody (Fig. 1A).
87 We checked for equal protein loading on the total Akt Western blot. The amount of
88 phosphorylated Akt increased until 30 min p.i., and then decreased; at 4 h p.i., the amount of
89 phosphorylated Akt present was similar to that in mock-infected cells analyzed at the same
90 time point. To check that the virus stock used in this study did not contain host-derived
91 components that may activate Akt signaling pathway, we depleted the virus suspension of PV
92 using an anti-PV antibody and infected cells with either the depleted or non-depleted
93 suspension. In contrast to cells infected with the non-depleted stock, no Akt activation (30
94 min p.i.) was detected in cells treated with the depleted suspension (Fig. 1A, bottom, left). We
95 also checked that poliovirus, purified by isopycnic CsCl gradient centrifugation (9), could
96 promote Akt activation (30 min p.i.), at an efficiency similar to that obtained with the virus
97 preparations used in this study (Fig. 1A, bottom, right). We then investigated whether Akt
98 activation in response to PV infection occurred through the PI3K pathway, by treating IMR5
99 cells with a specific PI3K inhibitor, wortmannin (5), at a concentration of 100 nM and 500
100 nM, 2 h before mock or virus infection. The concentration of the inhibitor was maintained
101 during the adsorption period and PV infection. Cell lysates were collected 30 min after
102 infection and subjected to Western blot analysis for the detection of Akt phosphorylation (Fig.
103 1B, top). Wortmannin inhibited Akt phosphorylation at both concentrations without altering
104 total Akt levels. The activation of Akt in response to PV infection was illustrated by
105 immunofluorescence staining, 30 min p.i., with the same anti-phospho (Ser473)-Akt antibody.
106 Representative staining patterns for mock-infected and PV-infected IMR5 cells treated with

107 wortmannin or left untreated are presented (Fig. 1B, bottom). As expected,
108 immunofluorescence staining was detected only in infected cells in the absence of
109 wortmannin. Thus, the rapid PV-induced phosphorylation of Akt involves a PI3K-dependent
110 mechanism.

111 We investigated whether PV adsorption onto IMR5 cells induced Akt activation in the
112 absence of PV replication by assessing Akt phosphorylation after the addition of UV-
113 inactivated PV (UV cross-linked at 6,000 $\mu\text{J}/\text{cm}^2$) to IMR5 cells at a dilution corresponding to
114 an MOI of 10 TCID₅₀ per cell (6). The complete abolition of viral infectivity by UV light
115 treatment was confirmed by titration assay with undiluted viral suspension. We also checked
116 that UV inactivation did not modify virus adsorption on cells, by comparing the binding
117 efficiency of infectious and UV light-treated PV labeled with [³⁵S]methionine (data not
118 shown). Akt phosphorylation was induced in IMR5 cells 30 minutes after the addition of UV-
119 inactivated PV, with an efficiency similar to that observed with infectious PV (Fig. 2). Thus,
120 PV-cell receptor interaction alone is sufficient to induce Akt phosphorylation in the absence
121 of viral replication.

122

123 **PI3K/Akt signaling pathway limits the amplitude of Bax activation, cytochrome *c*** 124 **release and apoptosis in PV-infected IMR5 cells**

125 We assessed the role of the PI3K/Akt signaling pathway in regulating the
126 mitochondrial pathway of apoptosis in PV-infected cells, by blocking PI3K activation with
127 wortmannin. The mitochondrial pathway is regulated by members of the Bcl-2 family,
128 including the proapoptotic protein Bax, which promotes the release of cytochrome *c*. Bax-
129 mediated cell death involves several well-controlled steps, including a conformational change
130 resulting in exposure of the NH₂-terminus. Mock- and PV-infected IMR5 cells were left
131 untreated or were treated with 100 nM wortmannin for 2 h before PV infection. The

132 concentration of the inhibitor was maintained throughout both PV adsorption and replication.
133 At 8 h p.i., a time point at which Bax activation is known to occur in PV-infected cells (6),
134 whole-cell lysates were prepared in a lysis buffer containing 1% of the zwitterionic detergent
135 CHAPS, which has no effect on Bax conformation (22). Bax was then immunoprecipitated
136 with an anti-Bax antibody (6A7) that specifically recognizes Bax protein with an exposed
137 NH₂ terminus. The Bax protein immunoprecipitated from mock- and PV-infected cells was
138 visualized by Western blotting (Fig. 3A, top). No activated Bax was detected in the
139 immunoprecipitates from mock-infected cells. Consistent with our previous report (6), Bax
140 was immunoprecipitated with the 6A7 antibody at 8 h p.i., indicating that PV infection was
141 responsible for inducing the change in Bax conformation. Wortmannin enhanced Bax
142 activation in IMR5-infected cells, without affecting the total amount of Bax (Fig. 3A,
143 bottom). The effect of wortmannin on cytochrome *c* efflux from the mitochondria of PV-
144 infected cells was also investigated. Whole-cell extracts from mock- or PV-infected cells were
145 fractionated at 8 h p.i., to separate the cytosolic fraction from the heavy membrane fraction,
146 including mitochondria, as previously described (6). Cytochrome *c* release was analyzed by
147 Western blotting the cytosolic fraction. Much more cytochrome *c* was released in response to
148 PV infection in cells treated with wortmannin than in untreated infected cells (Fig. 3B). These
149 results suggest that PI3K may inhibit Bax-dependent MOMP during the PV infection of
150 IMR5 cells.

151 We investigated the possible involvement of PV-mediated PI3K activation in the
152 inhibition of apoptosis, by analyzing the kinetics of apoptosis in mock infected and infected
153 cells treated or not treated with the specific PI3K inhibitor, wortmannin (Fig. 3C). Adherent
154 and detached cells were harvested at the indicated times p.i. and apoptosis was analyzed by
155 assessing chromatin condensation and fragmentation by flow cytometry after acridine orange
156 (AO) nuclear dye staining, as previously described (6). We found that levels of PV-induced

157 apoptosis were higher in infected cells treated with wortmannin than in untreated infected
158 cells. To confirm the role of PI3K/Akt signaling pathway in limiting PV-induced apoptosis,
159 we down-regulated Akt expression with a specific siRNA. Western blot analysis with a
160 specific antibody showed that Akt expression in IMR5 cells transfected with Akt siRNA was
161 significantly weaker than in cells transfected with a nontargeted control siRNA (Fig. 3D, left).
162 As expected, following PV infection (8 h p.i.), apoptosis levels were higher in Akt
163 knockdown cells than in nontargeted control siRNA-transfected cells (Fig. 3D, right). These
164 results suggest that PI3K/Akt pathway plays a role in inhibiting the mitochondrial apoptotic
165 pathway in PV-infected IMR5 cells.

166

167 **The PI3K/Akt signaling pathway does not affect PV growth, but delays PV release**

168 We evaluated the effects of PI3K/Akt signaling on the amount of total virus produced
169 in IMR5 cells, by determining the kinetics of total virus yield by TCID₅₀ assays in the
170 presence or absence of wortmannin. PI3K/Akt pathway inhibition had no effect on the total
171 amount of virus produced (Fig. 4). As PV-induced apoptosis levels were higher in infected
172 cells treated with wortmannin than in untreated infected cells, we assessed the possible effects
173 of the increase in apoptosis levels on externalization of the virus. Viruses were released earlier
174 in the presence of wortmannin (Fig. 4). Thus, PI3K/Akt seems to delay viral release without
175 affecting virus production.

176

177 **The PI3K/Akt signaling pathway limits JNK activation in PV-infected cells**

178 We have shown that Bax-dependent activation of the mitochondrial pathway of
179 apoptosis is mediated by early JNK activation (6). JNK activation peaks 30 min p.i and then
180 decreases in IMR5 neuroblastoma cells. It is possible that the PI3K/Akt pathway down
181 regulates the JNK pathway, as recently reported in nonviral models (2, 23).

182 We assessed the effects of PI3K/Akt on JNK activation in PV-infected cells, by
183 treating cells with wortmannin. JNK activation was investigated 30 min p.i., by Western
184 blotting whole-cell lysates with an antibody against phosphorylated forms of JNK (Fig. 5A).
185 As expected, phosphorylated JNK was detected 30 min p.i.. Larger amounts of
186 phosphorylated JNK were found in infected cells treated with wortmannin than in untreated
187 cells. Thus, activation of the PI3K/Akt pathway limits JNK activation in PV-infected IMR5
188 cells.

189

190 **JNK activation is limited by the Akt-mediated phosphorylation of ASK1 in PV-infected** 191 **cells**

192 We then examined the possibility that a kinase, upstream of JNK, was inhibited by
193 Akt, causing the observed limited JNK phosphorylation in PV-infected cells. Apoptosis
194 signal-regulating kinase 1 (ASK1) has been shown to be a key regulator of the JNK pathway
195 amenable to inhibition by Akt-mediated phosphorylation at Ser83 in nonviral systems (2, 23).
196 We assessed the possible involvement of ASK1 in JNK activation in PV-infected IMR5 cells,
197 by down-regulating ASK1 expression using specific siRNA (37). Western blot analysis with a
198 specific antibody showed that ASK1 levels were significantly lower in IMR5 cells transfected
199 with ASK1 siRNA than in cells transfected with a nontargeted control siRNA (Fig. 5B, left).
200 Moreover, following PV infection, JNK activation in ASK1 knockdown cells was weaker
201 than in cells transfected with the nontargeted control siRNA (Fig. 5B, right). Thus, ASK1
202 plays an important role in JNK activation following PV infection in IMR5 cells.

203 We then investigated the possible limitation of ASK1 activity by PI3K/Akt-mediated
204 phosphorylation at Ser83 in PV-infected cells. The kinetics of ASK1 phosphorylation at
205 Ser83 in PV-infected cells was analyzed by Western blotting with a specific antibody against
206 phosphorylated ASK1 (Fig. 5C). A transient increase in the level of ASK1 phosphorylation

207 was evident 30 minutes after infection, consistent with the pattern of Akt activation.
208 Furthermore, treatment of the cells with the PI3K inhibitor wortmannin abolished the increase
209 in ASK1 phosphorylation in PV-infected cells (Fig. 5D). Altogether, these results indicate
210 that the PI3K/Akt pathway negatively regulates JNK activation by phosphorylating and
211 inactivating ASK1 in PV-infected IMR5 cells.

212 This study provides evidence that the early PI3K/Akt survival pathway limits the
213 magnitude of PV-induced JNK activation and cell death in IMR5 cells. We previously
214 showed that PV-cell receptor interaction alone is sufficient to induce JNK phosphorylation, as
215 for Akt activation. However, we also showed that JNK phosphorylation is necessary, but not
216 sufficient, to trigger apoptosis that seems to require the active replication of PV. As
217 previously reported by Agol's group (1, 39), several different courses of events may influence
218 apoptosis in PV-infected cells between 30 min and 6-8 h p.i. These events may involve the
219 interplay between cellular and viral proteins (7, 15, 20, 33, 34, 42). Thus, the early PI3K/Akt
220 survival pathway seems to act upstream of this unidentified interplay. The PI3K/Akt pathway
221 has been shown to play an antiapoptotic role in several viral infections (11). However, this is
222 the first report, to our knowledge, of the limitation of JNK activation by PI3K/Akt mediating
223 a survival pathway during a viral infection. We have also shown that the cross-talk between
224 the PI3K/Akt and JNK pathways involved ASK1 inhibition. In poliomyelitis, this survival
225 pathway may limit the spread of PV-induced damage in the CNS.

226

227 We thank S. Susin and V. Yuste (Institut Pasteur, Paris, France) for providing IMR5 cells and
228 F. Delpeyroux (Institut Pasteur, Paris, France) for the anti-PV antibody. We also thank J.M.
229 Panaud (Institut Pasteur, Paris, France) for assistance with fluorescent microscopy. A. A. was
230 supported by grants from the Ministère de l'Education Nationale, de la Recherche et de la

231 Technologie. This work was supported by grants from the Institut Pasteur (PTR 120) and
232 Danone Research, Centre Daniel Carasso.

233

234 **REFERENCES**

235

- 236 1. **Agol, V. I., G. A. Belov, K. Bienz, D. Egger, M. S. Kolesnikova, L. I. Romanova,**
237 **L. V. Sladkova, and E. A. Tolskaya.** 2000. Competing death programs in poliovirus-
238 infected cells: commitment switch in the middle of the infectious cycle. *J Virol*
239 **74:5534-41.**
- 240 2. **Aikin, R., D. Maysinger, and L. Rosenberg.** 2004. Cross-talk between
241 phosphatidylinositol 3-kinase/AKT and c-jun NH2-terminal kinase mediates survival
242 of isolated human islets. *Endocrinology* **145:4522-31.**
- 243 3. **Alessi, D. R., M. Andjelkovic, B. Caudwell, P. Cron, N. Morrice, P. Cohen, and B.**
244 **A. Hemmings.** 1996. Mechanism of activation of protein kinase B by insulin and IGF-
245 1. *Embo J* **15:6541-51.**
- 246 4. **Ammendolia, M. G., A. Tinari, A. Calcabrini, and F. Superti.** 1999. Poliovirus
247 infection induces apoptosis in CaCo-2 cells. *J. Med. Virol.* **59:122-129.**
- 248 5. **Arcaro, A., and M. P. Wymann.** 1993. Wortmannin is a potent phosphatidylinositol
249 3-kinase inhibitor: the role of phosphatidylinositol 3,4,5-trisphosphate in neutrophil
250 responses. *Biochem J* **296 (Pt 2):297-301.**
- 251 6. **Autret, A., S. Martin-Latil, L. Mousson, A. Wirotius, F. Petit, D. Arnoult, F.**
252 **Colbère-Garapin, J. Estaquier, and B. Blondel.** 2007. Poliovirus induces Bax-
253 dependent cell death mediated by c-Jun NH2-terminal kinase. *J Virol* **81:7504-16.**
- 254 7. **Barco, A., E. Feduchi, and L. Carrasco.** 2000. Poliovirus Protease 3Cpro Kills Cells
255 by Apoptosis. *Virology* **266:352-360.**

- 256 8. **Belov, G. A., L. I. Romanova, E. A. Tolskaya, M. S. Kolesnikova, Y. A. Lazebnik,**
257 **and V. I. Agol.** 2003. The major apoptotic pathway activated and suppressed by
258 poliovirus. *J Virol* **77**:45-56.
- 259 9. **Blondel, B., O. Akacem, R. Crainic, P. Couillin, and F. Horodniceanu.** 1983.
260 Detection by monoclonal antibodies of an antigenic determinant critical for poliovirus
261 neutralization present on VP1 and on heat-inactivated virions. *Virology* **126**:707-10.
- 262 10. **Blondel, B., F. Colbère-Garapin, T. Couderc, A. Wirotius, and F. Guivel-**
263 **Benhassine.** 2005. Poliovirus, pathogenesis of poliomyelitis, and apoptosis. *Curr Top*
264 *Microbiol Immunol* **289**:25-56.
- 265 11. **Cooray, S.** 2004. The pivotal role of phosphatidylinositol 3-kinase-Akt signal
266 transduction in virus survival. *J Gen Virol* **85**:1065-76.
- 267 12. **Couderc, T., F. Guivel-Benhassine, V. Calaora, A. S. Gosselin, and B. Blondel.**
268 2002. An ex vivo murine model to study poliovirus-induced apoptosis in nerve cells. *J*
269 *Gen Virol* **83**:1925-30.
- 270 13. **Dahl, J., A. Jurczak, L. A. Cheng, D. C. Baker, and T. L. Benjamin.** 1998.
271 Evidence of a role for phosphatidylinositol 3-kinase activation in the blocking of
272 apoptosis by polyomavirus middle T antigen. *J Virol* **72**:3221-6.
- 273 14. **Datta, S. R., A. Brunet, and M. E. Greenberg.** 1999. Cellular survival: a play in
274 three Akts. *Genes Dev* **13**:2905-27.
- 275 15. **Dodd, D. A., T. H. Giddings, Jr., and K. Kirkegaard.** 2001. Poliovirus 3A protein
276 limits interleukin-6 (IL-6), IL-8, and beta interferon secretion during viral infection. *J*
277 *Virol* **75**:8158-65.
- 278 16. **Dudek, H., S. R. Datta, T. F. Franke, M. J. Birnbaum, R. Yao, G. M. Cooper, R.**
279 **A. Segal, D. R. Kaplan, and M. E. Greenberg.** 1997. Regulation of neuronal
280 survival by the serine-threonine protein kinase Akt. *Science* **275**:661-5.

- 281 17. **Ehrhardt, C., H. Marjuki, T. Wolff, B. Nurnberg, O. Planz, S. Pleschka, and S.**
282 **Ludwig.** 2006. Bivalent role of the phosphatidylinositol-3-kinase (PI3K) during
283 influenza virus infection and host cell defence. *Cell Microbiol* **8**:1336-48.
- 284 18. **Esfandiarei, M., H. Luo, B. Yanagawa, A. Suarez, D. Dabiri, J. Zhang, and B. M.**
285 **McManus.** 2004. Protein kinase B/Akt regulates coxsackievirus B3 replication
286 through a mechanism which is not caspase dependent. *J Virol* **78**:4289-98.
- 287 19. **Girard, S., T. Couderc, J. Destombes, D. Thiesson, F. Delpeyroux, and B.**
288 **Blondel.** 1999. Poliovirus induces apoptosis in the mouse central nervous system. *J.*
289 *Virol.* **73**:6066-6072.
- 290 20. **Goldstaub, D., A. Gradi, Z. Bercovitch, Z. Grossmann, Y. Nophar, S. Luria, N.**
291 **Sonenberg, and C. Kahana.** 2000. Poliovirus 2A Protease Induces Apoptotic Cell
292 Death. *Mol. Cell. Biol.* **20**:1271-1277.
- 293 21. **Gosselin, A. S., Y. Simonin, F. Guivel-Benhassine, V. Rincheval, J. L. Vayssiere,**
294 **B. Mignotte, F. Colbère-Garapin, T. Couderc, and B. Blondel.** 2003. Poliovirus-
295 induced apoptosis is reduced in cells expressing a mutant CD155 selected during
296 persistent poliovirus infection in neuroblastoma cells. *J Virol* **77**:790-8.
- 297 22. **Hsu, Y. T., and R. J. Youle.** 1998. Bax in murine thymus is a soluble monomeric
298 protein that displays differential detergent-induced conformations. *J Biol Chem*
299 **273**:10777-83.
- 300 23. **Kim, A. H., G. Khursigara, X. Sun, T. F. Franke, and M. V. Chao.** 2001. Akt
301 phosphorylates and negatively regulates apoptosis signal-regulating kinase 1. *Mol Cell*
302 *Biol* **21**:893-901.
- 303 24. **Koike, S., H. Horie, I. Ise, A. Okitsu, M. Yoshida, N. Iizuka, K. Takeuchi, T.**
304 **Takegami, and A. Nomoto.** 1990. The poliovirus receptor protein is produced both as
305 membrane-bound and secreted forms. *Embo J* **9**:3217-24.

- 306 25. **Koike, S., I. Ise, Y. Sato, H. Yonekawa, O. Gotoh, and A. Nomoto.** 1992. A 2nd
307 gene for the African green monkey poliovirus receptor that has no putative N-
308 glycosylation site in the functional N-terminal immunoglobulin-like domain. *J. Virol.*
309 **66:7059-7066.**
- 310 26. **Lawlor, M. A., and D. R. Alessi.** 2001. PKB/Akt: a key mediator of cell
311 proliferation, survival and insulin responses? *J Cell Sci* **114:2903-10.**
- 312 27. **Lee, C. J., C. L. Liao, and Y. L. Lin.** 2005. Flavivirus activates phosphatidylinositol
313 3-kinase signaling to block caspase-dependent apoptotic cell death at the early stage of
314 virus infection. *J Virol* **79:8388-99.**
- 315 28. **Lopez, M., F. Eberle, M. G. Mattei, J. Gabert, F. Birg, F. Bardin, C. Maroc, and**
316 **P. Dubreuil.** 1995. Complementary DNA characterization and chromosomal
317 localization of a human gene related to the poliovirus receptor-encoding gene. *Gene*
318 **155:261-5.**
- 319 29. **Lopez-Guerrero, J. A., M. Alonso, F. Martin-Belmonte, and L. Carrasco.** 2000.
320 Poliovirus induces apoptosis in the human U937 promonocytic cell line. *Virology*
321 **272:250-256.**
- 322 30. **Meili, R., P. Cron, B. A. Hemmings, and K. Ballmer-Hofer.** 1998. Protein kinase
323 B/Akt is activated by polyomavirus middle-T antigen via a phosphatidylinositol 3-
324 kinase-dependent mechanism. *Oncogene* **16:903-7.**
- 325 31. **Mendelsohn, C. L., E. Wimmer, and V. R. Racaniello.** 1989. Cellular receptor for
326 poliovirus: molecular cloning, nucleotide sequence, and expression of a new member
327 of the immunoglobulin superfamily. *Cell* **56:855-65.**
- 328 32. **Newcomb, D. C., U. Sajjan, S. Nanua, Y. Jia, A. M. Goldsmith, J. K. Bentley, and**
329 **M. B. Hershenson.** 2005. Phosphatidylinositol 3-kinase is required for rhinovirus-
330 induced airway epithelial cell interleukin-8 expression. *J Biol Chem* **280:36952-61.**

- 331 33. **Neznanov, N., K. P. Chumakov, A. Ullrich, V. I. Agol, and A. V. Gudkov.** 2002.
332 Unstable receptors disappear from cell surface during poliovirus infection. *Med Sci*
333 *Monit* **8**:BR391-6.
- 334 34. **Neznanov, N., A. Kondratova, K. M. Chumakov, B. Angres, B. Zhumabayeva, V.**
335 **I. Agol, and A. V. Gudkov.** 2001. Poliovirus protein 3A inhibits tumor necrosis
336 factor (TNF)-induced apoptosis by eliminating the TNF receptor from the cell surface.
337 *J Virol* **75**:10409-20.
- 338 35. **Peng, J. M., S. M. Liang, and C. M. Liang.** 2004. VP1 of foot-and-mouth disease
339 virus induces apoptosis via the Akt signaling pathway. *J Biol Chem* **279**:52168-74.
- 340 36. **Romanova, L. I., G. A. Belov, P. V. Lidsky, E. A. Tolskaya, M. S. Kolesnikova, A.**
341 **G. Evstafieva, A. B. Vartapetian, D. Egger, K. Bienz, and V. I. Agol.** 2005.
342 Variability in apoptotic response to poliovirus infection. *Virology* **331**:292-306.
- 343 37. **Saadatzadeh, M. R., K. Bijangi-Vishehsaraei, P. Hong, H. Bergmann, and L. S.**
344 **Haneline.** 2004. Oxidant hypersensitivity of Fanconi anemia type C-deficient cells is
345 dependent on a redox-regulated apoptotic pathway. *J Biol Chem* **279**:16805-12.
- 346 38. **Takahashi, K., H. Nakanishi, M. Miyahara, K. Mandai, K. Satoh, A. Satoh, H.**
347 **Nishioka, J. Aoki, A. Nomoto, A. Mizoguchi, and Y. Takai.** 1999. Nectin/PRR: an
348 immunoglobulin-like cell adhesion molecule recruited to cadherin-based adherens
349 junctions through interaction with Afadin, a PDZ domain-containing protein. *J Cell*
350 *Biol* **145**:539-49.
- 351 39. **Tolskaya, E. A., L. Romanova, M. S. Kolesnikova, T. A. Ivannikova, E. A.**
352 **Smirnova, N. T. Raikhlin, and V. I. Agol.** 1995. Apoptosis-inducing and apoptosis-
353 preventing functions of poliovirus. *J. Virol.* **69**:1181-1189.

- 354 40. **Tung, W. H., C. C. Sun, H. L. Hsieh, S. W. Wang, J. T. Horng, and C. M. Yang.**
355 2007. EV71 induces VCAM-1 expression via PDGF receptor, PI3-K/Akt, p38 MAPK,
356 JNK and NF-kappaB in vascular smooth muscle cells. *Cell Signal* **19**:2127-37.
- 357 41. **Wahid, R., M. J. Cannon, and M. Chow.** 2005. Dendritic cells and macrophages are
358 productively infected by poliovirus. *J Virol* **79**:401-9.
- 359 42. **Weidman, M., P. Yalamanchili, B. Ng, W. Tsai, and A. Dasgupta.** 2001. Poliovirus
360 3C protease-mediated degradation of transcriptional activator p53 requires a cellular
361 activity. *Virology* **291**:260-270.
- 362 43. **Wong, W. R., Y. Y. Chen, S. M. Yang, Y. L. Chen, and J. T. Horng.** 2005.
363 Phosphorylation of PI3K/Akt and MAPK/ERK in an early entry step of enterovirus
364 71. *Life Sci* **78**:82-90.

365

366 **FIGURE LEGENDS**

367

368 **Fig. 1. PV induces early Akt phosphorylation in a PI3K-dependent manner in IMR5**
369 **neuroblastoma cells**

370 **(A)** Kinetics of Akt activation in PV-infected neuronal cells. (Top) Akt activation was
371 analyzed in whole-cell lysates at the indicated times p.i., by Western blotting with a specific
372 anti-phospho (Ser473)-Akt antibody (Cell Signaling). Whole-cell lysates from mock-infected
373 cells were analyzed at 30 min (first lane) and 240 min (last lane) post mock-infection,
374 respectively. Blots were then stripped and reprobbed with an antibody recognizing all forms of
375 Akt (Cell Signaling), to confirm equal protein loading. (Bottom) Western blot analyses of Akt
376 activation 30 min p.i.. (Left) Cells were infected with viral stock (PV) or viral stock depleted
377 of PV (PV^{depleted}) with anti-PV antibody. (Right) Cells were infected with viral stock (PV) or
378 CsCl-purified PV (PV^{purified}). **(B)** Inhibition of Akt phosphorylation during PV infection in

379 IMR5 cells treated with the PI3K inhibitor, wortmannin (Calbiochem, 100 nM and 500 nM).
380 (Top) Cells were incubated or not incubated with the PI3K inhibitor for 2 h before PV
381 infection, and the concentration of the inhibitor was maintained during the adsorption period
382 and throughout PV infection. Levels of phospho (Ser473)-Akt in whole-cell lysates were
383 determined by Western blotting, 30 min p.i.. Blots were then stripped and reprobed with an
384 antibody recognizing all forms of Akt, to confirm equal protein loading. (Bottom) Mock- and
385 PV-infected IMR5 cells (30 min p.i.), treated or not treated with wortmannin (100 nM), were
386 stained for immunofluorescence with a specific antibody against phospho (Ser473)-Akt and a
387 secondary, fluorescein isothiocyanate-conjugated antibody (green) (middle panel). Nuclei
388 were stained with 4',6-diamidino-2-phenylindole (DAPI) (blue) (left panel). Merge, overlay of
389 the DAPI image with the anti-phospho (Ser473)-Akt image (right panel).

390

391 **Fig. 2. UV-inactivated PV induces early Akt activation in IMR5 cells.** Akt activation was
392 analyzed by Western blotting whole-cell lysates from cells infected with infectious or UV-
393 inactivated PV (30 min p.i.) with specific anti-phospho (Ser473)-Akt antibody. Blots were
394 then stripped and reprobed with an antibody recognizing all forms of Akt, to confirm equal
395 protein loading.

396

397 **Fig. 3. Inhibition of the PI3K/Akt signaling pathway enhances PV-induced apoptosis in**
398 **IMR5 cells**

399 **(A)** Enhancement of Bax activation in PV-infected cells treated with wortmannin. (Top) Cells
400 were uninfected or infected with PV (8 h p.i.) in the presence or absence of wortmannin (100
401 nM). Cells were lysed in immunoprecipitation buffer. Conformationally active Bax protein
402 was immunoprecipitated (IP) with anti-Bax 6A7 antibody (Santa-Cruz) and precipitates were
403 immunoblotted with anti-Bax antibody. The asterisk indicates immunoglobulin light chains.

404 (Bottom) Whole-cell lysates not incubated with 6A7 antibody were similarly tested for total
405 Bax by immunoblotting with a specific antibody (Upstate) to check that the amounts of Bax
406 protein in samples before immunoprecipitation were equivalent. Actin was used as a control
407 for protein loading. **(B)** Greater cytochrome *c* (Cyt *c*) release in PV-infected cells treated with
408 wortmannin. Cytochrome *c* release was analyzed in cytosolic fractions of mock-infected and
409 PV-infected IMR5 cells (8 h p.i.) treated or not treated with wortmannin (100 nM) by
410 Western blotting with a specific antibody (BD Pharmingen). Actin was used as a protein
411 loading control. Protein levels were determined by densitometry and plotted as ratios relative
412 to the actin levels. **(C)** Enhancement of apoptosis in PV-infected cells treated with
413 wortmannin. Mock-infected and PV-infected IMR5 cells treated (black) or not treated (light
414 gray) with wortmannin (100 nM) were analyzed at the indicated times p.i. by flow cytometry
415 after Acridine Orange (AO, Molecular Probes) staining, and the increase (*n*-fold) in apoptosis
416 was calculated as the ratio of the percentage of PV-infected IMR5 cells that were apoptotic to
417 the percentage of mock-infected cells that were apoptotic. Data are means from three
418 independent experiments. Error bars represent the standard errors of the means. *, $P < 0.05$ by
419 Student's *t* test comparing untreated IMR5 cells to treated IMR5 cells. **(D)** Higher levels of
420 apoptosis were observed after the knockdown of Akt expression in PV-infected cells. (Left)
421 IMR5 cells were transfected with Akt siRNA (Cell Signaling) or nontargeted control siRNA
422 (Cell Signaling) or left untreated. Akt protein was then assayed by immunoblotting with
423 extracts from nontargeted control siRNA-transfected, Akt siRNA-transfected or untreated
424 cells. Actin was used as a protein loading control. (Right) Cells were uninfected or were
425 infected (8 h p.i.) with PV 72 h after transfection, and cells were analyzed by flow cytometry
426 after AO staining and the increase (*n*-fold) in apoptosis was calculated as the ratio of the
427 percentage of PV-infected IMR5 cells that were apoptotic to the percentage of mock-infected
428 cells that were apoptotic. Data are means from three independent experiments. Error bars

429 represent the standard errors of the means. *, $P < 0.05$ by Student's *t* test comparing untreated
430 IMR5 cells to treated IMR5 cells.

431

432 **Fig. 4. Effect of PI3K/Akt signaling inhibition on PV growth and externalization**




433 IMR5 cells were infected with PV in the presence or absence of wortmannin (100 nM). Total
434 virus yield (extracellular and intracellular) was determined by TCID₅₀ assay at the indicated
435 times after three cycles of freezing and thawing to release intracellular viruses. Extracellular
436 virus titer was determined from the supernatant of PV-infected cells at the indicated times
437 after the removal of detached cells by centrifugation. Each point represents the mean virus
438 titers for two independent experiments. Standard errors of the mean are indicated, * $P < 0.05$ by
439 a Student *t* test comparing untreated to treated IMR5 cells.

440

441 **Fig. 5. The PI3K/Akt signaling pathway limits JNK activation by promoting ASK1**
442 **phosphorylation in PV-infected IMR5 cells**

443 **(A)** JNK activation levels are higher in PV-infected cells treated with wortmannin. Cells were
444 uninfected or infected with PV (30 min p.i.) in the presence or absence of wortmannin (100
445 nM). JNK activation was analyzed in whole-cell lysates, by Western blotting with a specific
446 anti-phospho (Thr183/Tyr185)-JNK (p46 [JNK1] and p54 [JNK2/3]) antibody, as previously
447 described (6). Blots were then stripped and re probed with an antibody recognizing all forms
448 of JNK, to confirm equal protein loading. **(B)** Inhibition of JNK activation after the
449 knockdown of ASK1 expression in PV-infected IMR5 cells. (Left) IMR5 cells were
450 transfected with ASK1 siRNA (37) or nontargeted control siRNA (Cell Signaling) or left
451 untreated. ASK1 protein was then assayed by immunoblotting with extracts from nontargeted
452 control siRNA-transfected, ASK1 siRNA-transfected or untreated cells. Actin was used as a
453 protein loading control. (Right) Untreated, nontargeted control and ASK1 siRNA transfected

454 IMR5 cells were uninfected or infected with PV. JNK activation was analyzed (30 min p.i.) in
455 whole-cell lysates, by Western blotting with a specific anti-phospho (Thr183/Tyr185)-JNK
456 antibody. Blots were then stripped and reprobbed with an antibody recognizing all forms of
457 JNK, to confirm equal protein loading. Phosphorylated JNK protein levels were determined
458 by densitometry and plotted as the ratios, relative to the levels of total JNK. Phosphorylated
459 JNK levels following PV infection in untreated cells were taken as 100%. Data are means
460 from three independent experiments. Error bars represent the standard errors of the means. *,
461 $P < 0.05$ by Student's *t* test comparing nontargeted control siRNA-transfected IMR5 cells to
462 ASK1 transfected IMR5 cells. **(C)** Phosphorylation of ASK1 in PV-infected neuronal cells.
463 ASK1 phosphorylation was analyzed in whole-cell lysates at the indicated times p.i., by
464 Western blotting with a specific anti-phospho (Ser83)-ASK1 antibody (Cell Signaling). Blots
465 were then stripped and reprobbed with an antibody recognizing all forms of ASK1 (Cell
466 Signaling), to confirm equal protein loading. **(D)** Inhibition of PV-induced ASK1
467 phosphorylation by the PI3K/Akt pathway inhibitor wortmannin. Cells were uninfected or
468 infected with PV in the presence or absence of wortmannin (100 nM). ASK1 phosphorylation
469 was analyzed (30 min p.i.) in whole-cell lysates by Western blotting with a specific anti-
470 phospho (Ser83)-ASK1 antibody. Blots were then stripped and reprobbed with an antibody
471 recognizing all forms of ASK1, to confirm equal protein loading. Phosphorylated ASK1
472 protein levels were determined by densitometry, and plotted as the ratios relative to the levels
473 of total ASK1.

A**B****Figure 1**








Figure 2

A**B****C****D****Figure 3**








Figure 4

A**B****C****D****Figure 5**