

Preferential Amplification of CD8 Effector-T Cells after Transcutaneous Application of an Inactivated Influenza Vaccine: A Randomized Phase I Trial

Behazine Combadière^{1,2,3,*}, Annika Vogt^{4,9}, Brice Mahé^{1,2,3}, Dominique Costagliola^{2,5}, Sabrina Hadam⁴, Olivia Bonduelle^{1,2,3}, Wolfram Sterry⁴, Shlomo Staszewski⁶, Hans Schaefer⁴, Sylvie van der Werf⁷, Christine Katlama^{2,3,8}, Brigitte Autran^{1,2,3,9}, Ulrike Blume-Peytavi⁴

1 Institut National de Santé et de Recherche Médicale, INSERM U945, Paris, France, **2** Université Pierre et Marie Curie (UPMC - Univ Paris 06), Paris, France, **3** Assistance Publique – Hôpitaux de Paris (AP/HP), Laboratory of Immunity and Infection, Paris, France, **4** Clinical Research Center for Hair and Skin Science, Department of Dermatology and Allergy, Charité - Universitätsmedizin Berlin, Berlin, Germany, **5** INSERM U720, Paris, France, **6** Hôpital Pitié Salpêtrière, Service des Maladies Infectieuses, Paris, France, **7** Pasteur Institute of Paris, Université Denis Diderot – Paris VII, Centre National de Recherche Scientifique URA1966, Laboratoire de génétique moléculaire des virus respiratoires, Paris, France, **8** HIV Treatment & Research Unit, Department of Internal Medicine II, Hospital of the Johann Wolfgang Goethe-University Frankfurt am Main, Frankfurt am Main, Germany, **9** Centre d'Investigation Clinique at INSERM U945, Paris, France

Abstract

Background: Current conventional vaccination approaches do not induce potent CD8 T-cell responses for fighting mostly variable viral diseases such as influenza, avian influenza viruses or HIV. Following our recent study on vaccine penetration by targeting of vaccine to human hair follicular ducts surrounded by Langerhans cells, we tested in the first randomized Phase-Ia trial based on hair follicle penetration (namely transcutaneous route) the induction of virus-specific CD8 T cell responses.

Methods and Findings: We chose the inactivated influenza vaccine – a conventional licensed tetanus/influenza (TETAGRI[®]) vaccine – to compare the safety and immunogenicity of transcutaneous (TC) versus IM immunization in two randomized controlled, multi-center Phase I trials including 24 healthy-volunteers and 12 HIV-infected patients. Vaccination was performed by application of inactivated influenza vaccine according to a standard protocol allowing the opening of the hair duct for the TC route or needle-injection for the IM route. We demonstrated that the safety of the two routes was similar. We showed the superiority of TC application, but not the IM route, to induce a significant increase in influenza-specific CD8 cytokine-producing cells in healthy-volunteers and in HIV-infected patients. However, these routes did not differ significantly for the induction of influenza-specific CD4 responses, and neutralizing antibodies were induced only by the IM route. The CD8 cell response is thus the major immune response observed after TC vaccination.

Conclusions: This Phase Ia clinical trial (Manon05) testing an anti-influenza vaccine demonstrated that vaccines designed for antibody induction by the IM route, generate vaccine-specific CD8 T cells when administered transcutaneously. These results underline the necessity of adapting vaccination strategies to control complex infectious diseases when CD8 cellular responses are crucial. Our work opens up a key area for the development of preventive and therapeutic vaccines for diseases in which CD8 cells play a crucial role.

Trial Registration: Clinicaltrials.gov NCT00261001

Citation: Combadière B, Vogt A, Mahé B, Costagliola D, Hadam S, et al. (2010) Preferential Amplification of CD8 Effector-T Cells after Transcutaneous Application of an Inactivated Influenza Vaccine: A Randomized Phase I Trial. PLoS ONE 5(5): e10818. doi:10.1371/journal.pone.0010818

Editor: Esper Georges Kallas, University of Sao Paulo, Brazil

Received: June 23, 2009; **Accepted:** March 5, 2010; **Published:** May 26, 2010

Copyright: © 2010 Combadière et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was funded by ORVACS (Objectifs Recherches VACCins SIDA). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. B. Mahe was supported by a grant from the Ministère de la Recherche et des Technologies. B. Combadière is a recipient of a Young Investigator Award from the Agence Nationale de Recherche, Fondation pour la Recherche Médicale and Interface-INSERM-AP/HP programs.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: behazine.combadiere@upmc.fr

These authors contributed equally to this work.

Introduction

Inducing CD8 T cell-mediated protective responses would be beneficial in eliminating infected cells and limiting virus or cancer dissemination. Classical preventive vaccines, however, except for live viral vectors and multiple DNA immunizations, are designed to generate neutralizing antibodies. The use of live attenuated

vaccines known to induce strong CD8 T cell responses is limited by the risk of uncontrolled virus dissemination in immunocompromised individuals (e.g., with HIV or elderly) as well as by vector or pathogen-specific pre-existing immunity that limits the efficacy of vaccine administration or readministration [1–4]. The development of successful vaccines against HIV, malaria, tuberculosis, and cancers will require efficient, potent, and durable T cell

responses [5–8]. In some cases involving high virus variability, the cross-reactivity of CD8 responses may be beneficial for recall responses [7,9]. Although there is still no clear definition of the quality of effector T cells required for protection, it is commonly accepted that one of its fundamental characteristics is the magnitude and the nature of T cell responses [10]. Recently, benchmarks were determined for primary CD8+ T cell responses in humans induced by two of the most effective vaccines ever developed, those against yellow fever and smallpox [11]. The importance of these responses has been shown in many viral diseases and cancers, in both mouse and human models [12–15], and their persistence has been observed in the absence of circulating antigens [16–19]. The generation of such immune cells is thus of crucial interest in studying long-term immune responses to pathogens and in vaccine development.

Recent advances in understanding the central role of antigen-presenting cells (APCs) in the skin have prompted numerous studies of this organ as an immunization route [20–23]. It has been suggested that differential targeting of epidermal or dermal APCs might also produce differential immune responses [21,23]. The main routes of immunization in humans – the muscle and the subcutaneous layer – are low in dendritic cells (DCs), and vaccines injected by these routes generally require adjuvant to augment DC recruitment and activation and to improve their immunogenicity [24]. Moreover, recent reports of the involvement of epithelial DCs in CD8 cell cross-priming suggests that vaccination via the cutaneous route may help to induce cellular immune responses [25–27]. Numerous concepts for vaccine delivery to the skin have thus been developed, but have not yet met expectations.

Hence, strong evidence indicates that targeting vaccine to the skin should effectively induce cellular immune responses [24,28]. Glenn and collaborators elegantly demonstrated the efficacy of transcutaneous (TC) immunization in inducing humoral immune responses in humans [25,29–34]. Frerichs et al [35] recently introduced a skin preparation system for improved TC vaccine delivery based on skin surface abrasion with silicone carbide particles, eliciting humoral responses. However, the induction of T cell immune responses, so well documented in murine models after TC immunization, remains to be shown in human.

We previously demonstrated that penetration of topically applied nanoparticles increased after application of cyanoacrylate skin surface stripping (CSSS) to human skin explants: the particles entered epidermal Langerhans cells (LCs), possibly via hair follicles [36]. We recently proposed that the cellular responses we observed to vaccine compounds were induced by the vaccine's penetration through hair follicular ducts, which are surrounded by APCs (LCs and DCs) [36,37]. In contrast to the interfollicular epidermis, the hair follicle infundibulum must be considered highly permeable, and skin DCs, including epidermal LCs, all residing in and around the hair follicles, may thus be highly accessible to topically applied vaccines [38]. In a first pilot study by our groups, we targeted an inactivated influenza vaccine at the hair follicles, with a protocol that used only a single CSSS procedure before the vaccine application. We found that TC vaccination after one CSSS procedure was safe and, most interestingly, effective in inducing cellular immune responses [37].

Questions nonetheless remain about how best to shape the nature and quality of human immune responses to vaccines. In particular, we wondered whether the site of antigen delivery would affect the nature and the quality of the immune response and whether this method of targeting the skin's APCs would be capable of inducing CD8 T cell responses to a conventional inactivated influenza vaccine designed to induce vaccine-specific antibodies. We thus conducted two randomized controlled phase I clinical

trials simultaneously in two groups of healthy individuals and a group of HIV-infected patients. Because this strategy can be of major help for HIV vaccination, we proposed to include HIV patients in parallel to healthy subjects, for safety and immunogenicity of this new method of vaccination. The transcutaneous vaccination protocol used is based on a single CSSS procedure that allows the opening of the hair follicular duct prior to application of a combined tetanus and influenza (TETAGRIP®) vaccine, compared with the conventional intramuscular (IM) immunization. The safety of the TC vaccine application was confirmed. More importantly, we demonstrated for the first time that TC application of an influenza vaccine induced a significant increase in influenza-specific CD8 responses compared with the IM route. The effects of this application on the intensity and quality of the influenza-specific effector T cells were studied in detail.

Results

Safety of TC compared with IM vaccination by inactivated influenza/tetanus vaccine in healthy volunteers and HIV-infected patients

In this randomized, investigator-blinded comparative Phase I study, we first evaluated the safety of TC compared to IM administration of a licensed non-adjuvanted tetanus+inactivated influenza vaccine (TETAGRIP®) in healthy volunteers (cohort I) and in HIV-infected patients (cohort II). All 24 healthy individuals completed the study protocol (12 in each group). The HIV-infected cohort was terminated early due to the onset of influenza season, and only 14 patients completed the study protocol: 6 in the TC group and 8 in the IM group. Table 1 summarizes the demographic and baseline characteristics of each group.

The primary safety endpoint was clinical local and systemic tolerance to vaccine administration by each route. Safety analyses of all 24 healthy volunteers and all 14 chronically HIV-infected patients who completed the study showed that the TC mode of administration was well tolerated in both populations (Table 2). All reactions and adverse events during the follow-up period (D1 to D28) were recorded. No serious adverse events were reported in the 28 days after TETAGRIP® administration in any recipient – healthy volunteer or HIV+ patient, TC or IM administration. Moderate local reactions occurred in 3 TC vaccinated healthy individuals (erythema at D1 in one volunteer and erythema at D3 in another one, erythema and swelling in a third volunteer). Mild local reactions occurred in IM vaccinated subjects. Three moderate systemic reactions were recorded for cohort I subjects, as described for both IM and TC groups in Table 2. Axillary adenopathy occurred in one TC vaccinated subject at D7 and D14 (Table subjects 2).

There was no significant increase in local adverse events, including erythema, itching, pain, swelling, and axillary node enlargement, or in the incidence of systemic (grade ≥ 3) adverse events (e.g., fever, myalgia, and diarrhea) in either cohort after TC compared to IM administration.

In Cohort II composed of HIV-infected individuals, moderate local reactions occurred in 6 TC vaccinated subjects. Two moderate systemic reactions were recorded for cohort II subjects, which are all detailed in Table 2.

Overall, the tolerance of the investigational TC route of vaccine administration was good.

Defect in induction of influenza-specific neutralizing Ab responses by the TC but not by IM vaccination route

Influenza vaccines are designed to induce neutralizing antibody (NAbs) responses after IM or subcutaneous (SC) administration in

Table 1. Demographic and Baseline Characteristics.

Cohort I		TC vaccination			IM vaccination	
Healthy Volunteers	n	median [range]	Q1–Q3	n	median [range]	Q1–Q3
Age	12	31 [22–40]	25–37	12	28 [18–40]	24–32
BMI*	12	22 [21–25]	21–23	12	22 [21–24]	21–23
Phototype II	8			8		
phototype III	4			4		
Phototype IV						
Cohort I		TC vaccination			IM vaccination	
HIV+ volunteers	n	median [range]	Q1–Q3	n	median [range]	Q1–Q3
Age	6	31[31–44]	33–44	8	40 [32–45]	37–45
BMI*	6	22 [21–26]	21–25	8	24 [21–25]	21–25
CD4 T cells						
Nadir (cells/ml)	6	308 [265–436]	274–365	8	266 [203–440]	226–347
baseline (cells/ml)	6	1022 [623–1257]	623–1257	8	724 [544–1082]	577–834
Baseline (%)	6	42 [35–51]	35–51	8	31 [20–40]	27–40
Vial Load (copies/ml)	6	<400		8	<400	
Phototype II	2			6		
phototype III	4			2		
Phototype IV						

*BMI Body mass Index.

doi:10.1371/journal.pone.0010818.t001

humans. NAb responses are therefore the reference criteria for evaluating the efficacy of influenza and tetanus vaccines. NAb responses were evaluated by an independent National Influenza reference center. Serum samples from D0 to D28 were simultaneously tested for strain-specific inhibition of HAI after

TETAGRIP® administration by both routes to the 24 healthy subjects (Table 3) and the 14 HIV+ patients (Table 4).

Results are reported for the geometric mean titers (GMT) of the A/CALIFORNIA/7/2004 H3N3 and A/NEW CALEDONIA/20/99 H1N1 strains at D14 and D28 in healthy volunteers

Table 2. Summary of clinical safety.

		Cohort I - Healthy Volunteers			Cohort II - HIV+ volunteers		
	n	12	12	24	12	12	24
	Severity	TC	IM	Total	TC	IM	Total
Local reactions	None	2	7	10	0	5	7
(p = 0.05)	Mild	7	5	11	0	3	5
	Moderate	3	0	3*	6	0	6#
	Severe	0	0	0	0	0	0
Systemic events	None	6	4	10	3	3	6
(p = 0.07)	Mild	4	7	11	2	4	6
	Moderate	2	1	3**	1	1	2##
	Severe	0	0	0	0	0	0

Statistical significance was set at $p < 0.05$.

*Moderate local reactions occurred in 3 TC vaccinated healthy individuals and included erythema at D1 in one volunteer and erythema at D3 in another one. The third volunteer experienced erythema and swelling at D1 and erythema at D3. No local reactions occurred in IM vaccinated subjects.

**Three moderate systemic reactions were recorded for cohort I subjects. The subject receiving IM vaccination experienced pain while breathing after sport accident on D7 and D14. Only volunteer experienced vomiting at D28 after TC vaccination. Axillary adenopathy occurred in one TC vaccinated subject at D7 and D14 and was the only systemic event, which was considered as related to the mode of administration by the investigator.

#Moderate local reactions occurred in 6 TC vaccinated HIV-infected individuals and included erythema at D3 in one volunteer, erythema at D14 in another volunteer as well as itching at D1 and D3 in two subjects. One volunteer experienced Itching at D1 followed by erythema at D3, D7, D14 and D21. One erythema and desquamation in D14, D21 and D28. No local reactions occurred in IM vaccinated subjects.

##Two moderate systemic reactions were recorded for cohort II subjects. Only volunteer reported an upper respiratory tract infection at D28 after IM vaccination. One TC vaccinated individual experienced malaise and myalgia at D1 after vaccination.

doi:10.1371/journal.pone.0010818.t002

Table 3. Anti-influenza specific neutralizing antibodies in healthy individuals.

	A/NEW CALEDONIA			A/CALIFORNIA		
	TC	IM	<i>p</i> value	TC	IM	<i>p</i> value
Prior vaccination – Day 0						
GMT	15.5	23.1	0.31	20	36.8	0.14
Seroconversion rate	2/12 (17%)	4/12 (33%)	0.64	3/13 (25%)	8/12 (67%)	0.10
Post vaccination – Day 28						
GMT	15.5	123.8	<0.001	20	208.2	0.001
Mean GMT increase >2.5	1.0	5.36	<0.001	1.0	5.66	0.003
Seroconversion rate >70%	2/12 (17%)	10/12 (83%)	0.003	3/12 (25%)	9/12 (75%)	0.003
Seroconversion rate	0/8 (0%)	5/7 (71%)	0.007	0/6 (0%)	1/4 (25%)	0.40
Significant increase in Ab titers	0/4 (0%)	3/5 (60%)	0.017	0/6 (0%)	6/8 (75%)	0.01
Seroconversion rate or significant Increase in Ab titers	0/12 (0%)	8/12 (67%)	0.001	0/12 (0%)	7/12 (58%)	0.005

Statistical significance was set at $p < .05$.
doi:10.1371/journal.pone.0010818.t003

(Cohort I, Table 3). There were no significant differences at baseline for antibody titers between the two study arms. IM vaccination induced the expected influenza-specific NAb responses to hemagglutinin with 75% seroprotection and seroconversion rates together with a significant increase in the antibody titers at D14, while TC vaccination did not. Indeed, TC vaccination did not induce any detectable influenza-specific NAb responses.

There were no significant differences at baseline for antibody titers between the two study arms in HIV-infected individuals and we found similar results among the cohort of HIV-infected volunteers with the absence of NAb responses in the TC group compared to IM group (Cohort II, Table 4).

Similar differences between the TC and IM groups were observed for the induction of NAb responses against tetanus toxin, as reported in supplemental Table S1. The two study arms (TC versus IM) thus differed significantly for all the NAb parameters we measured.

Thus, we found a striking absence of humoral responses to TC compared with IM vaccination.

Selective amplification of CD8 T cell responses after TC compared with IM vaccination in healthy individuals

The epidermal route of immunization described in our study is intended to target mostly epidermal LCs [36]. These APCs have

been shown to be more potent in inducing CD8 cells in *in vitro* studies [20–23]. In addition, effector CD4 and CD8 cells both provide cellular immune responses and can secrete multiple cytokines that reflect the quality of the effector-cell compartment of the immune responses. Multiparametric flow cytometry assays were performed to determine the relative importance of the subpopulations of influenza-specific CD3+CD4+ and CD3+CD8+ T cells that produce IL-2, IFN- γ , and TNF- α . Overlapping peptides were designed for three major influenza antigens included in the seasonal vaccine: i) H3, a recent strain not contained in seasonal influenza vaccines in Europe over the past five years, ii) H1, repeatedly present in influenza seasonal vaccines over the past five years, and iii) NP. We further analyzed IL-2, IFN- γ , and TNF- α production by CD4 and CD8 cells after *ex vivo* stimulation of T cells with overlapping 20-mer peptides of H3, H1, and NP. Because of the high variability of baseline influenza-specific T cells in healthy individuals, we measured the course of influenza-specific T cell responses from D0. Strikingly, we found that the frequencies of H3-, H1-, and NP-specific CD8 cells producing cytokines (IL-2, IFN- γ , and TNF- α) were significantly higher after TC than after IM vaccination (H3 $p = 0.0164$, H1 $p = 0.031$, NP $p = 0.007$), mainly because the levels of CD8 responses after IM administration was extremely low (Figure 1, upper panels). Influenza-specific CD4 responses were similar for both routes (H3 $p = 0.719$, H1 $p = 0.408$, NP $p = 0.299$). In addition, we found a higher proportion of positive

Table 4. Anti-influenza specific neutralizing antibodies in HIV-infected individuals.

	A/NEW CALEDONIA			A/CALIFORNIA		
	TC	IM	<i>p</i> value	TC	IM	<i>p</i> value
Prior vaccination – Day 0						
GMT	13.4	10.1	ns	29.0	19.2	ns
Seroconversion rate	0/6 (0%)	0/8 (0%)	ns	3/6 (50%)	3/8 (38%)	ns
Post vaccination – Day 28						
GMT	14.0	49.6	<0.001	26.8	129.1	<0.001
Mean GMT increase >2.5	1.0	4.9	<0.001	0.9	6.7	0.003
Seroconversion rate >70%	1/6 (17%)	5/8 (63%)	0.005	2/6 (33%)	6/8 (75%)	0.005

Statistical significance was set at $p < .05$.
doi:10.1371/journal.pone.0010818.t004

responders after TC compared with IM immunization, as depicted in Figure 1 (upper panels) (χ^2 test, H3 $p = 0.02$, H1 $p = 0.035$, NP $p = 0.035$) for influenza-specific CD8 but not CD4 responses (Fig 2, lower panels). Note the very high frequencies of CD8+cytokine+ cells against H3 proteins after TC vaccination. Overall, we observed preferential induction of CD8 responses against all three protein compounds included in the inactivated influenza vaccine when administered by the TC but not the IM route.

Representative flow cytometric analyses are shown for H3-specific CD3+CD8+ and H3-specific CD3+CD4+ cells by each vaccination route (Figure 2A, 2B). Besides the absence of any increase in influenza-specific CD8 cells after IM vaccination, we also observed significant differences in the distribution of single (SP), double (DP) and triple (TP) cytokine producing CD4 cells on D28 according to the vaccination route. Pie chart analyses summarize the differential distribution of influenza-specific CD4 cells and underline the impact of vaccination route on the quality of vaccine-specific effector T cells (Figure 2C and 2D).

In addition, a tetanus-specific cellular response was tested by IFN- γ -ELISPOT assays, as shown in supplemental Figure S1. Because the cellular response level was below the detection level of the ELISPOT assay, we did not perform further analysis of the quality of the tetanus-specific T cell responses.

Finally, we analyzed the quality of cellular immune responses in HIV+ volunteers (cohort II) after *ex vivo* stimulation of peripheral blood cells with the same overlapping peptides covering H3 (Figure 3), H1, and NP (data not shown). Intracellular cytokine staining assays were performed to assess the production of IL-2, IFN- γ , and TNF- α in CD4 and CD8 cells. We found that H3-specific CD8 cells tended to be slightly more frequent after TC than after IM vaccination in these volunteers ($p = 0.09$ at day 14 and $p = 0.2$ at day 28) (Figure 3). This was not the case for H3-specific CD4 responses ($p = 0.59$), although the small number of individuals tested made it very difficult to obtain significant results (Figure 3). Similar results were observed for H1 and NP proteins (data not shown). Note that the number of HIV+ volunteers able




Figure 1. Differential induction of CD4 and CD8 T cell responses after TC vaccine application compared to IM immunization in healthy individuals. Intracytoplasmic cytokine staining (ICS) of influenza-specific effector CD4 and CD8 responses was performed on frozen PBMC samples from vaccinated individuals: 10/12 from the TC group and 7–9/12 from IM group with 90% cell viability after thawing. Three million cells were stimulated with the overlapping peptide covering H3, H1, and NP for 12 hours at 37°C. Brefeldin A was added 4 h before harvesting. ICS was performed by flow-cytometric assays on CD3+CD4+ (left panels) and CD3+CD8+ T cells (Right panels). At least 1,000,000 live events according to forward and side scatter parameters were accumulated and analyzed (M&M section). The expression of IFN- γ , TNF- α , and/or IL-2 (triple+double+single cytokine positive cells) by influenza-specific T cells was analyzed with the Boolean gating function of FlowJo software. Results are shown as percentages of cytokine-producing T cells (Δ Day 28–Day 0) after subtracting the unstimulated cell background. Mann-Whitney test was used to compare continuous variables between the groups. Significance was set at $p < 0.05$. Responders are determined when (Δ Day 28–Day 0) were > 0 . The χ^2 test was used to define categorical variables between TC and IM groups. doi:10.1371/journal.pone.0010818.g001





A**B****C****D**

Figure 2. Flow cytometric representation of influenza-specific T cell response at day 28 post-vaccination by TC and IM routes. A, B) Representative flow cytometric analysis of cytokine-producing influenza-specific effector CD4 and CD8 responses. Experiments were performed on frozen PBMCs from individuals vaccinated by TC and IM routes as described in figure 1. Results are shown for a representative healthy individual with a TC route (A) and an IM route (B) vaccination at day 28. C, D) Pie chart analyses of single (white), double (gray) and triple (black)-cytokine positive cells for CD4 (C) and CD8 (D) effector cells specific for the indicated influenza protein. The expression of IFN- γ , TNF- α , and/or IL-2 (triple+double+single cytokine positive cells) by influenza-specific T cells was analyzed with the Boolean gating function of FlowJo software. NA: not applicable.

doi:10.1371/journal.pone.0010818.g002

to be tested was reduced because of the short period available until the flu season.

These observations suggest that the effect of the TC route on preferential amplification of vaccine-specific CD8 cells in HIV+ individuals was similar to that in healthy individuals. They need, however, to be confirmed in a larger sample of such subjects. Overall, this result demonstrates for the first time that the route of immunization affects the quality of immune responses and in particular the quality of T cells.

Discussion

We showed that hair follicular targeting of an inactivated influenza vaccine by the transcutaneous route induced preferentially CD8 effector T cells rather than humoral immune responses when compared to a conventional IM route, in a randomized, controlled Phase I clinical trial in healthy volunteers. Our work is the first demonstration that the immunization route in humans affects the magnitude and quality of CD8 T cell responses as well as the intrinsic quality of cytokine-producing CD4 effector cells after TC and IM vaccination with a seasonal influenza vaccine. We also found similar results after influenza vaccination in HIV-infected patients.

The importance of anti-viral CD8 cells as well as recent influenza pandemics (H5N1 and H1N1swine) have raised the need of re-evaluating cellular responses [39–41]. Numerous reviews and studies have highlighted the importance of respiratory virus-specific effector and memory T cell responses in humans and mice [41] that participate to the accelerated viral clearance as well as enhanced survival after viral challenge [41–43]. In addition, according to current concepts in HIV vaccine design, a broadly targeting vaccine-specific CD8 response would restrict acute HIV replication [44–47]. The HIV-infected population would constitute one of the key populations that would benefit from such novel vaccination strategies.

The aim of this approach is not yet to propose a vaccination strategy for a seasonal influenza epidemic, but rather a proof-of-concept study that the route of immunization will help in directing immune responses towards a larger pool of CD8 responses against infectious diseases. Conventional inactivated influenza vaccine was used as an example for vaccination because of its ease to use and wide acceptability by volunteers for clinical trials. The strength of our work resides on the use of this type of vaccine, that has been designed for antibody responses and not CD8 responses, rather the use of live viruses.

Vaccines are classically injected into a muscle, where the local deposit of vaccine compound must be captured by locally present and recruited APCs. Muscles, however, have very few DCs, and vaccines generally require adjuvant to improve their immunogenicity. Because our method of vaccine application results in opening the hair follicle, it should make skin DCs, including epidermal LCs, which reside in and around the hair follicles, highly accessible to topically applied vaccines [36,48]. Recent studies suggest that DC subsets play a role in generating adaptive immune responses of quantitatively and qualitatively distinct types [49,50]. The elicitation of T cell responses by epicutaneous

immunization [34,51,52] suggests that LCs play a role. It was recently shown that LCs cross-present exogenous antigen to CD8+ T cells in murine models [51,52]. Dermal DCs (in the dermis) and LCs (in the epidermis) have different migratory paths through the lymphoid tissues [53]. In addition, LCs generated *in vitro* and derived from CD34+ precursors can cross-present exogenous antigen to CD8+ cells [54] more potently than either dermal or monocyte-derived DCs [55]. The efficacy of cellular immune responses also appears to rely on strong antigen-induced priming of T cells by efficient targeting and activation of professional APCs [6,56].

In this study, vaccine was applied by a two-step protocol that included a single CSSS procedure on the upper arm followed by topical application of a total dose of a conventional influenza/tetanus vaccine. The safety and tolerability of this CSSS procedure in human volunteers was consistent with our previous findings in a pilot study [37] of a limited number of healthy volunteers, where we used CSSS and an anti-influenza vaccine (not combined with any other product). Our first published pilot study was conducted to define the surface area required for successful vaccination by this method [37]. Previous studies have already demonstrated the success of CSSS application in penetration of human skin [57,58]. This trial, however, is the first to demonstrate the safety of this CSSS procedure in HIV-infected patients. Skin physiology measurements confirmed that the skin parameters we measured did not differ significantly between the two cohorts (healthy volunteers and HIV-infected patients) or the two study arms (TC versus IM administration). The TC vaccination protocol was also safe in this specific patient population.

In recent years, various studies have recognized the feasibility of TC vaccination in mice and even in humans (for review see [24]). Safety and humoral responses to adjuvanted or live-inactivated vaccines have been widely studied in humans, but not the induction of cellular responses. In Phase I/II studies of a live-attenuated measles vaccine in humans, Etchard et al. [59] recently showed that TC – but not subcutaneous – immunization failed to induce serum antibodies and induced only limited mucosal IgA and IFN- γ responses. Yagi and colleagues explored the induction of T cell immune responses in five melanoma patients [60]. They showed that five percutaneous immunizations (monthly application of HIV and melanoma MHC-Class I restricted peptides) allowed the induction of peptide-specific CD8 cells. In that study, CSSS procedures led to complete removal of the stratum corneum, on skin areas as large as 100 cm². That study also used substantially more than our single vaccine application; however, we showed here that one application of influenza vaccine is sufficient to induce cellular responses by the TC route. Further dose-dependent vaccination needs to be performed in the future.

We previously explored the induction by TC and IM vaccination of cellular immune responses in 11 healthy volunteers (n = 6 TC and n = 4 IM) by a seasonal anti-influenza vaccine [37]. Interestingly, TC vaccination induced both CD4 and CD8 T cell responses, whereas IM injection induced only effector CD4 T cells. Nonetheless, the small sample size prevented effective statistical analysis. The Phase I trial reported here showed a significant difference according to route of administration in the




Figure 3. Differential induction of CD4 and CD8 T cell responses after TC and IM vaccine application in HIV-infected volunteers. Intracytoplasmic cytokine staining (ICS) of influenza-specific effector CD4 and CD8 responses was performed on frozen PBMCs from vaccinated individuals. Experiments were performed on 12 HIV-infected individuals for whom cell sample viability was at least 90% after thawing. Three million cells were stimulated with the overlapping peptide covering H3, H1, and NP for 12 hours at 37°C. Brefeldin A (5 µg/ml) was added 4 h before harvesting. ICS was performed for IFN-γ by flow-cytometric assay on CD3+CD4+ (left panels) and CD3+CD8+ T cells (Right panels). At least 1,000,000 live events according to forward and side scatter parameters were accumulated and analyzed (M&M section). The expression of IFN-γ, TNF-α, and/or IL-2 (triple+double+single cytokine positive cells) by influenza-specific T cells was analyzed with the Boolean gating function of FlowJo software. Results are shown as percentages of cytokine-producing T cells (Δ Day 28–Day 0) after subtracting the unstimulated cell background. Mann-Whitney tests were used to compare continuous variables between the groups. Significance was set at $p < 0.05$. doi:10.1371/journal.pone.0010818.g003

magnitude of the CD8 effector T cell population directed against three distinct influenza vaccine proteins: hemagglutinins H3 and H1 and NP. CD4 and CD8 cell response against the influenza virus have been thoroughly described. It has been shown that CD8 cells make it possible to eliminate the virus [5,61–64] as well as to control secondary infection by a lethal influenza strain in the absence of B cells and antibodies [65,66]. CD8 cells specific for influenza proteins may recognize viral epitopes such as nucleoprotein (NP), polymerase acid (PA), matrix protein (M), and nonstructural proteins [67–71]. In influenza infections in murine models, protection by CD8 T cells has been shown to be derived from restricting the dissemination of influenza A as well as of influenza variants [72–76]. In elderly humans, cellular immune responses against influenza correlated with protection against influenza virus and thus indicated the limitations of using serum antibody responses alone to measure vaccine efficacy [77,78].

Beyond the importance of CD8 immune responses in pandemic influenza and vaccination in the elderly and other immunocompromised individuals, the magnitude of antigen-specific T cell responses is a measure of vaccination efficacy against viral diseases and also cancers. Efforts to improve vaccination efficacy will help in the fight against these diseases. However, it does not reflect the functional abilities that can be analyzed for an antigen-specific T cell population. Currently flow cytometric techniques allow the analysis of multiple functions. IFN-γ, TNF, and IL-2 are analyzed most often to assess cellular immunity to infectious diseases [79,80]. It has been suggested that the quality of T cells is crucial for determining the outcome of infectious diseases [10]. However, quality is a critical point that has not yet been defined in terms of control of viral diseases. Most reports of the multifunctionality of T cells are based on studies of HIV infection or vaccination, and it remains difficult to expand or generalize these results to other infectious diseases or vaccinations. Combinations of markers can paint a more detailed picture of antigen-specific T cells. We found that the quality of polyfunctional CD8 and CD4 effector/memory cells against influenza proteins (H3, H1, and NP) was similar at baseline in both arms of the study and thus confirmed the relative homogeneity of influenza-specific effector/memory cells before vaccination. The TC and IM routes of vaccination that induced differential cytokine profiles that suggested that the quality of CD4 cell stimulation by these routes might differ. The induction of CD8 responses after IM vaccination was extremely limited and did not allow further analysis.

Nevertheless, vaccination by the TC route did not induce NAb responses. This result may also be related in part to the strength of immunization or the differential quality of the APCs, which can dictate immunological outcome.

A major obstacle to skin vaccine delivery is the stratum corneum, an important constituent of the skin barrier. Multiple approaches have explored ways of overcoming it. Interestingly, Fan et al. found that topical vaccination requires the presence of intact hair follicles, which are the most relevant physiological breaks in the skin barrier. They also highlighted the operation of efficient mechanisms within the follicle for the induction of immune responses against DNA vaccines [81]. Numerous studies have recognized the importance of hair follicles in percutaneous penetration processes [36,38,82,83]. The CSSS technique facilitates follicular penetration by removing cellular debris and sebum from the hair follicle openings [82,83]. In contrast to previous approaches, however, CSSS as it was used in this study had two important effects on the skin. It removed approximately 30% of the stratum corneum, inducing mild barrier disruption without damaging the viable epidermis and its associated cell populations (LCs), and it removed cellular debris from hair follicles, hereby

increasing the number of hair follicles available for penetration. This idea of utilizing the reservoir function of the hair follicle for TC vaccination strategies is perfectly in line with recent reports by Naito *et al.*, who found that prolongation of antigen presence increased the efficacy of TC immunization in mice [84]. While it remains difficult to evaluate the amount of antigen to which the immune system is actually exposed, this study supports the experimental evidence from many studies that TC vaccination may in fact allow dose sparing with equivalent T cell responses, self-administration, and immune enhancement, even in elderly patients with less responsive immune systems [85,86].

Materials and Methods

Study Design

The primary objective of this study was to evaluate the safety of a newly developed protocol for TC vaccine application in healthy volunteers (cohort I) and HIV-infected individuals (cohort II) by using a commercially available tetanus/influenza vaccine (Tetagr[®]), in comparison to conventional IM injections. The secondary objectives included first a comparison of the immunogenicity of TC and IM vaccination by assessing the tetanus- and influenza-specific antibody titers and cellular immune responses, and second, an evaluation of skin physiology in the patients in the TC groups. The study included two cohorts of volunteers composed of healthy individuals (cohort I, $n = 24$), and HIV-infected patients (cohort II, $n = 14$), each randomized in two arms *i.e.* TC versus IM vaccination, of 12 subjects per arm, as defined by the Data Management and Statistics Center, according to an SAS procedure plan. The volunteers were recruited at two investigational centers (Department of Dermatology and Allergy, Charité – Universitätsmedizin Berlin, Germany, and HIV Treatment & Research Unit, Department of Internal Medicine II, Johann Wolfgang Goethe University Hospital, Frankfurt, Germany). After vaccination at D0, safety data were collected at D1, D3, D14, D21, D28, and blood samples were taken for immunological analyses at indicated time points (D0, D14 and D28). Skin physiological measurements were performed before and after vaccination at indicated time points. The protocol for this trial and supporting CONSORT checklist are available as supporting information (Checklist S1, Flowchart S1 and Protocol S1).

Ethics committee approval, health authorities

The trial was conducted in accordance with the latest Declaration of Helsinki, GCP, and ICH regulatory guidelines. The study protocol, its first amendment, and the informed consent and patient information forms were reviewed and approved by the independent Ethics Committee of Charité – Universitätsmedizin Berlin, Campus Mitte, and submitted to the Ethics Committee of Johann Goethe University, Frankfurt, Germany. Approval was further obtained from the Paul-Ehrlich Institute, Germany (Federal Agency of Sera and Vaccines). Written informed consent was obtained from each volunteer before study entry. The clinical trial has the following identification number NCT00261001.

Study population

Volunteers from both cohorts had to meet the following eligibility criteria: men aged 18–45 years, body mass index (BMI) 21–26, skin phototype I–IV, clinical examination and interview for medical history, no tetanus vaccination within the past four years, no influenza vaccination within the past year, no psychological, familial, sociological or geographic condition that might impede compliance with the study protocol, and written informed consent. Men with excessive terminal hair growth on the

investigational sites were excluded from the study. Further exclusion criteria were any acute or chronic illness, skin conditions or allergies, or local or systemic treatments that might interfere with the study protocol, and past or planned sessions of ultraviolet or sun exposure within six weeks of study entry. Healthy volunteers with a negative HIV test within three months of study entry were included in cohort I. Inclusion in cohort II (HIV-infected patients) required HIV-infection, positive HIV serology, CD4 cell nadir $>200/\text{mm}^3$, effective antiretroviral treatment with a minimum of three drugs for <1 year prior to study entry with a plasma HIV RNA $<400\text{cp}/\text{ml}$ during the 6 months before the inclusion and CD4+ counts >350 cells/ mm^3 for 12 months.

Vaccine

TETAGRIP[®] is a commercially available influenza/tetanus vaccine (Sanofi-Pasteur, France). The vaccine is provided as injectable suspension of one vaccine dose containing tetanus anatoxin and inactivated influenza virus type A and type B fragments of influenza antigens equivalent to 15 μg of 2 type-A (A/CALIFORNIA/7/2004 H3N3 and A/NEW CALEDONIA/20/99 H1N1) and 1 type-B (B/SHANGAI/361/2002) virus hemagglutinin subunits in saline solution. For TC and IM vaccination, 0.5 ml of the vaccine was used, as provided by the manufacturer.

TC and IM vaccination

TC vaccination was performed as described elsewhere, with a newly developed standard operating procedure based on CSSS [36,37]. Briefly, two investigational sites of 4×4 cm each were delimited on the external part of the upper left arm with a permanent skin marker (Skin marker H7003 Falc). The investigational sites and the surrounding skin (2 cm on the top and bottom of each investigational site and 1 cm on both sides) were lightly shaved with a dry razor (Disposable razor, Art.-No. 182 H, Wilkinson Sword GmbH, Germany). One CSSS was performed with 190 mg cyanoacrylate (Superglue, UHU GmbH & Co. KG, Germany), which was spread evenly on the skin surface with a microscope slide. Adhesive tape (6 $\times$ 5 cm, Art.-No. 571176-00000, Tesa Beiersdorf, Germany) was then applied and massaged with a rubber roll to improve adhesion (10 times). After the glue hardened for 20 minutes, tape and glue were removed from the skin surface. A silicone barrier was placed to protect the area around the investigational sites (Window-Colourpaste, Art.-No. 4469/ko, Max-Bringmann GmbH & Co., Wendelstein, Germany) to avoid spreading the vaccine. TETAGRIP[®] vaccine (250 μl) was applied in droplets from the original syringe provided by the manufacturer onto the skin surface of each investigational site (16 drops per investigational site, each drop approximately 16 μl) and then gently massaged with gloved fingertip (care & serve[®]) presaturated with vaccine for one minute. After an incubation time of 20 minutes, a protective hydrocolloid bandage (Comfeel[®] Plus Transparent 9 $\times$ 14 cm Art.-Nr.: 3542, Coloplast A/S, Denmark) was applied to the dried surface to protect the investigational sites for 24 h. The volunteers were instructed not to take a shower or bath and to avoid any activity that caused sweating or mechanical stress to the investigational site, e.g., physical exercise, for these 24 hours.

For the IM group, TETAGRIP[®] vaccine (0.5 ml) as provided by the manufacturer was injected intramuscularly into the deltoid muscle of the left arm after careful disinfection, following the most recent ICH Good Clinical Practices.

Clinical safety

After each immunization the volunteers remained under medical supervision for at least 30 minutes. Safety data (local

and systemic reactions, adverse events) were assessed and recorded on D1, D3, D7, D14, D21, and D28. Each visit included an interview, review of the diary cards, and a physical examination of the volunteer. Local reactions were graded 0–3 (0 = none, 1 = mild, 2 = moderate, 3 = severe) according to the occurrence of erythema, pruritus, burning, or desquamation. Systemic reactions (e.g., rash, pain, fever, headache, shivering, diarrhea, or malaise) were similarly graded 0–3. Severe adverse events were graded 1–4 for each organ system according to clinical and laboratory parameters defined in standard toxicity tables.

Skin physiology measurements

Transepidermal water loss (TEWL), stratum corneum hydration, skin pH, and sebum production were assessed before vaccination on D0 with a Multi Probe Adapter MPA® (Courage-Khazaka, Cologne, Germany). All skin measurements were performed according to the manufacturer's recommendation.

HAI titers

Serum antibody against influenza was measured at the Institut Pasteur, the French national reference for influenza (Centre National de Référence de la Grippe) by a standard microtiter hemagglutination inhibition (HAI) assay, as previously described [87]. HAI antibody titers were determined before and after influenza vaccination in all volunteers. Serum samples from D0, D14, and D28 were simultaneously tested for strain-specific HAI. Results are reported for the A/CALIFORNIA/7/2004 strain. The humoral response to this strain contained in the influenza vaccine was assessed by calculating the geometric mean titers before vaccination and at D14 and D28, and the fold increases in the titer at D14 and D28.

Synthetic peptide design

Fifty-six overlapping 20-mer peptides covering the entire H1 hemagglutinin protein (strain A/NEW/CALEDONIA/20/99) – included in European influenza vaccines over the past five years – were synthesized (Altergen, France), and a super pool was generated. Overlapping 20-mer peptides [88] (32 for H3 and 7 for nucleoprotein, NP) covering MHC Class I and II peptides of the H3 hemagglutinin protein (strain A/CALIFORNIA/7/2004) (not included in European influenza vaccines over the past five years) and NP (strain A/NEW/CALEDONIA/20/99) were designed based on epitopes described in the literature for 48 H3 and 44 NP peptides. To visualize the protein region rich in T cell epitopes, sequences were aligned (Multalin® Software, INRA, France) between the H3 hemagglutinin protein (strain A/CALIFORNIA/7/2004) and all related T cell epitopes, thus refining the 20-mer overlapping peptides. All peptides were synthesized by Eurogentec, France.

Intracellular cytokine staining

Immunomonitoring studies were performed for the number of individuals indicated below, that is, for the subjects whose frozen blood samples were properly stored and had a cell viability superior to 90% on thawing. All experiments were performed blinded to study arm. Experiments were performed on frozen PBMC samples from most subjects in both cohorts: i) in cohort I, 10/12 members of the TC group and 7 to 9/12 members of the IM group and ii) in cohort II, 6/6 of the TC group and 6/8 of the IM group. Frozen PBMCs were thawed in RPMI (Life Technologies, France) containing 5% FCS (Seromed, Germany), 2 mmol/l L-glutamine (Gibco BRL, Life Technology, Scotland), and antibiotics (1000 UI/ml penicillin sodium, 1 mg/ml strepto-

mycin sulfate, and 250 ng/ml amphotericin B). Cells were stimulated with three different pools of 20-mer peptides (2 µg/ml) for H1, H3, and NP antigens. Brefeldin A (5 µg/ml) (Sigma Chemical Co., France) was added to the well four hours before harvesting to detect intracellular cytokines. Then cells were stained in PBS 1 × for 10 min at RT, and membrane markers were added for 20 min at 4°C. Next 100 µl of Fix and Perm Medium A (Caltag, France) was added to each sample for 10 min at RT. Cells were washed, resuspended with 100 µl of Fix and Perm Medium B (Caltag, France), and incubated with intracellular monoclonal antibodies (Abs) specific for cytokine detection for 20 min at RT. The following panel of eight Abs was used: CD4-AmCyan, CD8-PacificBlue, CD27-APC, CD45RA-ECD, IL2-FITC, IFN-γ-Alexa700, TNF-α-PE-C7. Flow cytometric analyses were done with LSRII flow cytometers (Becton Dickinson, Immunocytometry Systems). Analyses were performed with FlowJo software (Tree Star). The live lymphocyte gate (at least 1,000,000 live events) was set based on forward and side scatter for further analysis. The expression of IFN-γ, TNF-α, and IL-2 (triple+double+single cytokine-positive cells) by influenza-specific T cells was analyzed with the Boolean gating function of FlowJo software as described previously [89,90,91]. Results are shown as percentages of cytokine-producing T cells (Δ Day 28-Day 0) after subtracting the background unstimulated cells. Responders are determined when (Δ Day 28-Day 0) were superior to 0.

Statistical Analysis

The data analysis consisted of a comparison of safety and immunogenicity data between study arms. All statistical analysis was performed using SPSS 11 software or Prism 4.0c for Mac OS X for data handling and graphic representation. The analysis variables consisted of baseline variables, primary endpoints (safety variables), and secondary endpoints (immunogenicity variables). For baseline variables, descriptive analyses were performed (% , median, interquartiles, ranges) by arm in each cohort. For safety and immunogenicity the two arms were compared using non-parametric tests: Fisher exact test for qualitative variables, Mann-Whitney tests for continuous variables and χ^2 -test for categorical variables. Statistical significance was set at $p < 0.05$.

Supporting Information

Table S1 Supplemental Data Table.

Found at: doi:10.1371/journal.pone.0010818.s001 (0.04 MB DOC)

Figure S1 Tetanus-specific cellular immune responses after TC and IM vaccination.

Found at: doi:10.1371/journal.pone.0010818.s002 (0.12 MB TIF)

Protocol S1 Trial Protocol.

Found at: doi:10.1371/journal.pone.0010818.s003 (0.65 MB PDF)

Checklist S1 Consort Checklist.

Found at: doi:10.1371/journal.pone.0010818.s004 (0.19 MB DOC)

Flowchart S1 Flowchart cohort I and cohort II.

Found at: doi:10.1371/journal.pone.0010818.s005 (0.11 MB PDF)

Acknowledgments

We thank members of ORVACS (Objective Recherche Vaccination contre le SIDA) for their helpful advice and discussions. We thank Jo-ann Cahn and Dr Darragh Duffy for English editing.

Author Contributions

Conceived and designed the experiments: BC AV DC SS HS CK BA UBP. Performed the experiments: BC BM SH OB SvdW. Analyzed the

data: BC AV BM DC SvdW. Contributed reagents/materials/analysis tools: BC AV WS SS UBP. Wrote the paper: BC AV BM BA. Head of the department: WS.

References

- Wharton M, Strikas RA, Harpaz R, Rotz LD, Schwartz B, et al. (2003) Recommendations for using smallpox vaccine in a pre-event vaccination program. Supplemental recommendations of the Advisory Committee on Immunization Practices (ACIP) and the Healthcare Infection Control Practices Advisory Committee (HICPAC). *MMWR Recomm Rep* 52: 1–16.
- Patterson S, Papagatsias T, Benlahrech A (2009) Use of adenovirus in vaccines for HIV. *Handb Exp Pharmacol*. pp 275–293.
- Harro CD, Robertson MN, Lally MA, O'Neill LD, Edupuganti S, et al. (2009) Safety and immunogenicity of adenovirus-vectored near-consensus HIV type 1 clade B gag vaccines in healthy adults. *AIDS Res Hum Retroviruses* 25: 103–114.
- White A (2008) Why vaccines are not the answer - the failure of V520 and the importance of cell-mediated immunity in the fight against HIV. *Med Hypotheses* 71: 909–913.
- Doherty PC, Hou S, Tripp RA (1994) CD8+ T-cell memory to viruses. *Curr Opin Immunol* 6: 545–552.
- Kaech SM, Wherry EJ, Ahmed R (2002) Effector and memory T-cell differentiation: implications for vaccine development. *Nat Rev Immunol* 2: 251–262.
- Robinson HL (2003) T cells versus HIV-1: fighting exhaustion as well as escape. *Nat Immunol* 4: 12–13.
- Post J, Ratnarajah S, Lloyd AR (2009) Immunological determinants of the outcomes from primary hepatitis C infection. *Cell Mol Life Sci* 66: 733–756.
- Brown LE, Kelso A (2009) Prospects for an influenza vaccine that induces cross-protective cytotoxic T lymphocytes. *Immunol Cell Biol*.
- Seder RA, Darrah PA, Roederer M (2008) T-cell quality in memory and protection: implications for vaccine design. *Nat Rev Immunol* 8: 247–258.
- Miller JD, van der Most RG, Akondy RS, Glidewell JT, Albott S, et al. (2008) Human effector and memory CD8+ T cell responses to smallpox and yellow fever vaccines. *Immunity* 28: 710–722.
- Godelaine D, Carrasco J, Lucas S, Karanikas V, Schuler-Thurner B, et al. (2003) Polyclonal CTL responses observed in melanoma patients vaccinated with dendritic cells pulsed with a MAGE-3A1 peptide. *J Immunol* 171: 4893–4897.
- Boon AC, de Mutsert G, Fouchier RA, Osterhaus AD, Rimmelzwaan GF (2005) Functional profile of human influenza virus-specific cytotoxic T lymphocyte activity is influenced by interleukin-2 concentration and epitope specificity. *Clin Exp Immunol* 142: 45–52.
- Boon AC, de Mutsert G, van Baarle D, Smith DJ, Lapedes AS, et al. (2004) Recognition of homo- and heterosubtypic variants of influenza A viruses by human CD8+ T lymphocytes. *J Immunol* 172: 2453–2460.
- Germeau C, Ma W, Schiavetti F, Lurquin C, Henry E, et al. (2005) High frequency of antitumor T cells in the blood of melanoma patients before and after vaccination with tumor antigens. *J Exp Med* 201: 241–248.
- Lau LL, Jamieson BD, Somasundaram T, Ahmed R (1994) Cytotoxic T-cell memory without antigen. *Nature* 369: 648–652.
- Hammarlund E, Lewis MW, Hansen SG, Strelow LI, Nelson JA, et al. (2003) Duration of antiviral immunity after smallpox vaccination. *Nat Med*.
- Combadiere B, Boissonnas A, Carcelain G, Lefranc E, Samri A, et al. (2004) Distinct time effects of vaccination on long-term proliferative and IFN- γ -producing T cell memory to smallpox in humans. *J Exp Med* 199: 1585–1593.
- Demkowicz WE, Jr., Littau RA, Wang J, Ennis FA (1996) Human cytotoxic T-cell memory: long-lived responses to vaccinia virus. *J Virol* 70: 2627–2631.
- Banchereau J, Palucka AK (2005) Dendritic cells as therapeutic vaccines against cancer. *Nat Rev Immunol* 5: 296–306.
- Banchereau J, Steinman RM (1998) Dendritic cells and the control of immunity. *Nature* 392: 245–252.
- Kupper TS, Fuhlbrigge RC (2004) Immune surveillance in the skin: mechanisms and clinical consequences. *Nat Rev Immunol* 4: 211–222.
- Klechevsky E, Morita R, Liu M, Cao Y, Coquery S, et al. (2008) Functional specializations of human epidermal Langerhans cells and CD14+ dermal dendritic cells. *Immunity* 29: 497–510.
- Combadiere B, Mahe B (2008) Particle-based vaccines for transcutaneous vaccination. *Comp Immunol Microbiol Infect Dis* 31: 293–315.
- Belyakov IM, Hammond SA, Ahlers JD, Glenn GM, Berzofsky JA (2004) Transcutaneous immunization induces mucosal CTLs and protective immunity by migration of primed skin dendritic cells. *J Clin Invest* 113: 998–1007.
- Berzofsky JA, Ahlers JD, Belyakov IM (2001) Strategies for designing and optimizing next generation vaccines. *Nat Rev Immunol* 1: 209–219.
- Le Borgne M, Etchart N, Goubier A, Lira SA, Sirard JC, et al. (2006) Dendritic cells rapidly recruited into epithelial tissues via CCR6/CCL20 are responsible for CD8+ T cell crosspriming in vivo. *Immunity* 24: 191–201.
- Bos JD, Kapsenberg ML (1993) The skin immune system: progress in cutaneous biology. *Immunol Today* 14: 75–78.
- Glenn GM, Rao M, Matyas GR, Alving CR (1998) Skin immunization made possible by cholera toxin. *Nature* 391: 851.
- Glenn GM, Scharton-Kersten T, Vassell R, Mallett CP, Hale TL, et al. (1998) Transcutaneous immunization with cholera toxin protects mice against lethal mucosal toxin challenge. *J Immunol* 161: 3211–3214.
- Glenn GM, Taylor DN, Li X, Frankel S, Montemarano A, et al. (2000) Transcutaneous immunization: a human vaccine delivery strategy using a patch. *Nat Med* 6: 1403–1406.
- Hammond SA, Walwender D, Alving CR, Glenn GM (2001) Transcutaneous immunization: T cell responses and boosting of existing immunity. *Vaccine* 19: 2701–2707.
- Matyas GR, Friedlander AM, Glenn GM, Little S, Yu J, et al. (2004) Needle-free skin patch vaccination method for anthrax. *Infect Immun* 72: 1181–1183.
- Glenn GM, Flyer DC, Ellingsworth LR, Frech SA, Frerichs DM, et al. (2007) Transcutaneous immunization with heat-labile enterotoxin: development of a needle-free vaccine patch. *Expert Rev Vaccines* 6: 809–819.
- Frerichs DM, Ellingsworth LR, Frech SA, Flyer DC, Villar CP, et al. (2008) Controlled, single-step, stratum corneum disruption as a pretreatment for immunization via a patch. *Vaccine* 26: 2782–2787.
- Vogt A, Combadiere B, Hadam S, Stieler KM, Lademann J, et al. (2006) 40 nm, but not 750 or 1,500 nm, nanoparticles enter epidermal CD1a+ cells after transcutaneous application on human skin. *J Invest Dermatol* 126: 1316–1322.
- Vogt A, Mahe B, Costagliola D, Bonduelle O, Hadam S, et al. (2008) Transcutaneous anti-influenza vaccination promotes both CD4 and CD8 T cell immune responses in humans. *J Immunol* 180: 1482–1489.
- Vogt A, Mandt N, Lademann J, Schaefer H, Blume-Peytavi U (2005) Follicular targeting—a promising tool in selective dermatotherapy. *J Invest Dermatol Symp Proc* 10: 252–255.
- Doherty PC, Turner SJ, Webby RG, Thomas PG (2006) Influenza and the challenge for immunology. *Nat Immunol* 7: 449–455.
- Greenbaum JA, Kotturi MF, Kim Y, Oseroff C, Vaughan K, et al. (2009) Pre-existing immunity against swine-origin H1N1 influenza viruses in the general human population. *Proc Natl Acad Sci U S A* 106: 20365–20370.
- Kohlmeier JE, Woodland DL (2009) Immunity to respiratory viruses. *Annu Rev Immunol* 27: 61–82.
- Cerwenka A, Morgan TM, Dutton RW (1999) Naive, effector, and memory CD8 T cells in protection against pulmonary influenza virus infection: homing properties rather than initial frequencies are crucial. *J Immunol* 163: 5535–5543.
- Kohlmeier JE, Rumsey LM, Chan MA, Benedict SH (2003) The outcome of T-cell costimulation through intercellular adhesion molecule-1 differs from costimulation through leucocyte function-associated antigen-1. *Immunology* 108: 152–157.
- Walker BD, Burton DR (2008) Toward an AIDS vaccine. *Science* 320: 760–764.
- Goulder PJ, Watkins DI (2008) Impact of MHC class I diversity on immune control of immunodeficiency virus replication. *Nat Rev Immunol* 8: 619–630.
- Watkins DI, Burton DR, Kallas EG, Moore JP, Koff WC (2008) Nonhuman primate models and the failure of the Merck HIV-1 vaccine in humans. *Nat Med* 14: 617–621.
- Hansen SG, Vieville C, Whizin N, Coyne-Johnson L, Siess DC, et al. (2009) Effector memory T cell responses are associated with protection of rhesus monkeys from mucosal simian immunodeficiency virus challenge. *Nat Med* 15: 293–299.
- Mahe B, Vogt A, Liard C, Duffy D, Abadie V, et al. (2008) Nanoparticle-Based Targeting of Vaccine Compounds to Skin Antigen-Presenting Cells By Hair Follicles and their Transport in Mice. *J Invest Dermatol*.
- Pulendran B, Smith JL, Caspari G, Brasel K, Pettit D, et al. (1999) Distinct dendritic cell subsets differentially regulate the class of immune response in vivo. *Proc Natl Acad Sci U S A* 96: 1036–1041.
- Dudziak D, Kamphorst AO, Heidkamp GF, Buchholz VR, Trumpfheller C, et al. (2007) Differential antigen processing by dendritic cell subsets in vivo. *Science* 315: 107–111.
- Stoitzner P, Tripp CH, Douillard P, Sacland S, Romani N (2005) Migratory Langerhans cells in mouse lymph nodes in steady state and inflammation. *J Invest Dermatol* 125: 116–125.
- Stoitzner P, Tripp CH, Eberhart A, Price KM, Jung JY, et al. (2006) Langerhans cells cross-present antigen derived from skin. *Proc Natl Acad Sci U S A* 103: 7783–7788.
- Kissenpfennig A, Malissen B (2006) Langerhans cells—revisiting the paradigm using genetically engineered mice. *Trends Immunol* 27: 132–139.
- Matsuo M, Nagata Y, Sato E, Atanackovic D, Valmori D, et al. (2004) IFN- γ -langerhans enables cross-presentation of exogenous protein antigen in human Langerhans cells by potentiating maturation. *Proc Natl Acad Sci U S A* 101: 14467–14472.
- Ratzinger G, Baggers J, de Cos MA, Yuan J, Dao T, et al. (2004) Mature human Langerhans cells derived from CD34+ hematopoietic progenitors stimulate greater cytolytic T lymphocyte activity in the absence of bioactive IL-12p70, by either single peptide presentation or cross-priming, than do dermal-interstitial or monocyte-derived dendritic cells. *J Immunol* 173: 2780–2791.

56. Belyakov IM, Ahlers JD, Berzofsky JA (2004) Mucosal AIDS vaccines: current status and future directions. *Expert Rev Vaccines* 3: S65–73.
57. Otberg N, Richter H, Schaefer H, Blume-Peytavi U, Sterry W, et al. (2004) Variations of hair follicle size and distribution in different body sites. *J Invest Dermatol* 122: 14–19.
58. Mangelsdorf S, Otberg N, Maibach HI, Sinkgraven R, Sterry W, et al. (2006) Ethnic variation in vellus hair follicle size and distribution. *Skin Pharmacol Physiol* 19: 159–167.
59. Etchart N, Hennino A, Friede M, Dahel K, Dupouy M, et al. (2007) Safety and efficacy of transcutaneous vaccination using a patch with the live-attenuated measles vaccine in humans. *Vaccine* 25: 6891–6899.
60. Yagi H, Hashizume H, Horibe T, Yoshinari Y, Hata M, et al. (2006) Induction of therapeutically relevant cytotoxic T lymphocytes in humans by percutaneous peptide immunization. *Cancer Res* 66: 10136–10144.
61. Doherty PC, Topham DJ, Tripp RA, Cardin RD, Brooks JW, et al. (1997) Effector CD4+ and CD8+ T-cell mechanisms in the control of respiratory virus infections. *Immunol Rev* 159: 105–117.
62. Hou S, Hyland L, Ryan KW, Portner A, Doherty PC (1994) Virus-specific CD8+ T-cell memory determined by clonal burst size. *Nature* 369: 652–654.
63. Haanen JB, Wolkers MC, Kruisbeek AM, Schumacher TN (1999) Selective expansion of cross-reactive CD8(+) memory T cells by viral variants. *J Exp Med* 190: 1319–1328.
64. Watabe S, Xin KQ, Ihata A, Liu LJ, Honsho A, et al. (2001) Protection against influenza virus challenge by topical application of influenza DNA vaccine. *Vaccine* 19: 4434–4444.
65. Graham MB, Braciale TJ (1997) Resistance to and recovery from lethal influenza virus infection in B lymphocyte-deficient mice. *J Exp Med* 186: 2063–2068.
66. Epstein SL, Lo CY, Mispion JA, Bennink JR (1998) Mechanism of protective immunity against influenza virus infection in mice without antibodies. *J Immunol* 160: 322–327.
67. Pala P, Townsend AR, Askonas BA (1986) Viral recognition by influenza A virus cross-reactive cytotoxic T (Tc) cells: the proportion of Tc cells that recognize nucleoprotein varies between individual mice. *Eur J Immunol* 16: 193–198.
68. Townsend ARM, Rothbard J, Gotch FM, Bahadur G, Wraith D, et al. (1986) The epitopes of influenza nucleoprotein recognized by cytotoxic T lymphocytes can be defined with short synthetic peptides. *Cell* 44: 959–968.
69. Belz GT, Doherty PC (2001) Virus-specific and bystander CD8+ T-cell proliferation in the acute and persistent phases of a gammaherpesvirus infection. *J Virol* 75: 4435–4438.
70. Belz GT, Xie W, Doherty PC (2001) Diversity of epitope and cytokine profiles for primary and secondary influenza A virus-specific CD8+ T cell responses. *J Immunol* 166: 4627–4633.
71. Belz GT, Xie W, Altman JD, Doherty PC (2000) A previously unrecognized H-2D(b)-restricted peptide prominent in the primary influenza A virus-specific CD8(+) T-cell response is much less apparent following secondary challenge. *J Virol* 74: 3486–3493.
72. Tamura M, Kuwano K, Kurane I, Ennis FA (1998) Definition of amino acid residues on the epitope responsible for recognition by influenza A virus H1-specific, H2-specific, and H1- and H2-cross-reactive murine cytotoxic T-lymphocyte clones. *J Virol* 72: 9404–9406.
73. Kuwano K, Tamura M, Ennis FA (1990) Cross-reactive protection against influenza A virus infections by an NS1-specific CTL clone. *Virology* 178: 174–179.
74. Kuwano K, Scott M, Young JF, Ennis FA (1989) Active immunization against virus infections due to antigenic drift by induction of crossreactive cytotoxic T lymphocytes. *J Exp Med* 169: 1361–1371.
75. Kuwano K, Braciale TJ, Ennis FA (1989) Cytotoxic T lymphocytes recognize a cross-reactive epitope on the transmembrane region of influenza H1 and H2 hemagglutinins. *Viral Immunol* 2: 163–173.
76. Kuwano K, Scott M, Young JF, Ennis FA (1988) HA2 subunit of influenza A H1 and H2 subtype viruses induces a protective cross-reactive cytotoxic T lymphocyte response. *J Immunol* 140: 1264–1268.
77. Murasko DM, Bernstein ED, Gardner EM, Gross P, Munk G, et al. (2002) Role of humoral and cell-mediated immunity in protection from influenza disease after immunization of healthy elderly. *Exp Gerontol* 37: 427–439.
78. Bernstein E, Kaye D, Abrutyn E, Gross P, Dorfman M, et al. (1999) Immune response to influenza vaccination in a large healthy elderly population. *Vaccine* 17: 82–94.
79. Makedonas G, Betts MR (2006) Polyfunctional analysis of human t cell responses: importance in vaccine immunogenicity and natural infection. *Springer Semin Immunopathol* 28: 209–219.
80. Harari A, Dutoit V, Cellerai C, Bart PA, Du Pasquier RA, et al. (2006) Functional signatures of protective antiviral T-cell immunity in human virus infections. *Immunol Rev* 211: 236–254.
81. Fan H, Lin Q, Morrissey GR, Khavari PA (1999) Immunization via hair follicles by topical application of naked DNA to normal skin. *Nat Biotechnol* 17: 870–872.
82. Lademann J, Otberg N, Jacobi U, Hoffman RM, Blume-Peytavi U (2005) Follicular penetration and targeting. *J Invest Dermatol Symp Proc* 10: 301–303.
83. Lademann J, Richter H, Schaefer UF, Blume-Peytavi U, Teichmann A, et al. (2006) Hair follicles - a long-term reservoir for drug delivery. *Skin Pharmacol Physiol* 19: 232–236.
84. Naito S, Maceyama J, Mizukami T, Takahashi M, Hamaguchi I, et al. (2007) Transcutaneous immunization by merely prolonging the duration of antigen presence on the skin of mice induces a potent antigen-specific antibody response even in the absence of an adjuvant. *Vaccine* 25: 8762–8770.
85. Glenn GM, Kenney RT (2006) Mass vaccination: solutions in the skin. *Curr Top Microbiol Immunol* 304: 247–268.
86. Kenney RT, Frech SA, Muenz LR, Villar CP, Glenn GM (2004) Dose sparing with intradermal injection of influenza vaccine. *N Engl J Med* 351: 2295–2301.
87. Halperin SA, Smith B, Clarke K, Treanor J, Mabrouk T, et al. (2005) Phase I, randomized, controlled trial to study the reactogenicity and immunogenicity of a nasal, inactivated trivalent influenza virus vaccine in healthy adults. *Hum Vaccin* 1: 37–42.
88. Draenert R, Altfeld M, Brander C, Basgoz N, Corcoran C, et al. (2003) Comparison of overlapping peptide sets for detection of antiviral CD8 and CD4 T cell responses. *J Immunol Methods* 275: 19–29.
89. Beavis AJ, Kalejta RF (2001) Simultaneous analysis of the cyan, green, and yellow fluorescent proteins. *Curr Protoc Cytom Chapter 1: Unit 1* 16.
90. Appay V, Rowland-Jones SL (2002) The assessment of antigen-specific CD8+ T cells through the combination of MHC class I tetramer and intracellular staining. *J Immunol Methods* 268: 9–19.
91. Appay V, Nixon DF, Donahoe SM, Gillespie GM, Dong T, et al. (2000) HIV-specific CD8(+) T cells produce antiviral cytokines but are impaired in cytolytic function. *J Exp Med* 192: 63–75.