



Lack of efflux mediated quinolone resistance in *Salmonella enterica* serovars Typhi and Paratyphi A.

Sylvie Baucheron, Isabelle Monchaux, Simon Le Hello, François-Xavier Weill,
Axel Cloeckaert

► To cite this version:

Sylvie Baucheron, Isabelle Monchaux, Simon Le Hello, François-Xavier Weill, Axel Cloeckaert. Lack of efflux mediated quinolone resistance in *Salmonella enterica* serovars Typhi and Paratyphi A.. *Frontiers in Microbiology*, 2013, 5, pp.12. 10.3389/fmicb.2014.00012 . pasteur-01104960

HAL Id: pasteur-01104960

<https://pasteur.hal.science/pasteur-01104960>

Submitted on 19 Jan 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Lack of efflux mediated quinolone resistance in *Salmonella enterica* serovars Typhi and Paratyphi A

Sylvie Baucheron^{1,2*}, Isabelle Monchaux^{1,2}, Simon Le Hello³, François-Xavier Weill³ and Axel Cloeckaert^{1,2*}

¹ UMR1282 Infectiologie et Santé Publique, Institut National de la Recherche Agronomique, Nouzilly, France

² UMR1282 Infectiologie et Santé Publique, Université François Rabelais de Tours, Tours, France

³ Institut Pasteur, Unité des Bactéries Pathogènes Entériques, Centre National de Référence des *Escherichia coli*, *Shigella* et *Salmonella*, Paris, France

Edited by:

Kunihiko Nishino, Osaka University, Japan

Reviewed by:

Kunihiko Nishino, Osaka University, Japan

Peter Heisig, University of Hamburg, Germany

*Correspondence:

Sylvie Baucheron and Axel Cloeckaert, Unité Infectiologie et Santé Publique site 213, Institut National de la Recherche Agronomique, 37380 Nouzilly, France
e-mail: sylvie.baucheron@tours.inra.fr;
axel.cloeckaert@tours.inra.fr

Salmonella enterica serovars Typhi and Paratyphi A isolates from human patients in France displaying different levels of resistance to quinolones or fluoroquinolones were studied for resistance mechanisms to these antimicrobial agents. All resistant isolates carried either single or multiple target gene mutations (i.e., in *gyrA*, *gyrB*, or *parC*) correlating with the resistance levels observed. Active efflux, through upregulation of multipartite efflux systems, has also been previously reported as contributing mechanism for other serovars. Therefore, we investigated also the occurrence of non-target gene mutations in regulatory regions affecting efflux pump expression. However, no mutation was detected in these regions in both Typhi and Paratyphi isolates of this study. Besides, no overexpression of the major efflux systems was observed for these isolates. Nevertheless, a large deletion of 2334 bp was identified in the *acrS-acrE* region of all *S. Typhi* strains but which did not affect the resistance phenotype. As being specific to *S. Typhi*, this deletion could be used for specific molecular detection purposes. In conclusion, the different levels of quinolone or FQ resistance in both *S. Typhi* and *S. Paratyphi A* seem to rely only on target modifications.

Keywords: *Salmonella*, ciprofloxacin, transcriptional regulatory genes, *acrS*, efflux pumps

INTRODUCTION

Enteric fever caused by the human-adapted pathogens *Salmonella enterica* serovars Typhi (*S. Typhi*) and Paratyphi A (*S. Paratyphi A*), B, and C, remains a major health problem (Crump and Mintz, 2010). A global epidemiologic study estimated that during the year 2000 typhoid fever caused 21.7 million illnesses and 21,700 deaths and paratyphoid fever caused 5.4 million illnesses (Crump et al., 2004). During the past decade *S. Paratyphi A* was responsible for a growing proportion of enteric fever in Asia (Ochiai et al., 2005; Crump and Mintz, 2010). Enteric fever being associated with poor sanitation and unsafe food and water, it particularly affects children and adolescents in developing countries of Asia, Africa and Latin America (Crump et al., 2004; Bhan et al., 2005; Crump and Mintz, 2010). In developed countries, patients are most often ill-returned travellers or migrant workers (Bhan et al., 2005; Connor and Schwartz, 2005; Hassing et al., 2013).

To treat these infections, fluoroquinolones (FQ) and third-generation cephalosporins have been considered as first-line drugs, owing to the resistance to ampicillin, chloramphenicol, and trimethoprim/sulfamethoxazole that appeared during the 1980s (Hassing et al., 2011, 2013). Multidrug resistance (MDR) in *S. Typhi* is encoded mainly by resistance genes carried by large conjugative plasmids and has been reported worldwide (Le et al., 2007). As a consequence of a widespread FQ usage, *S. Typhi* and *S. Paratyphi A* isolates resistant to nalidixic acid (NAL^R, minimum inhibitory concentration [MIC] > 16 mg/L) and with decreased susceptibility to ciprofloxacin (CIP^{DS}, MIC 0.125–1.0 mg/L) have also emerged. Such NAL^R-CIP^{DS} *S. Typhi*

and *S. Paratyphi A* have been isolated in endemic areas and also in developed countries (Roumagnac et al., 2006; Le et al., 2007; Gaborieau et al., 2010; Accou-Demartin et al., 2011; Hassing et al., 2011, 2013).

Resistance to quinolones in *Salmonella* spp. is mostly attributed to point mutations in the quinolone resistance-determining regions (QRDRs) of the target genes *gyrA*, *gyrB*, *parC*, and *parE* (Cloeckaert and Chaslus-Dancla, 2001; Piddock, 2002; Velge et al., 2005; Giraud et al., 2006). For the *gyrA* gene, coding the A subunit of DNA gyrase, a single mutation resulting in an amino acid substitution at the position 83 (Serine to Phenylalanine or to Tyrosine) or at the position 87 (Aspartic acid to Asparagine or Glycine) has been the most frequently described in NAL^R-CIP^{DS} *S. Typhi* and *S. Paratyphi A* isolates (Bhan et al., 2005; Renuka et al., 2005; Le et al., 2007; Gaborieau et al., 2010; Hassing et al., 2011). A second mutation leading to the amino acid change at the position 80 (Serine to Isoleucine or to Arginine) of the ParC subunit of topoisomerase IV was described to increase the CIP MIC (≥ 0.5 mg/L) in *S. Typhi* and *S. Paratyphi A* human isolates from India (Gaïnd et al., 2006). Whereas three mutations, i.e., a double mutation in *gyrA* at both codons 83 and 87 and one mutation in *parC*, were shown to confer CIP resistance (MIC > 1 mg/L) in *S. Typhi* and *S. Paratyphi A* human isolates from India or from Taiwan (Gaïnd et al., 2006; Lee et al., 2013).

Moreover, the varying levels of CIP resistance observed in *S. Typhi* and *S. Paratyphi A* isolates with only a single *gyrA* mutation suggest that other mechanisms could be involved in quinolone resistance in this serovar (Renuka et al., 2005).

Resistance to FQ in *S. Typhimurium* has also been attributed to active efflux mechanism, due to overproduction of the AcrAB-TolC efflux system (Giraud et al., 2000, 2006; Cloeckaert and Chaslus-Dancla, 2001; Piddock, 2006). We have previously reported the contribution of the AcrAB-TolC efflux system in resistance to FQ in several MDR epidemic clones of *S. Typhimurium*, such as *S. Typhimurium* of phage types DT204 or DT104 (Baucheron et al., 2002, 2004a,b). Among the chromosomal loci affecting AcrAB-TolC expression, the *ramRA* locus appears to be the most important in *Salmonella* spp. (Abouzeed et al., 2008; Kehrenberg et al., 2009). *ramR* encodes a repressor protein (RamR) belonging to the TetR family of repressor proteins, and has been shown to be the local repressor protein of *ramA* transcription (Abouzeed et al., 2008; Baucheron et al., 2012); while *ramA* encodes a transcriptional activator protein (RamA) belonging to the AraC/XylS family of regulatory proteins (Nikaido et al., 2008; Bailey et al., 2010). The latter is involved in upregulating expression of the AcrAB-TolC system (Nikaido et al., 2008; Bailey et al., 2010). Several mutations in *ramR* or its binding site upstream of *ramA*, affecting expression of this efflux system, have been detected in clinical isolates of serovar Typhimurium or Kentucky and of minor serovars Hadar, Infantis, Livingstone, or Schwarzengrund (Abouzeed et al., 2008; Kehrenberg et al., 2009;

Hentschke et al., 2010; Akiyama and Khan, 2012; Baucheron et al., 2013).

In the present study, we have characterized mechanisms involved in resistance to quinolones or fluoroquinolones in 21 *S. Typhi* and *S. Paratyphi A* strains displaying different levels of resistance to these drugs and isolated from patients in France during the period 1997–2008. For a subset of strains, with suspected increased efflux activity, we investigated the occurrence of mutations in the global *ram*, *sox* and *mar* regulatory loci of AcrAB-TolC, and in the local *acrR* and *acrS* repressor genes of the AcrAB and AcrEF pumps, respectively (Abouzeed et al., 2008; Kehrenberg et al., 2009).

MATERIALS AND METHODS

BACTERIAL STRAINS

The twenty one strains including 16 *S. Typhi* and 5 *S. Paratyphi A* selected for this study were collected by the French National Reference Center for *Salmonella*, Institut Pasteur, Paris, France. They were isolated in France from travellers or migrants between 1997 and 2008 (Table 1). The selection was made to obtain diversity in terms of geographic origin, year of isolation, genetic lineages (haplotype for *S. Typhi*; Roumagnac et al., 2006), and phenotype of resistance to quinolones (Table 1).

Table 1 | *Salmonella enterica* serovars Typhi and Paratyphi A strains analyzed in this study.

Strain	Country	Year of isolation	Haplo type	Antimicrobial resistance pattern	MIC (mg/L)		Substitution(s) in the QRDR of:				AcrA production ratio*	
					NAL	CIP	GyrA	GyrB	ParC	ParE		
SALMONELLA TYPHI												
06-423	India	2006	ND	Pansusceptible	4	0.015	WT	WT	WT	WT	1	
06-426	India	2006	ND	CIP ^{DS}	16	0.125	WT	S464Y	WT	WT	1	
02-1180	India	2002	H45	NALCIP ^{DS}	64	0.125	D87G	WT	WT	WT	1	
05-3275	Morocco	2005	H6	NALCIP ^{DS}	64	0.125	D87N	WT	WT	WT	1	
4(02)MB	Vietnam	1997	H58	ASCSTmpTeNAL	128	0.03	S83Y	WT	WT	WT	0.5	
222(97)MN	Vietnam	1996	ND	ASCSTmpTeNALCIP ^{DS}	128	0.125	S83F	WT	WT	WT	0.5	
43(97)MN	Vietnam	1996	H63	ASCSTmpTeNALCIP ^{DS}	128	0.125	S83F	WT	WT	WT	0.5	
98-3139	Mexico	1998	H50	NALCIP ^{DS}	128	0.125	S83F	WT	WT	WT	0.5	
02-7744	India	2002	H52	NALCIP ^{DS}	128	0.125	S83F	WT	WT	WT	0.5	
226(97)MN	Vietnam	1996	H61	ASCSTmpTeNALCIP ^{DS}	128	0.25	S83F	WT	WT	WT	0.5	
97-2307	India	1997	H63	NALCIP ^{DS}	256	0.125	S83F	WT	WT	WT	0.5	
318(98)MB	Vietnam	1998	H58	ASCSTmpTeNALCIP ^{DS}	512	0.25	S83Y	WT	WT	WT	1	
39(98)MN	Vietnam	1998	H58	ASCSTmpTeNALCIP ^{DS}	512	0.25	S83F	WT	WT	WT	1	
4(02)MN	Vietnam	2000	H58	ASCSTmpTeNALCIP ^{DS}	1024	0.25	S83F	WT	WT	D420N	1	
5(04)MN	Vietnam	2004	ND	NALCIP ^{DS}	1024	0.25	S83F	WT	WT	D420N	1	
04-2176	India	2004	H58	SSpSulTmpTeNALCIP	1024	8	S83F; D87N	WT	S80I	WT	1	
SALMONELLA PARATYPHI A												
08-8903	Senegal	2008		Pansusceptible	8	0.030	WT	WT	WT	WT	2	
07-6329	Burkina Faso	2007		CIP ^{DS}	16	0.25	WT	S464F	WT	WT	2	
05-208	India	2005		NALCIP ^{DS}	256	0.50	S83F	WT	WT	WT	2	
08-4271	Guinea Bissau	2008		NACIP ^{DS}	1024	1	S83F	WT	WT	WT	2	
08-2580	India	2008		NALCIP ^{DS}	1024	1	S83F	WT	WT	WT	3	

*AcrA expression was measured by dot blotting with an anti-AcrA polyclonal antibody.

ND, not determined.

WT, wild type.

A, amoxicillin; S, streptomycin; Sp, spectinomycin; C, chloramphenicol; Sul, sulfamethoxazole; Tmp, trimethoprim; Te, tetracycline; NAL, nalidixic acid; CIP, ciprofloxacin; CIP^{DS}, decreased susceptibility to ciprofloxacin.

D, aspartic acid; F, phenylalanine; G, glycine; I, isoleucine; N, asparagine; S, serine; Y, tyrosine.

ANTIMICROBIAL SUSCEPTIBILITY TESTING

Antimicrobial susceptibility was investigated by the standard disk diffusion method according to the recommendations of the Antibigram Committee of the French Society for Microbiology (CA-SFM) (www.sfm-microbiologie.org/). The MICs of NAL and CIP were determined by the standard agar doubling dilution method as described previously (Baucheron et al., 2002). The NAL^R isolates were defined as having a MIC > 16 mg/L. The CIP^R isolates were defined as having a MIC > 1 mg/L and CIP^{DS} isolates as having a MIC comprised between 0.125 and 1.0 mg/L (Accou-Demartin et al., 2011; Hassing et al., 2013). MICs of these antibiotics were also determined in the presence of the efflux pump inhibitor Phe-Arg- β -naphthylamide (PA β N, Sigma) at the following concentrations: 10, 20, 30, 40, 50, and 60 mg/L.

ASSESSMENT OF TARGET-AFFECTING MECHANISMS

Mutations in the QRDRs of *gyrA*, *gyrB*, *parC*, and *parE* genes were detected as described previously (Le et al., 2007; Song et al., 2010; Accou-Demartin et al., 2011).

The search of plasmid-mediated quinolone resistances genes, *qnrA*, *qnrB*, *qnrS*, *qnrD*, *aac(6')*-Ib-cr, and *qepA* was performed as described previously (Accou-Demartin et al., 2011).

ASSESSMENT OF EFFLUX MECHANISMS

Efflux pump production was assessed by dot blot using an anti-AcrA polyclonal antibody as described previously (Abouzeed et al., 2008). Occurrence of mutations affecting *acrAB*, *acrEF*, and *tolC* expression was determined by PCR and sequencing the regulatory regions *ramR-ramA*, *acrR-acrA*, *marC-marO-marR-marA*, *soxS-soxR*, and *acrS-acrE* using primers listed in (Table 2).

Table 2 | Primers used for PCRs.

Primer used and target region	Primer	Nucleotide position relative to the LT2 strain genome sequence *	Oligonucleotide sequence(s) (5'–3')	Size (bp)	Annealing temp (°C)	Reference
DETECTION OF MUTATIONS						
<i>ramR-ramA</i>	ram5	638085	TCGGTAAAAGGCAGTTCAG	958	60	Baucheron et al., 2013
	ramA6	639042	GTCGATAACCTGAGCGGAAA			
<i>acrR-acrA</i>	acrR1	533463	CAGTGGTTCCGTTTTAGTG	992	58	Olliver et al., 2005
	acrR2	534454	ACAGAATAGCGACACAGAAA			
<i>marC-marO-marR-marA</i>	marR1	1597459	CAGTGTTGCGTCTGGACATC	787	60	Baucheron et al., 2013
	marR2	1598245	GCTAACGGGAGCAGTACGAC			
<i>soxS-soxR</i>	sox1	4503970	CTACAGGCGGTGACGGTAAT	915	60	Baucheron et al., 2013
	sox2	4504884	CGGCGCTTTAGTTTTAGGTG			
<i>acrS-acrE</i>	acrS3	3559106	AAAACGAACGGGAACGTATG	2874 ***	58	This study
	acrS4	3561978	ACAAACATACCGGGAAGCAG			
qRT-PCR						
<i>gmk</i>	gmk-f	3933294	TTGGCAGGGAGGCGTTT	62	60	Baucheron et al., 2012
	gmk-r	3933355	GCGCGAAGTGCCGTAGTAAT			
<i>gyrB</i>	gyrB-f	4040275	TCTCCTCACAGACCAAAGATAAGCT	81	60	Baucheron et al., 2012
	gyrB-r	4040195	CGCTCAGCAGTTCGTTTCATC			
<i>rrs</i>	rrs-f	NA**	CCAGCAGCCGCGGTAAT	57	60	Baucheron et al., 2012
	rrs-r	NA**	TTTACGCCCAGTAATTCGATT			
<i>ramA</i>	ramA-f	639180	GCGTGAACGGAAGCTAAAAC	167	60	Baucheron et al., 2012
	ramA-r	639346	GGCCATGCTTTTCTTTACGA			
<i>ramR</i>	ramR-f	638623	TAACGCAGGTGTTGCAGAAG	192	64	Baucheron et al., 2012
	ramR-r	638432	TGGTTCAGACCCCAACTGAT			
<i>acrA</i>	acrA-f	533120	GAAACCGCACGTATCAACCT	220	60	Baucheron et al., 2012
	acrA-r	532901	CCTGTTTCAGCGAACCATTT			
<i>acrB</i>	acrB-f	531348	TCGTGTTCTCGTGATGTACCT	68	66	Baucheron et al., 2012
	acrB-r	531281	AACCGCAATAGTCGGAATCAA			
<i>acrF</i>	acrF-f	3563042	GCTCTGTCGTCCATCTCAAAGA	70	66	This study
	acrF-r	3563111	CGCGCTACAACGTTATAGTTTTCA			
<i>tolC</i>	tolC-f	3349107	GCCCGTGCGCAATATGAT	67	60	Baucheron et al., 2012
	tolC-r	3349173	CCGCGTTATCCAGGTTGTTG			

*GenBank NC_0031971.

**NA: Not Applicable due to the number of copies of this gene in *Salmonella*.

***2874 bp for *S. Typhimurium* or *S. Paratyphi A* and 539 bp for *S. Typhi* (see Figure 1).

Transcription levels of efflux related genes *acrA*, *acrB*, *acrF*, *tolC*, *ramA*, and *ramR* were done by quantitative reverse transcription-PCR (qRT-PCR) as described previously (Baucheron et al., 2012; Giraud et al., 2013). Primers used for qRT-PCR are listed in (Table 2).

RESULTS AND DISCUSSION

RESISTANCE PHENOTYPES AND TARGET-AFFECTING MECHANISMS

The twenty one *S. Typhi* and *S. Paratyphi A* strains of this study were isolated in France but acquired abroad, mainly in Asia and Africa (Table 1). Among the *S. Typhi* strains, all but two were NAL^R (MIC > 16 mg/L). One of the two strains was pan-susceptible and the second was CIP^{DS} but only categorized as intermediate for NAL (NAL^I, MIC 16 mg/L). Of the 14 NAL^R strains, one was CIP^R, 12 were CIP^{DS} and one was CIP^S (MIC 0.03 mg/L). Eight NAL^R *S. Typhi* strains were also multidrug resistant. The majority of the NAL^R *S. Typhi* strains belonged to haplotype H58 which had emerged in Southern Asia during the mid 1990s (Roumagnac et al., 2006; Le et al., 2007). Among the *S. Paratyphi A* strains, all but two were NAL^R. One of the two strains was pansusceptible and the second was NAL^I-CIP^{DS}. The three others were NAL^R-CIP^{DS}.

As shown in Table 1, both NAL^I-CIP^{DS} had a mutation resulting in an amino acid substitution at position 464 of GyrB: serine to tyrosine for the *S. Typhi* isolate and serine to phenylalanine for the *S. Paratyphi A* isolate. The most frequent mechanism of resistance of NAL^R-CIP^{DS} ($n = 17$) and NAL^R-CIP^S ($n = 1$) strains, whatever the serovar, was a substitution at position 83 (serine to phenylalanine, $n = 12$, 66.6%) of GyrA. Other GyrA modifications were observed at position 83 (serine to tyrosine) in two isolates or at position 87 (aspartic acid to glycine, aspartic acid to asparagine) in one isolate for each. As described previously, a single substitution in GyrA was associated with resistance to

nalidixic acid and decreased susceptibility to CIP (Bhan et al., 2005; Le et al., 2007; Gaborieau et al., 2010; Hassing et al., 2011). One exception was the *S. Typhi* strain 4 (02)MB, which was NAL^R-CIP^S (and not CIP^{DS}) despite a mutation in *gyrA* resulting in substitution serine to tyrosine at position 83.

Additional substitutions were found in ParE of 2 NAL^R-CIP^{DS} *S. Typhi* strains that led to amino acid substitution aspartic acid to asparagine at position 420. In both cases, a 2-fold increase of NAL MICs was observed.

In the CIP^R *S. Typhi* isolate, three mutations leading to a double substitution in GyrA at positions 83 (serine to phenylalanine) and 87 (aspartic acid to asparagine) and one substitution at the position 80 of ParC (serine to isoleucine), as observed in previous studies (Renuka et al., 2005; Gaind et al., 2006; Lee et al., 2013).

The NAL^R and CIP^{DS} *S. Typhi* and *S. Paratyphi A* strains harboring a single substitution in GyrA showed various values for NAL (64–1024 mg/L) and CIP (0.03–0.5 mg/L) MICs which suggested the presence of other mechanisms of resistance. Since the plasmid-mediated quinolone resistance-conferring genes *qnrA*, *qnrB*, *qnrD* or *qnrS*, *qepA*, and *aac(6′)-Ib-cr* were not detected, we investigated the role of the AcrAB-TolC efflux system.

INVOLVEMENT OF EFFLUX

None of the *S. Typhi* strains showed significant AcrA overproduction by dot blot, but nevertheless all *S. Paratyphi A* isolates showed a 2 or 3-fold increased AcrA production relative to the susceptible *S. Typhi* isolate (Table 1). Thus, overproduction of AcrA seems not to be involved in CIP^{DS} isolates compared with the susceptible isolates of *S. Typhi* or *S. Paratyphi A*. In presence of the efflux pump inhibitor PAβN (20 or 40 mg/L), the CIP MICs similarly decreased (4 or 8-fold) in CIP^{DS} and in susceptible strains (Table 3 and data not shown), which is in accordance with previous studies on *S. Typhimurium* and corresponds to

Table 3 | Study of efflux in a subset of *Salmonella enterica* serovars Typhi and Paratyphi A strains.

Strain	Antimicrobial resistance pattern ^a	MIC (mg/L) ^b		Substitution(s) in the QRDR of ^c :				<i>acrSE</i> sequencing	Transcription level of:					
		NAL	CIP	GyrA	GyrB	ParC	ParE		<i>acrA</i>	<i>acrF</i>	<i>acrB</i>	<i>tolC</i>	<i>ramA</i>	<i>ramR</i>
SALMONELLA TYPHI														
06–423	Pansusceptible	4	0.015 [0.004]	WT	WT	WT	WT	Deletion ^d	1.0	1.0	1.0	1.0	1.0	1.0
02–1180	NALCIP ^{DS}	64	0.125 [0.015]	D87G	WT	WT	WT	Deletion ^d	1.5	0.7	0.5	0.8	0.9	1.4
05–3275	NALCIP ^{DS}	64	0.125 [0.030]	D87N	WT	WT	WT	Deletion ^d	0.5	1.4	0.2	0.5	0.3	0.7
97–2307	NALCIP ^{DS}	256	0.125 [0.030]	S83F	WT	WT	WT	Deletion ^d	1.7	0.8	0.8	0.7	1.9	2.2
04–2176	SSpSulTmptTeNALCIP	1024	8 [2]	S83F; A87N	WT	S80I	WT	Deletion ^d	1.5	0.8	1.3	0.9	1.2	2.2
SALMONELLA PARATYPHI A														
08–8903	Pansusceptible	8	0.030 [0.008]	WT	WT	WT	WT	WT	1.0	1.0	1.0	1.0	1.0	1.0
07–6329	CIP ^{DS}	16	0.25 [0.060]	WT	S464F	WT	WT	WT	1.6	1.4	1.0	1.0	1.0	1.2
05–208	NALCIP ^{DS}	256	0.50 [0.030]	S83F	WT	WT	WT	WT	1.3	0.9	1.4	1.0	1.1	1.0
08–4271	NALCIP ^{DS}	1024	1 [0.25]	S83F	WT	WT	WT	WT	1.2	1.1	0.7	1.4	1.3	1.3
08–2580	NALCIP ^{DS}	1024	1 [0.25]	S83F	WT	WT	WT	WT	2.0	1.6	1.4	1.4	2.0	1.3

^a S, streptomycin; Sp, spectinomycin; Sul, sulfamethoxazole; Tmp, trimethoprim; Te, tetracycline; NAL, nalidixic acid; CIP, ciprofloxacin; CIP^{DS}, decreased susceptibility to ciprofloxacin.

^b Values in brackets are MICs in the presence of the efflux pump inhibitor PAβN at 40 mg/L.

^c WT, wild type; D, aspartic acid; F, phenylalanine; G, glycine; I, isoleucine; N, asparagine; S, serine.

^d 2334 bp deleted.




FIGURE 1 | Deletion identified in the *acrSE* region relative to the genome sequence of *S. Typhi* strain CT18 (GenBank NC_003198.1).

The sequence of the *acrSE* region of *S. Typhi* strain CT18 is compared to those of *S. Paratyphi A* strain ATCC9150 (GenBank NC_006511.1) and

S. Typhi strain LT2 (GenBank NC_003197.1). Small black arrows indicate primers positions used for PCR to amplify and sequence the *acrSE* region. The 2334 bp chromosomal deletion was found in all *S. Typhi* strains studied.

a decrease of resistance level observed for *acrB* or *tolC* deletion mutants (Baucheron et al., 2002, 2004b).

Despite a lack of evidence of increased efflux in the resistance phenotype, we measured by qRT-PCR the transcription levels of efflux related genes *acrA*, *acrF*, *acrB*, *tolC*, *ramA*, and *ramR* in CIP^{DS} non-MDR strains and in the CIP^R strain. No differences were detected in the transcription levels of these genes, between susceptible, CIP^{DS} and CIP^R strains, whatever the serovar (Table 3). In addition, no mutations were detected in the regulatory regions of the AcrAB-TolC efflux system. However, during the screening of the regulatory regions, we identified a single large deletion of 2334 bp in the *acrS*-*acrE* region of all *S. Typhi* strains, including the susceptible one (Table 3). This deletion encompassed the *acrS* gene, that encodes a transcriptional repressor, and a large part of the *acrE* gene that encodes the AcrE periplasmic lipoprotein, which is homologous to AcrA (Olliver et al., 2005). This 2334 bp deletion was also observed in the *acrS*-*acrE* region of sequenced genomes of MDR *S. Typhi* CT18 strain (Parkhill et al., 2001) and pansusceptible TY2 strain (Deng et al., 2003) (Figure 1). Previously, it has been shown that *acrS* deletion in *S. Typhimurium* does not affect *acrEF* expression (Olliver et al., 2005). Similarly the natural *acrSE* deletion detected in *S. Typhi* had no impact on the *acrF* transcription level as observed in this study. To our knowledge, this is the first description of such a natural *acrS*-*acrE* chromosomal deletion and seems specific to *S. Typhi* since it was not detected in all currently sequenced genomes of the other serovars (not shown).

CONCLUSIONS

The main mechanisms involved in quinolone or FQ resistance in both *S. Typhi* and *S. Paratyphi A* are target modifications. In contrast to what is seen in enteric pathogenic serovars, such as Typhimurium or the emerging CIP^R Kentucky ST198 clone (Baucheron et al., 2013), increased efflux pump production-mediated mechanisms seem to be totally absent in both *S. Typhi* and *S. Paratyphi A*. The deletion identified in the *acrSEF* region, although not involved in the resistance phenotype, may be helpful for the specific detection of *S. Typhi*.

ACKNOWLEDGMENTS

We are grateful to Dr. Hong Le for providing some of the studied strains and to Laëticia Fabre for excellent technical assistance. We would like to thank all corresponding laboratories of the French National Reference Center for *E. coli*, *Shigella*, and *Salmonella*. The French National Reference Center for *E. coli*, *Shigella*, and *Salmonella* is funded by the Institut Pasteur and the Institut de Veille Sanitaire. The “Unité des Bactéries Pathogènes Entériques” belongs to the “Integrative Biology of Emerging Infectious Diseases” Laboratory of Excellence funded by the French Government “Investissement d’Avenir” programme (grant no. ANR-10-LABX-62-IBED).

REFERENCES

- Abouzeed, Y. M., Baucheron, S., and Cloeckert, A. (2008). *ramR* mutations involved in efflux-mediated multidrug resistance in *Salmonella enterica* serovar Typhimurium. *Antimicrob. Agents Chemother.* 52, 2428–2434. doi: 10.1128/AAC.00084-08
- Accou-Demartin, M., Gaborieau, V., Song, Y., Roumagnac, P., Marchou, B., Achtman, M., et al. (2011). *Salmonella enterica* Serotype Typhi with nonclassical quinolone resistance phenotype. *Emerg. Infect. Dis.* 17, 1091–1094. doi: 10.3201/eid1706.101242
- Akiyama, T., and Khan, A. A. (2012). Molecular characterization of strains of fluoroquinolone-resistant *Salmonella enterica* serovar Schwarzengrund carrying multidrug resistance isolated from imported foods. *J. Antimicrob. Chemother.* 67, 101–110. doi: 10.1093/jac/dkr414
- Bailey, A. M., Ivens, A., Kingsley, R., Cottell, J. L., Wain, J., and Piddock, L. J. V. (2010). RamA, a member of the AraC/XylS family, influences both virulence and efflux in *Salmonella enterica* serovar Typhimurium. *J. Bacteriol.* 192, 1607–1616. doi: 10.1128/JB.01517-09
- Baucheron, S., Chaslus-Dancla, E., and Cloeckert, A. (2004a). Role of TolC and *parC* mutation in high-level fluoroquinolone resistance in *Salmonella enterica* serotype Typhimurium DT204. *J. Antimicrob. Chemother.* 53, 657–659. doi: 10.1093/jac/dkh122
- Baucheron, S., Tyler, S., Boyd, D., Mulvey, M. R., Chaslus-Dancla, E., and Cloeckert A. (2004b). AcrAB-TolC directs efflux-mediated multidrug resistance in *Salmonella enterica* serovar Typhimurium DT104. *Antimicrob. Agents Chemother.* 48, 3729–3735. doi: 10.1128/AAC.48.10.3729-3735.2004
- Baucheron, S., Coste, F., Canepa, S., Maurel, M. C., Giraud, E., Culard, F., et al. (2012). Binding of the RamR repressor to wild-type and mutated promoters of the RamA gene involved in efflux-mediated multidrug resistance in *Salmonella enterica* serovar Typhimurium. *Antimicrob. Agents Chemother.* 56, 942–948. doi: 10.1128/AAC.05444-11

- Baucheron, S., Imberechts, H., Chaslus-Dancla, E., and Cloeckaert, A. (2002). The AcrB multidrug transporter plays a major role in high-level fluoroquinolone resistance in *Salmonella enterica* serovar Typhimurium phage type DT204. *Microb. Drug Resist.* 8, 281–289. doi: 10.1089/10766290260469543
- Baucheron, S., Le Hello, S., Doublet, B., Giraud, E., Weill, F. X., and Cloeckaert, A. (2013). *ramR* mutations affecting fluoroquinolone susceptibility in epidemic multidrug-resistant *Salmonella enterica* serovar Kentucky ST198. *Front. Microbiol.* 4:213. doi: 10.3389/fmicb.2013.00213
- Bhan, M. K., Bahl, R., and Bhatnagar, S. (2005). Typhoid and paratyphoid fever. *Lancet* 366, 749–762. doi: 10.1016/S0140-6736(05)67181-4
- Cloeckaert, A., and Chaslus-Dancla, E. (2001). Mechanisms of quinolone resistance in *Salmonella*. *Vet. Res.* 32, 291–300. doi: 10.1051/vetres:2001105
- Connor, B. A., and Schwartz, E. (2005). Typhoid and paratyphoid fever in travellers. *Lancet Infect. Dis.* 5, 623–628. doi: 10.1016/S1473-3099(05)70239-5
- Crump, J. A., Luby, S. P., and Mintz, E. D. (2004). The global burden of typhoid fever. *Bull. World Health Organ.* 82, 346–353.
- Crump, J. A., and Mintz, E. D. (2010). Global trends in typhoid and paratyphoid fever. *Clin. Infect. Dis.* 50, 241–246. doi: 10.1086/649541
- Deng, W., Liou, S. R., Plunkett, G. 3rd., Mayhew, G. F., Rose, D. J., Burland, V., et al. (2003). Comparative genomics of *Salmonella enterica* serovar Typhi strains Ty2 and CT18. *J. Bacteriol.* 185, 2330–2337. doi: 10.1128/JB.185.7.2330-2337.2003
- Gaborieau, V., Weill, F. X., and Marchou, B. (2010). A case of *Salmonella enterica* serovar Typhi with decreased susceptibility to ciprofloxacin. *Med. Mal. Infect.* 40, 691–695. doi: 10.1016/j.medmal.2010.06.009
- Gaind, R., Paglietti, B., Murgia, M., Dawar, R., Uzzau, S., Cappuccinelli, P., et al. (2006). Molecular characterization of ciprofloxacin-resistant *Salmonella enterica* serovar Typhi and Paratyphi A causing enteric fever in India. *J. Antimicrob. Chemother.* 58, 1139–1144. doi: 10.1093/jac/dkl391
- Giraud, E., Baucheron, S., and Cloeckaert, A. (2006). Resistance to fluoroquinolones in *Salmonella*: emerging mechanisms and resistance prevention strategies. *Microbes Infect.* 8, 1937–1944. doi: 10.1016/j.micinf.2005.12.025
- Giraud, E., Baucheron, S., Virlogeux-Payant, I., Nishino, K., and Cloeckaert, A. (2013). Effects of natural mutations in the *ramRA* locus on invasiveness of epidemic fluoroquinolone-resistant *Salmonella enterica* serovar Typhimurium isolates. *J. Infect. Dis.* 207, 794–802. doi: 10.1093/infdis/jis755
- Giraud, E., Cloeckaert, A., Kerboeuf, D., and Chaslus-Dancla, E. (2000). Evidence for active efflux as the primary mechanism of resistance to ciprofloxacin in *Salmonella enterica* serovar Typhimurium. *Antimicrob. Agents Chemother.* 44, 1223–1228. doi: 10.1128/AAC.44.5.1223-1228.2000
- Hassing, R. J., Goessens, W. H., Mevius, D. J., van Pelt, W., Mouton, J. W., Verbon, A., et al. (2013). Decreased ciprofloxacin susceptibility in *Salmonella* Typhi and Paratyphi infections in ill-returned travellers: the impact on clinical outcome and future treatment options. *Eur. J. Clin. Microbiol. Infect. Dis.* 32, 1295–1301. doi: 10.1007/s10096-013-1878-9
- Hassing, R. J., Menezes, G. A., van Pelt, W., Petit, P. L., van Genderen, P. J., and Goessens, W. H. (2011). Analysis of mechanisms involved in reduced susceptibility to ciprofloxacin in *Salmonella enterica* serotypes Typhi and Paratyphi A isolates from travellers to Southeast Asia. *Int. J. Antimicrob. Agents* 37, 240–243. doi: 10.1016/j.ijantimicag.2010.10.026
- Hentschke, M., Christner, M., Sobottka, I., Aepfelbacher, M., and Rohde, H. (2010). Combined *ramR* mutation and presence of a Tn1721-associated tet(A) variant in a clinical isolate of *Salmonella enterica* serovar Hadar resistant to tigecycline. *Antimicrob. Agents Chemother.* 54, 1319–1322. doi: 10.1128/AAC.00993-09
- Kehrenberg, C., Cloeckaert, A., Klein, G., and Schwarz, S. (2009). Decreased fluoroquinolone susceptibility in mutants of *Salmonella* serovars other than Typhimurium: detection of novel mutations involved in modulated expression of *ramA* and *saxS*. *J. Antimicrob. Chemother.* 64, 1175–1180. doi: 10.1093/jac/dkp347
- Le, T. A., Fabre, L., Roumagnac, P., Grimont, P. A., Scavizzi, M. R., and Weill, F. X. (2007). Clonal expansion and microevolution of quinolone-resistant *Salmonella enterica* serotype Typhi in Vietnam from 1996 to 2004. *J. Clin. Microbiol.* 45, 3485–3492. doi: 10.1128/JCM.00948-07
- Lee, C. J., Su, L. H., Huang, Y. C., and Chiu, C. H. (2013). First isolation of ciprofloxacin-resistant *Salmonella enterica* serovar Typhi in Taiwan. *J. Microbiol. Immunol. Infect.* 46, 469–473. doi: 10.1016/j.jmii.2013.01.002
- Nikaido, E., Yamaguchi, A., and Nishino, K. (2008). AcrAB multidrug efflux pump regulation in *Salmonella enterica* serovar Typhimurium by RamA in response to environmental signals. *J. Biol. Chem.* 283, 24245–24253. doi: 10.1074/jbc.M804544200
- Ochiai, R. L., Wang, X., von Seidlein, L., Yang, J., Bhutta, Z. A., Bhattacharya, S. K., et al. (2005). *Salmonella* paratyphi A rates, Asia. *Emerg. Infect. Dis.* 11, 1764–1766. doi: 10.3201/eid1111.050168
- Olliver, A., Valle, M., Chaslus-Dancla, E., and Cloeckaert, A. (2005). Overexpression of the multidrug efflux operon *acrEF* by insertional activation with IS1 or IS10 elements in *Salmonella enterica* serovar typhimurium DT204 *acrB* mutants selected with fluoroquinolones. *Antimicrob. Agents Chemother.* 49, 289–301. doi: 10.1128/AAC.49.1.289-301.2005
- Parkhill, J., Dougan, G., James, K. D., Thomson, N. R., Pickard, D., Wain, J., et al. (2001). Complete genome sequence of a multiple drug resistant *Salmonella enterica* serovar Typhi CT18. *Nature* 413, 848–852. doi: 10.1038/35101607
- Piddock, L. J. V. (2002). Fluoroquinolone resistance in *Salmonella* serovars isolated from humans and food animals. *FEMS Microbiol. Rev.* 26, 3–16. doi: 10.1016/S0168-6445(01)00076-6
- Piddock, L. J. V. (2006). Clinically relevant chromosomally encoded multidrug resistance efflux pumps in bacteria. *Clin. Microbiol. Rev.* 19, 382–402. doi: 10.1128/CMR.19.2.382-402.2006
- Renuka, K., Sood, S., Das, B. K., and Kapil, A. (2005). High-level ciprofloxacin resistance in *Salmonella enterica* serotype Typhi in India. *J. Med. Microbiol.* 54, 999–1000. doi: 10.1099/jmm.0.45966-0
- Roumagnac, P., Weill, F. X., Dolecek, C., Baker, S., Brisse, S., Chinh, N. T., et al. (2006). Evolutionary history of *Salmonella* Typhi. *Science* 314, 1301–1304. doi: 10.1126/science.1134933
- Song, Y., Roumagnac, P., Weill, F. X., Wain, J., Dolecek, C., Mazzoni, C. J., et al. (2010). A multiplex single nucleotide polymorphism typing assay for detecting mutations that result in decreased fluoroquinolone susceptibility in *Salmonella enterica* serovars Typhi and Paratyphi A. *J. Antimicrob. Chemother.* 65, 1631–1641. doi: 10.1093/jac/dkq175
- Velge, P., Cloeckaert, A., and Barrow, P. (2005). Emergence of *Salmonella* epidemics: the problems related to *Salmonella enterica* serotype Enteritidis and multiple antibiotic resistance in other major serotypes. *Vet. Res.* 36, 267–288. doi: 10.1051/vetres:2005005

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 30 October 2013; accepted: 09 January 2014; published online: 27 January 2014.

Citation: Baucheron S, Monchaux I, Le Hello S, Weill F-X and Cloeckaert A (2014) Lack of efflux mediated quinolone resistance in *Salmonella enterica* serovars Typhi and Paratyphi A. *Front. Microbiol.* 5:12. doi: 10.3389/fmicb.2014.00012

This article was submitted to Antimicrobials, Resistance and Chemotherapy, a section of the journal Frontiers in Microbiology.

Copyright © 2014 Baucheron, Monchaux, Le Hello, Weill and Cloeckaert. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.