



## Discovery of hantavirus circulating among *Rattus rattus* in French Mayotte island, Indian Ocean.

Claudia Filippone, Guillaume Castel, Séverine Murri, Frédéric Beaulieux, Myriam Ermonval, Corinne Jallet, Emma L Wise, Richard J Ellis, Denise A Marston, Lorraine M McElhinney, et al.

### ► To cite this version:

Claudia Filippone, Guillaume Castel, Séverine Murri, Frédéric Beaulieux, Myriam Ermonval, et al.. Discovery of hantavirus circulating among *Rattus rattus* in French Mayotte island, Indian Ocean.. Journal of General Virology, 2016, 97 (5), pp.1060-1065. 10.1099/jgv.0.000440 . pasteur-01350075

HAL Id: pasteur-01350075

<https://pasteur.hal.science/pasteur-01350075>

Submitted on 28 May 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

Copyright

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Short Communication

# Discovery of hantavirus circulating among *Rattus rattus* in French Mayotte island, Indian Ocean

Claudia Filippone,<sup>1†</sup> Guillaume Castel,<sup>2†</sup> Séverine Murri,<sup>3</sup> Frédéric Beaulieux,<sup>4</sup> Myriam Ermonval,<sup>1</sup> Corinne Jallet,<sup>1</sup> Emma L. Wise,<sup>5</sup> Richard J. Ellis,<sup>5</sup> Denise A. Marston,<sup>5</sup> Lorraine M. McElhinney,<sup>5,6</sup> Anthony R. Fooks,<sup>5,6</sup> Amélie Desvars,<sup>7‡</sup> Lénaïg Halos,<sup>8</sup> Gwenaël Vourc'h,<sup>7</sup> Philippe Marianneau<sup>3</sup> and Noël Tordo<sup>1</sup>

### Correspondence

Noël Tordo  
ntordo@pasteur.fr

<sup>1</sup>Institut Pasteur, Unité des Stratégies Antivirales, F-75015, Paris, France

<sup>2</sup>INRA, UMR 1062 CBGP, F-34988 Montferrier-sur-Lez, France and Institut de Biologie Computationnelle, 34095 Montpellier, France

<sup>3</sup>Anses-Laboratoire de Lyon, Unité de Virologie, Lyon, France

<sup>4</sup>Centre de Recherche Clinique, Hôpital Croix Rousse, Lyon, France

<sup>5</sup>Wildlife Zoonoses and Vector-borne Diseases Research Group, Animal and Plant Health Agency, Woodham Lane, New Haw, Surrey, KT15 3NB, UK

<sup>6</sup>Department of Clinical Infection, Microbiology and Immunology, University of Liverpool, Liverpool, UK

<sup>7</sup>UR346 Animal Epidemiology, INRA, Saint Genès Champanelle, France

<sup>8</sup>Mérial, Lyon, France

Hantaviruses are emerging zoonotic viruses that cause human diseases. In this study, sera from 642 mammals from La Réunion and Mayotte islands (Indian Ocean) were screened for the presence of hantaviruses by molecular analysis. None of the mammals from La Réunion island was positive, but hantavirus genomic RNA was discovered in 29/160 (18 %) *Rattus rattus* from Mayotte island. The nucleoprotein coding region was sequenced from the liver and spleen of all positive individuals allowing epidemiological and intra-strain variability analyses. Phylogenetic analysis based on complete coding genomic sequences showed that this Murinae-associated hantavirus is a new variant of Thailand virus. Further studies are needed to investigate hantaviruses in rodent hosts and in Haemorrhagic Fever with Renal Syndrome (HFRS) human cases.

Received 5 November 2015

Accepted 27 February 2016

Viruses belonging to the genus *Hantavirus*, family *Bunyaviridae*, are negative-sense tri-segmented RNA viruses, with L, M and S segments encoding the RNA-dependent RNA polymerase, glycoproteins Gc and Gn, and nucleoprotein, respectively. Hantaviruses are known to circulate in small mammals (rodents, shrews, moles and bats) in Europe, Asia, America and, more recently discovered,

†These authors contributed equally to this work.

‡Present address: Research Institute of Wildlife Ecology, University of Veterinary Medicine, Savoyenstrasse 1, A-1160 Vienna, Austria.

The GenBank/EMBL/DDBJ accession numbers for the sequences determined in this study are KT719320–KT719370 (N gene coding sequence), KT779091–KT779093 and KU587796 (S, M and L complete coding sequence).

One supplementary figure is available with the online Supplementary Material.

Africa. Infection of humans occurs through inhalation of rodents' aerosolized excreta and can cause two severe pathologies: Haemorrhagic Fever with Renal Syndrome (HFRS) in Asia and Europe, and Hantavirus Cardiopulmonary Syndrome (HCPS), in the Americas (Kruger *et al.*, 2015).

An extensive capture of almost 4000 wild and domestic mammals was conducted in La Réunion, Maurice and Mayotte islands between 2006 and 2007, during Chikungunya virus (CHIKV) outbreaks. Although a low seroprevalence against CHIKV was detected in several non-human primates and rats (Vourc'h *et al.*, 2014), the sample collection was used for virus hunting. Here, we report on an investigation of hantaviruses in 642 small wild mammals captured in La Réunion (193 *Rattus rattus*, 44 *Rattus norvegicus*, 67 *Mus musculus*, 133 *Suncus murinus*, 45 *Tenrec ecaudatus*) and Mayotte (160 *Rattus rattus*) islands.

Pools of five individual sera were prepared for the collections from both La Réunion and Mayotte islands.

RNA was extracted from the pools using the MagAttract Viral RNA M48 kit and BioRobot M48 (Qiagen). The following reverse transcription (RT)-PCRs were performed for hantavirus screening: (i) consensus RT-PCR using degenerate primers targeting the L gene (Klempa *et al.*, 2006) for all samples; and (ii) specific RT-PCRs for insectivore hantaviruses targeting both the L and S segments (Kang *et al.*, 2009) for insectivore samples. Sera from positive pooled RNA were subsequently screened independently by RT-PCR. Although none of the samples from La

Réunion was positive for hantavirus RNA, we observed hantavirus L gene amplicons for 18 % (29/160) of *Rattus rattus* samples from Mayotte island. Rat species identification was confirmed by a RT-PCR targeting the cytochrome c oxidase gene and using the RodentSEA Identification Tool ([http://www.ceropath.org/barcoding\\_tool/rodentsea](http://www.ceropath.org/barcoding_tool/rodentsea)) for three representative animals (Fig. S1, available in the online Supplementary Material).

The characteristics of the 29 positive individuals are described in Table 1. Thirteen positive individuals (44.8 %) were males and 16 (55.2 %) were females, reflecting the gender proportion within all captured animals

**Table 1.** Description of hantavirus positive *Rattus rattus* in Mayotte island

Epidemiological information, geo-coordinates of captures and molecular analysis (RT-PCR; sequencing) are indicated for each individual.

| <i>R. rattus</i> ID | Sex | Age* | Capture area† | Latitude  | Longitude | RT-PCR (L)‡ |        | RT-PCR (S)/ sequences§ |        | GenBank accession numbers |
|---------------------|-----|------|---------------|-----------|-----------|-------------|--------|------------------------|--------|---------------------------|
|                     |     |      |               |           |           | Serum       | Spleen | Liver                  | Spleen |                           |
| 463                 | M   | a    | SE - 1        | -12.98468 | 45.18128  | +           | +      | +                      | +      | KT719351-KT719352         |
| 468                 | M   | a    | SE - 1        | -12.98468 | 45.18128  | +           | -      | +                      | +      | KT719353-KT719354         |
| 469                 | M   | j    | SE - 1        | -12.98468 | 45.18128  | +           | +      | +                      | +      | KT719355-KT719356         |
| 470                 | F   | a    | SE - 1        | -12.98468 | 45.18128  | +           | +      | +                      | +      | KT719357-KT719358         |
| 90                  | F   | a    | SE - 2        | -12.97011 | 45.16034  | +           | +      | +                      | +      | KT719366-KT719367         |
| 412                 | F   | a    | SE - 4        | -12.91093 | 45.1904   | +           | +      | +                      | +      | KT719329-KT719330         |
| 422                 | M   | a    | SE - 4        | -12.91093 | 45.1904   | +           | +      | +                      | +      | KT719331-KT719332         |
| 423                 | M   | a    | SE - 4        | -12.91093 | 45.1904   | +           | +      | +                      | +      | KT719333-KT719334         |
| 96                  | M   | a    | SE - 5        | -12.92727 | 45.177    | +           | +      | +                      | -      | KT719368                  |
| 99                  | M   | a    | SE - 5        | -12.92727 | 45.177    | +           | +      | +                      | +      | KT719369-KT719370         |
| 102                 | F   | a    | SE - 5        | -12.92727 | 45.177    | +           | -      | +                      | -      | KT719320                  |
| 425                 | F   | a    | SE - 5        | -12.92727 | 45.177    | +           | +      | +                      | +      | KT719335                  |
| 426                 | F   | a    | SE - 5        | -12.92727 | 45.177    | +           | +      | +                      | -      | KT719336                  |
| 428                 | F   | a    | SE - 5        | -12.92727 | 45.177    | +           | +      | +                      | +      | KT719337-KT719338         |
| 429                 | F   | NA   | SE - 5        | -12.92727 | 45.177    | +           | +      | +                      | -      | KT719339                  |
| 376                 | M   | a    | W - 6         | -12.83119 | 45.13716  | +           | +      | +                      | +      | KT719321-KT719322         |
| 379                 | M   | a    | W - 6         | -12.83119 | 45.13716  | +           | +      | +                      | +      | KT719323-KT719324         |
| 382                 | M   | a    | W - 6         | -12.83119 | 45.13716  | +           | +      | -                      | +      | KT719325-KT719326         |
| 408                 | F   | a    | W - 7         | -12.87164 | 45.11745  | +           | +      | +                      | +      | KT719327-KT719328         |
| 441                 | M   | a    | N - 8         | -12.70017 | 45.12168  | +           | +      | +                      | +      | KT719340-KT719341         |
| 444                 | F   | a    | N - 8         | -12.70017 | 45.12168  | +           | +      | -                      | +      | KT719342                  |
| 445                 | M   | a    | N - 8         | -12.70017 | 45.12168  | +           | +      | +                      | +      | KT719343-KT719344         |
| 449                 | F   | a    | N - 8         | -12.70017 | 45.12168  | +           | +      | +                      | +      | KT719345-KT719346         |
| 456                 | F   | a    | N - 8         | -12.70017 | 45.12168  | +           | +      | +                      | +      | KT719347-KT719348         |
| 457                 | F   | a    | N - 8         | -12.70017 | 45.12168  | +           | +      | +                      | +¶     | KT719349-KT719350         |
| 76                  | M   | a    | W - 9         | -12.76478 | 45.10652  | +           | +      | -                      | +      | KT719363                  |
| 82                  | F   | a    | W - 9         | -12.76478 | 45.10652  | +           | +      | +                      | +      | KT719364-KT719365         |
| 493                 | F   | a    | W - 9         | -12.76478 | 45.10652  | +           | +      | +                      | +¶     | KT719359-KT719360         |
| 494                 | F   | a    | W - 9         | -12.76478 | 45.10652  | +           | +      | +                      | +      | KT719361-KT719362         |

\*a, Adult; j, juvenile; NA, information not available.

†SE, South-east; N, north; W, west.

‡Klempa *et al.* (2006).

§GenBank accession numbers KT719320-KT719370.

||Different sequences from spleen and liver of the same individual. E382: C/T spleen/liver at nt 945 (corresponding to nt 957 of the coding region) of the S segment. This polymorphism is a silent mutation.


¶Only half the length of the sequence obtained (562 nt vs 1005 nt).

(44 and 56 %, respectively). No significant association was observed between gender and infection ( $P=0.923$ ), therefore male and female black rats were equally susceptible to hantavirus infection (13/70 : 18.6 % and 16/89 : 18.0 %, respectively). Of the 28 individuals with known age, 27 (96.4 %) were adults, which was significantly higher ( $P<0.05$ ) than the proportion of adults within the whole sampled population (58.8 %). Despite the limited number of individuals, our results support the hypothesis that hantavirus infection is not acquired vertically or during the neonatal stage, as demonstrated previously experimentally (Taruishi *et al.*, 2008).

A precise location of each capture according to geo-coordinate (latitude and longitude) values (Table 1) allowed the differentiation of nine geographical areas across Mayotte island according to the virus circulation (Fig. 1). Five areas were in the south-east: areas 1 (prevalence 11.8 %), 2 (16.7 %), 3 (0 %), 4 (23.1 %) and 5 (26.9 %); three were in the west of the island: areas 6 (13.6 %), 7 (5.6 %) and 9 (17.4 %); and one was in the north: area 8 (35.3 %). We have noticed two main prevalence foci in the north (area 8) and south-east (area 5) of the island. However, the relatively low number of samples per area did not allow statistical comparison tests to be applied to the regional prevalence data. Additional samples and further ecological

analysis are required to fully elucidate hantavirus transmission and circulation in Mayotte island.

To explore further the hantavirus genetic diversity in *Rattus rattus* from Mayotte island, RNA was extracted (QIAamp Viral RNA Extraction kit and QIAcube; Qiagen) from homogenized (TissueLyser; Qiagen) liver and spleen from the 29 positive animals. The majority of the spleen (27/29=93.1 %) and liver (26/29=90.0 %) samples tested positive using the nested RT-PCR targeting the L segment (Table 1). Both degenerate and specific primers were designed along the S segment by multiple alignments of the Murinae-associated hantavirus sequences (primers available on request) to obtain the nucleoprotein coding sequence from at least one organ per animal. Sequences from both organs (22/26) were identical to each other, except for one individual (E382) where a C/T polymorphism was observed (spleen/liver) at position 957 of the coding region on the S segment (Table 1), this mutation being silent at the amino acid level. The determined partial N gene coding sequences (1005 nt) within the S segment (nt 13–1017), were used. The method used was the maximum-likelihood method with PhyML v.3.0 (Guindon *et al.*, 2010) under the GTR (general time reversible) substitution model with a  $\gamma$ -distribution model among site rate heterogeneity and a proportion of invariant sites




**Fig. 1.** Hantavirus prevalence and phylogeography in *Rattus rattus* from Mayotte island. Prevalence and intra-strain variability of MAYOV hantavirus according to localization of the captured *R. rattus*. Phylogenetic analysis was done using partial S segment sequences (nt 13–1017). Anjouzorobe virus, previously detected in Madagascar (GenBank accession no. KC490916) was used as an outgroup. Bar, mean number of nucleotide substitutions per site. Colour codes correspond between the map and the tree.

(GTR + G + I), as determined by MEGA v.6.0 (Tamura *et al.*, 2013) and with a statistical approximate likelihood ratio test of branch support. Fig. 1 shows the phylogenetic pattern merged with the virus distribution across the island. The genetic cluster of most of the hantavirus sequences appeared to follow a geographical trajectory with a probable gradient of the virus distribution from north to south. However, individuals E96 and E379 did not follow this pattern, branching out of their group.

More exhaustive genetic information for this newly observed hantavirus was obtained from a representative individual (E469) by Next-Generation Sequencing (NGS) (Illumina MiSeq; Illumina). RNA was extracted and depleted of genomic DNA and ribosomal RNA using a protocol optimized for RNA viruses (Marston *et al.*, 2013). Double-stranded cDNA and subsequent library preparation for Illumina sequencing were undertaken as described previously (Marston *et al.*, 2013). Host sequences were removed by

mapping to the *Rattus rattus* reference genome (BWA-mem v.0.7.5). BLAST analysis of the assembled contigs of non-host reads (Velvet v.1.2.10) was used to select the most similar reference sequence [Anjozorobe virus (ANJV)]. Iterative mapping and intermediate consensus alignment was used to generate the final consensus sequence. RT-PCR targeting with Sanger sequencing was undertaken to fill missing gaps in the sequence. Complete protein coding sequences (S, M and L) of the newly identified virus isolate, named Mayotte virus (MAYOV), were obtained. Sequences from the S and M segments were used for phylogenetic analysis (Fig. 2a, b), following the methodology described above. This analysis showed that MAYOV clusters within the Thailand hantavirus clade closer to the isolates circulating in the Indian Ocean, South-East Asia (Hugot *et al.*, 2006; Plyusnina *et al.*, 2009; Johansson *et al.*, 2010) and Madagascar (Reynes *et al.*, 2014), rather than the Murinae-associated hantaviruses circulating in continental Africa (Klempa *et al.*, 2006, 2012; Witkowski *et al.*, 2014). The data suggested that both



**Fig. 2.** Phylogenetic analysis of Murinae-associated hantaviruses including MAYOV. Phylogenetic analysis was performed for the S (a) and M (b) segments among the Murinae-associated hantaviruses. MAYOV sequences were obtained by next-generation sequencing (S, 142 reads; M, 306 reads) and Sanger sequencing. The sequences used for phylogenetic analysis were obtained from GenBank: Sangassou virus (SANGV) (N: JQ082303, JQ082300; Gc-Gn: JQ082301); Dobrava virus (DOBV)/Saaremaa virus (SAAV) (N: AJ269550, AJ009773, AJ616854, L41916, AJ410619, NC005233; Gc-Gn: AY168578, AJ009774, L33685, AY168577); Amur/Soochong virus (ASV) (N: AB127997, AY675349, AY675350; Gc-Gn: AB127994, AY675353, DQ056293); Hantaan virus (HTNV) (N: AF329390, AF366568, JN712306, EU363809, EU092221; Gc-Gn: AF035831, AF366569, EU363818, EU092225); Gou virus (GOUV) (N: AF184988; Gc-Gn: AF145977); Seoul virus (SEOV) (N: JQ665919, AF187082, FJ803202, FJ803207, GQ279380, FJ803215; Gc-Gn: JQ665895, AF187081, GU592931, GQ274942, KF387724); Thailand virus (THAIIV) (N: AM397664, AB186420; Gc-Gn: L08756); Serang virus (SERV) (N: AM998808, GO274941; Gc-Gn: GO274939); Anjozorobe (N: KC490914-KC490918; Gc-Gn: KC490919-KC490921). The Kazan strain of Puumala virus (Z84204, Z84205) was used as an outgroup for both trees. Bars, mean number of nucleotide substitutions per site.

MAYOV and Anjozorobe virus, the closest variant within the Thailand hantavirus clade, may have resulted from a westward expansion of an ancestral South-East Asian hantavirus introduced with *R. rattus* (Cheke & Hume, 2008; Tollenaere *et al.*, 2010). Pairwise sequence identities between their coding regions, calculated with the BLASTN tool available online at <http://blast.ncbi.nlm.nih.gov/Blast.cgi>, were 91/99 % (nucleotide/amino acid identity) for the S segment, 90–91/97–98 % (nucleotide/amino acid identity) for the M segment (for ATD49 and AT261 Anjozorobe virus strains, respectively) and 90/99 % (nucleotide/amino acid identity) for the L segment. It is likely that exchanges between the Indian Ocean islands may have occurred as indicated by the phylogenetic trees in Fig. 2: Fig. 2(a) indicates that MAYOV and Anjozorobe virus have a common ancestor from which MAYOV diverged first, whereas Fig. 2(b) is less definitive and shows only the common ancestor of these two viral strains.

## Conclusions

This study was performed in the framework of a campaign aimed at elucidating potential reservoirs of CHIKV, which induced several outbreaks in the Indian Ocean between 2005 and 2007. While screening for hantaviruses, we demonstrated the prevalence (18 %) of a newly identified isolate in *R. rattus* on Mayotte island.

Interestingly, antibodies against hantaviruses were reported previously in rats from the Eastern Horn of Africa (Rodier *et al.*, 1993). One hypothesis is that hantaviruses were introduced by the shipping route to East Africa through infected *R. rattus*, originating from South-East Asia, via the Middle East during Arabian trade in the 10th century, and then shipped in the same reservoir host to the Indian Ocean islands (Rollin *et al.*, 1986; Brouat *et al.*, 2014).

MAYOV could have been acquired earlier by *R. rattus* through a spillover infection event from other hantavirus rodent reservoirs such as *Bandicota indica* for Thailand virus and *Rattus tanezumi* for Jurong/Serang virus in South-East Asia (Thailand, Indonesia) (Hugot *et al.*, 2006; Plyusnina *et al.*, 2009; Johansson *et al.*, 2010). The poor intrinsic genetic variability of MAYOV in *R. rattus* reflects a limited evolution and suggests a relatively recent colonization of Mayotte island. Further studies are required to explore hantavirus and rodent genetic diversity to understand better the origin of MAYOV and its adaptation to local hosts. Identifying the presence of hantavirus in Mayotte island is of great importance to evaluate its distribution in the tropical Indian Ocean region. A more extensive surveillance of rodents and/or other reservoirs is required to fully understand the circulation of these zoonotic viruses and to assess their potential risk of transmission to humans and subsequent occurrence of HFRS.

## Acknowledgements

This study was supported by the EU FP7 programmes: EDENext (no. 261504), EMPERIE (no. 223498), ANTIGONE (no. 278976)

and the Research Infrastructure Grant European Virus Archive (no. 19 228292). We thank Sandra Lacote (Virology Unit, Anses-Laboratoire, Lyon, France) for helping during RNA extraction. A. R. F. was supported by the Research and Policy for Infectious Disease Dynamics Program (RAPIDD), Science and Technology Directorate, US Department of Homeland Security and Fogarty International Center, US National Institutes of Health. Sampling was supported by the French Research National Agency ChikAni (ANR no. 06SEST06). We thank all the people who have participated in the collection of data in the field. We thank Dr Alexander Plyusnin for critical reading of the manuscript.

## References

- Brouat, C., Tollenaere, C., Estoup, A., Loiseau, A., Sommer, S., Soanandrasana, R., Rahalison, L., Rajerison, M., Piry, S. & other authors (2014). Invasion genetics of a human commensal rodent: the black rat *Rattus rattus* in Madagascar. *Mol Ecol* **23**, 4153–4167.
- Cheke, A. & Hume, J. (2008). *Lost Land of the Dodo: The Ecological History of Mauritius, Réunion & Rodrigues*. London: T. & A.D. Poyser.
- Guindon, S., Dufayard, J. F., Lefort, V., Anisimova, M., Hordijk, W. & Gascuel, O. (2010). New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst Biol* **59**, 307–321.
- Hugot, J. P., Plyusnina, A., Herbreteau, V., Nemirov, K., Laakkonen, J., Lundkvist, A., Supputamongkol, Y., Henttonen, H. & Plyusnina, A. (2006). Genetic analysis of Thailand hantavirus in *Bandicota indica* trapped in Thailand. *Virol J* **3**, 72.
- Johansson, P., Yap, G., Low, H. T., Siew, C. C., Kek, R., Ng, L. C. & Bucht, G. (2010). Molecular characterization of two hantavirus strains from different *Rattus* species in Singapore. *Virol J* **7**, 15.
- Kang, H. J., Bennett, S. N., Sumibcay, L., Arai, S., Hope, A. G., Mocz, G., Song, J. W., Cook, J. A. & Yanagihara, R. (2009). Evolutionary insights from a genetically divergent hantavirus harbored by the European common mole (*Talpa europaea*). *PLoS One* **4**, e6149.
- Klempa, B., Fichet-Calvet, E., Lecompte, E., Auste, B., Aniskin, V., Meisel, H., Denys, C., Koivogui, L., ter Meulen, J. & Krüger, D. H. (2006). Hantavirus in African wood mouse, Guinea. *Emerg Infect Dis* **12**, 838–840.
- Klempa, B., Witkowski, P. T., Popugaeva, E., Auste, B., Koivogui, L., Fichet-Calvet, E., Strecker, T., Ter Meulen, J. & Krüger, D. H. (2012). Sangassou virus, the first hantavirus isolate from Africa, displays genetic and functional properties distinct from those of other Murinae-associated hantaviruses. *J Virol* **86**, 3819–3827.
- Kruger, D. H., Figueiredo, L. T., Song, J. W. & Klempa, B. (2015). Hantaviruses – globally emerging pathogens. *J Clin Virol* **64**, 128–136.
- Marston, D. A., McElhinney, L. M., Ellis, R. J., Horton, D. L., Wise, E. L., Leech, S. L., David, D., de Lamballerie, X. & Fooks, A. R. (2013). Next generation sequencing of viral RNA genomes. *BMC Genomics* **14**, 444.
- Plyusnina, A., Ibrahim, I. N. & Plyusnina, A. (2009). A newly recognized hantavirus in the Asian house rat (*Rattus tanezumi*) in Indonesia. *J Gen Virol* **90**, 205–209.
- Reynes, J. M., Razafinalambo, N. K., Lacoste, V., Olive, M. M., Barivelio, T. A., Soarimalala, V., Heraud, J. M. & Lavergne, A. (2014). Anjozorobe hantavirus, a new genetic variant of Thailand virus detected in rodents from Madagascar. *Vector Borne Zoonotic Dis* **14**, 212–219.
- Rodier, G., Soliman, A. K., Bouloumié, J. & Kremer, D. (1993). Presence of antibodies to Hantavirus in rat and human populations of Djibouti. *Trans R Soc Trop Med Hyg* **87**, 160–161.
- Rollin, P. E., Mathiot, C., Nawrocka, E., Ravaoalimalala, V. E., Coulanges, P., Sureau, P. & McCormick, J. B. (1986). [Hemorrhagic

fever with renal syndrome in Madagascar. First seroepidemiologic survey of rat populations]. *Arch Inst Pasteur Madagascar* 52, 181–186 (in French).

**Tamura, K., Stecher, G., Peterson, D., Filipski, A. & Kumar, S. (2013).** MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Mol Biol Evol* 30, 2725–2729.

**Taruishi, M., Yoshimatsu, K., Hatsuse, R., Okumura, M., Nakamura, I. & Arikawa, J. (2008).** Lack of vertical transmission of Hantaan virus from persistently infected dam to progeny in laboratory mice. *Arch Virol* 153, 1605–1609.

**Tollenaere, C., Brouat, C., Duplantier, J. M., Rahalison, L., Rahelinirina, S., Pascal, M., Moné, H., Mouahid, G., Leirs, H. &**

**Cosson, J.-F. (2010).** Phylogeography of the introduced species *Rattus rattus* in the western Indian Ocean, with special emphasis on the colonization history of Madagascar. *J Biogeogr* 37, 398–410.

**Vourc'h, G., Halos, L., Desvars, A., Boué, F., Pascal, M., Lecollinet, S., Zientara, S., Duval, T., Nzanza, A. & Brémont, M. (2014).** Chikungunya antibodies detected in non-human primates and rats in three Indian Ocean islands after the 2006 ChikV outbreak. *Vet Res* 45, 52.

**Witkowski, P. T., Klempa, B., Ithete, N. L., Auste, B., Mfune, J. K., Hoveka, J., Matthee, S., Preiser, W. & Kruger, D. H. (2014).** Hantaviruses in Africa. *Virus Res* 187, 34–42.