



HAL
open science

Inhibition of DPP4 activity in humans establishes its in vivo role in CXCL10 post-translational modification: prospective placebo-controlled clinical studies

Jérémie Decalf, Kristin V Tarbell, Armanda Casrouge, Jeffrey D Price, Grace Linder, Estelle Mottez, Philippe Sultanik, Vincent Mallet, Stanislas Pol, Darragh Duffy, et al.

► To cite this version:

Jérémie Decalf, Kristin V Tarbell, Armanda Casrouge, Jeffrey D Price, Grace Linder, et al.. Inhibition of DPP4 activity in humans establishes its in vivo role in CXCL10 post-translational modification: prospective placebo-controlled clinical studies. *EMBO Molecular Medicine*, 2016, 8 (6), pp.679 - 683. 10.15252/emmm.201506145 . pasteur-01380633

HAL Id: pasteur-01380633

<https://pasteur.hal.science/pasteur-01380633>

Submitted on 13 Oct 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Inhibition of DPP4 activity in humans establishes its *in vivo* role in CXCL10 post-translational modification: prospective placebo-controlled clinical studies

Jérémy Decalf^{1,2,†}, Kristin V Tarbell^{3,†}, Armanda Casrouge^{1,2}, Jeffrey D Price³, Grace Linder³, Estelle Mottez⁴, Philippe Sultanik⁵, Vincent Mallet⁵, Stanislas Pol⁵, Darragh Duffy^{1,2,4,*,‡} & Matthew L Albert^{1,2,4,6,**,‡}

Abstract

Biochemical experiments, animal models, and observational studies in humans all support a role of dipeptidyl peptidase 4 (DPP4) in the N-terminal truncation of CXCL10, which results in the generation of an antagonist form of the chemokine that limits T-cell and NK cell migration. Motivated by the ability to regulate lymphocyte trafficking *in vivo*, we conducted two prospective clinical trials to test the effects of DPP4 inhibition on CXCL10 processing in healthy donors and in chronic hepatitis C patients, a disease in which DPP4 levels are found to be elevated. Participants were treated daily with 100 mg sitagliptin, a clinically approved DPP4 inhibitor. Plasma samples were analyzed using an ultrasensitive single-molecule assay (Simoa) to distinguish the full-length CXCL10_{1–77} from the NH₂-truncated CXCL10_{3–77}, as compared to the total CXCL10 levels. Sitagliptin treatment resulted in a significant decrease in CXCL10_{3–77} concentration, a reciprocal increase in CXCL10_{1–77}, with only minimal effects on total levels of the chemokine. These data provide the first direct evidence that *in vivo* DPP4 inhibition in humans can preserve the bioactive form of CXCL10, offering new therapeutic opportunities for DPP4 inhibitors.

Keywords chemokines; clinical study; CXCL10; DPP4; post-translational modifications

Subject Categories Microbiology, Virology & Host Pathogen Interaction; Pharmacology & Drug Discovery; Post-translational Modifications, Proteolysis & Proteomics

DOI 10.15252/emmm.201506145 | Received 14 December 2015 | Revised 19 February 2016 | Accepted 16 March 2016 | Published online 9 April 2016

EMBO Mol Med (2016) 8: 679–683

Introduction

Chemokines play an essential role in cell migration. Regulation of their activity is particularly important during inflammatory responses, determining the recruitment of immune cells to lymphoid organs or targeting them toward injured tissues (Griffith *et al*, 2014). Post-translational modification of chemokines has been shown to regulate their activity; however, *in vivo* evidence remains limited to observational studies and experimental mouse models (Moelants *et al*, 2013). Dipeptidyl peptidase 4 (DPP4, also known as CD26) is a serine protease capable of removing the first two amino acids of proteins possessing a proline or alanine in the N-terminal penultimate position (Bongers *et al*, 1992). *In vitro* studies have shown that DPP4-mediated N-terminal truncation of the pro-inflammatory chemokine CXCL10 leads to the generation of an antagonist form (Proost *et al*, 2001; Casrouge *et al*, 2011). Moreover, recent *in vivo* work performed in mice has demonstrated that this truncation alters lymphocyte migration and limits infiltration of the tumor parenchyma, a phenomenon that could be reversed using the DPP4 inhibitor sitagliptin (Barreira da Silva *et al*, 2015).

A challenge for studying post-translational modifications of chemokines is the ability to specifically monitor the different protein forms in biological material. To overcome this, we developed

1 The Laboratory of Dendritic Cell Immunobiology, Institut Pasteur, Paris, France

2 INSERM U818, Paris, France

3 Diabetes Endocrinology and Obesity Branch, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, MD, USA

4 Center for Human Immunology, Institut Pasteur, Paris, France

5 Département d'Hépatologie, AP-HP, Hôpital Cochin, Université Paris Descartes, INSERM UMS20, Institut Pasteur, Paris, France

6 Department of Cancer Immunotherapy, Genentech, South San Francisco, CA, USA

*Corresponding author. Tel: +33 1 44 38 93 34; E-mail: darragh.duffy@pasteur.fr

**Corresponding author. Tel: +33 1 44 38 93 34; E-mail: albertm@pasteur.fr

†These authors contributed equally to this work

‡These authors contributed equally to this work

immunoassays that discriminate the full-length agonist form of CXCL10 (referred to as CXCL10₁₋₇₇, or long CXCL10) from the NH₂-truncated form generated by DPP4 cleavage (referred to as CXCL10₃₋₇₇, or short CXCL10) (Casrouge *et al*, 2012). Accordingly, we were able to show that elevated levels of short CXCL10 were associated with an increased DPP4 activity, both being negative predictors for viral clearance in chronic and acute hepatitis C (HCV) patients (Casrouge *et al*, 2011; Ragab *et al*, 2013; Riva *et al*, 2014). Taking advantage of the Simoa technology (Rissin *et al*, 2010), we have now developed ultrasensitive immunoassays with these antibodies, making possible the quantification of functional forms of CXCL10 in the plasma from healthy individuals.

To provide a proof of concept for DPP4 inhibition as a means to protect full-length CXCL10, we conducted a randomized placebo-controlled study in healthy individuals, with the primary scientific objective being the assessment of DPP4 inhibition on immunological parameters. A prior report on this cohort demonstrated an effective DPP4 inhibition and an increased concentration of active glucagon-like peptide 1, one of the key substrates implicated in insulin resistance (Price *et al*, 2013). In parallel, we undertook an investigational study to explore how sitagliptin treatment affects virus-induced CXCL10 in chronic HCV patients (cHCV), a clinical setting in which levels of both DPP4 and CXCL10₃₋₇₇ have been shown to be elevated (Ragab *et al*, 2013). The findings from this report establish the basis for repositioning DPP4 inhibitors as a potential immunotherapy.

Results

To quantify CXCL10 agonist and antagonist forms in healthy individuals, we implemented our unique immunoassays on the Simoa platform (Fig 1A). Using these assays, we determined median levels of total CXCL10 to be 60 pg/ml in healthy individuals (Fig 1B and C, SV and D0 time points), within the range of what has been previously reported using other techniques (Butera *et al*, 2005; Duffy *et al*, 2014). Long CXCL10 remained undetectable in most individuals, whereas short CXCL10 was detected in 30 out of 36 subjects (83%) with concentrations ranging from 2 to 75 pg/ml, suggesting active *in vivo* chemokine processing in healthy individuals. Importantly, all the plasma samples used for the analysis of CXCL10 levels were collected in tubes containing a DPP4 inhibitor to avoid potential extracorporeal CXCL10 processing.

In order to explore the role of DPP4 in CXCL10 truncation *in vivo*, we monitored the levels of short, long, and total CXCL10 in healthy individuals receiving a 28-day course of placebo or sitagliptin (Fig 1B and C). DPP4 inhibition in individuals receiving sitagliptin was confirmed by monitoring plasma DPP4 activity, which was previously published (Price *et al*, 2013) and showed an inhibition ranging from 9 to 80% (Fig EV1). We observed that 3 days after the onset of sitagliptin treatment, the concentrations of short CXCL10 dropped significantly in individuals receiving sitagliptin, but remained stable in donors receiving the placebo. Effect size analysis supported the strong impact of sitagliptin on short CXCL10 when compared to pre-therapy levels ($d = 0.92$). As detailed in Fig EV1, some donors showed sporadic increases in short CXCL10 during sitagliptin therapy, possibly reflecting a partial recovery of DPP4 activity (Herman *et al*, 2005). Moreover, three of the 27 donors showed elevated levels of short CXCL10 during sitagliptin treatment,

reflecting natural human variability in response to the treatment. Of note, despite the presence of short CXCL10, two of these three donors showed a strong DPP4 inhibition.


Interestingly, the inhibition of DPP4 was associated with the preservation of long CXCL10, as indicated by an increase in concentration compared to pre-therapy ($d = 0.27$), although these data have to be interpreted cautiously due to a large number of patients in which long CXCL10 remained undetectable. That said, this increase was observed at week 2 of treatment, but not at day 3 post-treatment initiation, perhaps reflecting a low level of newly secreted CXCL10 in healthy individuals. Of note, the levels of total CXCL10 were found to be slightly decreased during sitagliptin treatment compared to pre-therapy levels ($d = 0.33$), suggesting a limited biological impact of sitagliptin on total CXCL10 levels. That said, the strong impact of sitagliptin on short CXCL10 indicated that we selectively altered processing, rather than the production of the chemokine. Finally, the modulation of short and long CXCL10 was stable during the 28-day course of sitagliptin, returning to pre-therapy concentrations once treatment was terminated. These data provide direct evidence that DPP4 inhibition impacts *in vivo* N-terminal truncation of CXCL10.

The data obtained in healthy individuals suggest that CXCL10 processing by DPP4 is a rapid event, as it was strongly affected 72 h after the onset of sitagliptin therapy. Therefore, we assessed how sitagliptin might affect higher levels of CXCL10, a hallmark of inflammatory diseases (Van Raemdonck *et al*, 2015). To do so, we monitored CXCL10 forms in three cHCV patients receiving sitagliptin treatment (Fig 1D). Of note, as detailed in the study design section, additional patients could not be recruited for ethical reasons. As previously described (Casrouge *et al*, 2011), cHCV patients showed elevated concentrations of short, long, and total CXCL10 compared to healthy individuals (D0 time points). Interestingly, sitagliptin treatment led to a decrease in short CXCL10 and an increase in long CXCL10 (Fig 1D), a trend similar to what we observed in healthy individuals. Although striking, the impact of sitagliptin on CXCL10 forms did not influence HCV viral loads over the period monitored (Fig EV2).

Discussion

Overall, the finding that the agonist form CXCL10₁₋₇₇ was undetectable in the majority of healthy individuals indicates that CXCL10 and likely other chemokines are rapidly catabolized by DPP4. The corollary to this observation is that long CXCL10 may be considered as a marker of recently produced CXCL10. Moreover, the lower levels of long and short forms compared to the total plasma concentration of CXCL10 indicate that additional processing of the protein is probably occurring *in vivo*. Other proteases, such as matrix metalloproteinases (Van den Steen *et al*, 2003), have been shown to target CXCL10, but their activity *in vivo* and the impact on chemokine function remain unknown (see Mortier *et al* (2008) for review of subject). Action of other N-terminal aminoproteases could also explain the trimming of CXCL10₃₋₇₇, acting after DPP4 removes the penultimate proline residue, as shown in *in vitro* biochemical studies using CXCL11 (Proost *et al*, 2007).

Our findings expose a broader putative *in vivo* role of DPP4 in the regulation of cell trafficking. Notably, other chemokine substrates of



DPP4 have been defined as key mediators of stem cell migration. In mouse studies, DPP4 inhibition has been shown to improve the engraftment of CD34⁺ stem cells following umbilical cord blood transplantation, with the mechanism of action being the protection of agonist forms of CXCL12 (Farag *et al*, 2013). While our study does

not directly test a role for altered post-translational modification of CXCL12, the similar biological processes would suggest a direct impact on CXCR4-mediated leukocyte migration.

From a therapeutic perspective, our results establish a path toward the secondary use of DPP4 inhibitors. Indeed, our recent

Table 1. Clinical information.

Category	Treatment	Number of subjects	Gender (F/M)	Age - median (range)	HCV genotype
Healthy	Placebo	9	6/3	27 (20–61)	–
Healthy	Sitagliptin	27	16/11	36 (19–53)	–
cHCV	Sitagliptin	3	2/1	41 (38–55)	4

mouse studies have shown that preserving agonist CXCL10₁₋₇₇ increases lymphocyte infiltration into tumor parenchyma and results in an enhanced tumor immunity (Barreira da Silva *et al*, 2015). Thus, these drugs may offer an opportunity to increase the lymphocyte migration in settings such as chronic infection and cancer. As DPP4 inhibitors are a class of widely used therapeutics for the management of type II diabetes, they are considered to be safe and well tolerated.

In sum, our work provides the first *in vivo* experimental evidence of CXCL10 processing in humans and supports the clinical testing of sitagliptin as either an antitumor or autoimmune therapy. Further work will be required to better understand the impact of DPP4 inhibition *in vivo* and to evaluate how other forms of chemokine post-translational modifications could influence inflammatory responses.

Materials and Methods

Study design and protocols

The cohort of healthy individuals receiving sitagliptin or placebo (NCT00813228) has been previously described. This study was a double-blind, randomized trial approved by the institutional review board of NIDDK (Price *et al*, 2013). Participants received 100 mg of sitagliptin ($n = 27$) or placebo ($n = 9$) once daily for 28 days, and the blood samples were collected longitudinally. Chronic HCV patients receiving sitagliptin ($n = 3$) were recruited as part of the INSERM-sponsored C10-54 trial (NCT01567540). The three patients were chronically infected with HCV genotype 4 and were receiving pegylated-interferon alpha + ribavirin therapy when they enrolled in the study. These patients were considered difficult to treat, as they had previously failed to achieve early virological response (EVR) as defined by a two-log reduction in virus at 12 weeks post-treatment. Sitagliptin treatment (100 mg daily) was given for 3 weeks and the blood samples were collected longitudinally. Of note, the C10-54 trial was terminated prematurely due to the approval of sofosbuvir in France for HCV genotype 4-infected patients. The ethical implications that arose from the availability of this novel highly efficacious antiviral therapy prevented the continued recruitment of additional cHCV patients in our experimental clinical study. The respective studies were approved by the institutional review boards of the NIH and INSERM. All participants gave written informed consent prior to inclusion in the study, conformed to the principles set out in the WMA Declaration of Helsinki and the Department of Health and Human Services Belmont Report. Clinical characteristics are summarized in Table 1. Plasma was collected in BD P700 tubes, containing ethylenediamine tetraacetic acid (EDTA) and a DPP4 inhibitor to prevent extracorporeal cleavage of CXCL10. Plasma collected in sodium heparin tubes was used for monitoring DPP4 activity. Samples were stored at -80°C until analysis.

CXCL10 quantification

Plasma concentration of total (R&D clone 33036), long (CXCL10₁₋₇₇), and short (CXCL10₃₋₇₇) CXCL10 was measured using Simoa technology (Quanterix). Simoa assays were carried out as

The paper explained

Problem

The N-terminal truncation of CXCL10 by DPP4 results in the generation of an antagonist form of the chemokine that limits T-cell and NK cell migration toward infectious or tumor sites. In this work, we studied whether DPP4 inhibition *in vivo* by sitagliptin could limit the generation of the antagonist form of CXCL10 in humans, which could have the potential to boost T-cell and NK cell migration in certain pathological contexts.

Results

Participants were treated daily with 100 mg sitagliptin, and plasma samples were analyzed using an ultrasensitive single-molecule assay (Simoa) to distinguish levels of the agonist (CXCL10₁₋₇₇), antagonist (CXCL10₃₋₇₇), and total CXCL10 forms. Sitagliptin treatment resulted in a significant decrease in antagonist CXCL10 concentration and a reciprocal increase in the agonist form CXCL10₁₋₇₇ compared to placebo controls.

Impact

Our data provide the first *in vivo* evidence that DPP4 inhibition in humans can preserve the bioactive form of CXCL10. This offers new therapeutic opportunities for DPP4 inhibitors, which could be relevant for the development of novel cancer immunotherapies aiming at restoring immune cell migration.

previously described (Meissner *et al*, 2015), and additional optimization was conducted for this study; standard curves are presented in Fig 1A.

Statistical analysis

Differences in CXCL10 forms in healthy donors receiving either sitagliptin or placebo were assessed by nonparametric Friedman's test. No statistical analysis was performed on HCV patients due to the available sample size ($n = 3$). For healthy donors receiving sitagliptin, size effect analysis using Cohen's d was performed to assess the biological significance of modified CXCL10 levels. Three groups of samples were compared; the values were obtained pre-therapy (SV, D0), during therapy (D3, W2, and W4), and post-therapy (W9). Cohen's d estimates the biological significance of statistically significant differences to be small ($d \approx 0.2$), medium ($d \approx 0.5$), or large ($d \approx 1$).

Expanded View for this article is available online.

Acknowledgements

This work was supported by the Agence nationale de recherches sur le sida et les hépatites virales (ANRS) and the Ligue nationale contre le cancer and by the Intramural Research Program of the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK).

Author contributions

JD and AC performed the experiments; JD, KVT, AC, MLA, and DD analyzed the data; JP, GL, EM, PS, VM, SP, KVT, MLA, and DD designed and executed the clinical studies; JD, DD, and MLA wrote the manuscript.

Conflict of interest

The authors declare that they have no conflict of interest.

References

- Barreira da Silva R, Laird ME, Yatim N, Fiette L, Ingersoll MA, Albert ML (2015) Dipeptidylpeptidase 4 inhibition enhances lymphocyte trafficking, improving both naturally occurring tumor immunity and immunotherapy. *Nat Immunol* 16: 850–858
- Bongers J, Lambros T, Ahmad M, Heimer EP (1992) Kinetics of dipeptidyl peptidase IV proteolysis of growth hormone-releasing factor and analogs. *Biochim Biophys Acta* 1122: 147–153
- Butera D, Marukian S, Iwamaye AE, Hembrador E, Chambers TJ, Di Bisceglie AM, Charles ED, Talal AH, Jacobson IM, Rice CM et al (2005) Plasma chemokine levels correlate with the outcome of antiviral therapy in patients with hepatitis C. *Blood* 106: 1175–1182
- Casrouge A, Decalf J, Ahloulay M, Lababidi C, Mansour H, Vallet-Pichard A, Mallet V, Mottez E, Mapes J, Fontanet A et al (2011) Evidence for an antagonist form of the chemokine CXCL10 in patients chronically infected with HCV. *J Clin Invest* 121: 308–317
- Casrouge A, Bisiaux A, Stephen L, Schmolz M, Mapes J, Pfister C, Pol S, Mallet V, Albert ML (2012) Discrimination of agonist and antagonist forms of CXCL10 in biological samples. *Clin Exp Immunol* 167: 137–148
- Duffy D, Mamdouh R, Laird M, Soneson C, Le Foul r L, El-Daly M, Casrouge A, Decalf J, Abbas A, Eldin NS et al (2014) The ABCs of viral hepatitis that define biomarker signatures of acute viral hepatitis. *Hepatology* 59: 1273–1282
- Farag SS, Srivastava S, Messina-Graham S, Schwartz J, Robertson MJ, Abonour R, Cornetta K, Wood L, Secrest A, Strother RM et al (2013) In vivo DPP-4 inhibition to enhance engraftment of single-unit cord blood transplants in adults with hematological malignancies. *Stem Cells Dev* 22: 1007–1015
- Griffith JW, Sokol CL, Luster AD (2014) Chemokines and chemokine receptors: positioning cells for host defense and immunity. *Annu Rev Immunol* 32: 659–702
- Herman GA, Stevens C, Van Dyck K, Bergman A, Yi B, De Smet M, Snyder K, Hilliard D, Tanen M, Tanaka W et al (2005) Pharmacokinetics and pharmacodynamics of sitagliptin, an inhibitor of dipeptidyl peptidase IV, in healthy subjects: results from two randomized, double-blind, placebo-controlled studies with single oral doses. *Clin Pharmacol Ther* 78: 675–688
- Meissner EG, Decalf J, Casrouge A, Masur H, Kottlilil S, Albert ML, Duffy D (2015) Dynamic Changes of Post-Translationally Modified Forms of CXCL10 and Soluble DPP4 in HCV Subjects Receiving Interferon-Free Therapy. *PLoS ONE* 10: e0133236
- Moelants EAV, Mortier A, Van Damme J, Proost P (2013) In vivo regulation of chemokine activity by post-translational modification. *Immunol Cell Biol* 91: 402–407
- Mortier A, Van Damme J, Proost P (2008) Regulation of chemokine activity by posttranslational modification. *Pharmacol Ther* 120: 197–217
- Price JD, Linder G, Li WP, Zimmermann B, Rother KI, Malek R, Alattar M, Tarbell KV (2013) Effects of short-term sitagliptin treatment on immune parameters in healthy individuals, a randomized placebo-controlled study. *Clin Exp Immunol* 174: 120–128
- Proost P, Schutyser E, Menten P, Struyf S, Wuyts A, Opendakker G, Detheux M, Parmentier M, Durinx C, Lambeir AM et al (2001) Amino-terminal truncation of CXCR3 agonists impairs receptor signaling and lymphocyte chemotaxis, while preserving antiangiogenic properties. *Blood* 98: 3554–3561
- Proost P, Mortier A, Loos T, Vandercappellen J, Gouwy M, Ronsse I, Schutyser E, Put W, Parmentier M, Struyf S et al (2007) Proteolytic processing of CXCL11 by CD13/aminopeptidase N impairs CXCR3 and CXCR7 binding and signaling and reduces lymphocyte and endothelial cell migration. *Blood* 110: 37–44
- Ragab D, Laird M, Duffy D, Casrouge A, Mamdouh R, Abass A, Shenawy DE, Shebl AM, Elkashef WF, Zalata KR et al (2013) CXCL10 antagonism and plasma sDPP4V correlate with increasing liver disease in chronic HCV genotype 4 infected patients. *Cytokine* 63: 105–112
- Rissin DM, Kan CW, Campbell TG, Howes SC, Fournier DR, Song L, Piech T, Patel PP, Chang L, Rivnak AJ et al (2010) Single-molecule enzyme-linked immunosorbent assay detects serum proteins at subfemtomolar concentrations. *Nat Biotech* 28: 595–599
- Riva A, Laird M, Casrouge A, Ambrozaitis A, Williams R, Naoumov NV, Albert ML, Chokshi S (2014) Truncated CXCL10 is associated with failure to achieve spontaneous clearance of acute hepatitis C infection. *Hepatology* 60: 487–496
- Van den Steen PE, Husson SJ, Proost P, Van Damme J, Opendakker G (2003) Carboxyterminal cleavage of the chemokines MIG and IP-10 by gelatinase B and neutrophil collagenase. *Biochem Biophys Res Commun* 310: 889–896
- Van Raemdonck K, Van den Steen PE, Liekens S, Van Damme J, Struyf S (2015) CXCR3 ligands in disease and therapy. *Cytokine Growth Factor Rev* 26: 311–327



License: This is an open access article under the terms of the Creative Commons Attribution 4.0 License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.