

Strong Ifitm1 Expression In Cd4 T Cells In Hiv Controllers Is Correlated With Immune Activation

Etienne Canoui, Nicolas Noël, Camille Lécuroux, Faroudy Boufassa, Asier Sáez-Ciri3n, Christine Bourgeois, Olivier Lambotte, Anrs Co21 Codex Group

► **To cite this version:**

Etienne Canoui, Nicolas Noël, Camille Lécuroux, Faroudy Boufassa, Asier Sáez-Ciri3n, et al.. Strong Ifitm1 Expression In Cd4 T Cells In Hiv Controllers Is Correlated With Immune Activation. Journal of Acquired Immune Deficiency Syndromes, Lippincott, Williams

Wilkins, 2017, <10.1097/QAI.0000000000001166>. <pasteur-01420412>

HAL Id: pasteur-01420412

<https://hal-pasteur.archives-ouvertes.fr/pasteur-01420412>

Submitted on 16 May 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



1 **Strong *ifitm1* expression in CD4 T cells in HIV controllers is correlated with**
2 **immune activation**

3 Etienne Canoui^{1, *}, Nicolas Noël^{1,2,3,4 *}, Camille Lécuroux^{1,4}, Faroudy Boufassa^{2,5,6}, Asier
4 Sáez-Cirió⁷, Christine Bourgeois^{1,2,4}, Olivier Lambotte^{1, 2, 3,4 §} and the ANRS CO21 CODEX
5 study group

6
7 ¹: INSERM U1184, Immunologie des infections virales et autoimmunité, Le Kremlin-Bicêtre, France

8 ²: Université Paris Sud, Le Kremlin-Bicêtre, France

9 ³: Assistance Publique – Hôpitaux de Paris, Service de Médecine Interne et Immunologie Clinique,
10 Groupe Hospitalier Universitaire Paris Sud, Hôpital Bicêtre, Le Kremlin-Bicêtre, France

11 ⁴: CEA, DSV/iMETI, Division of Immuno-Virology, IDMIT, France

12 ⁵: INSERM U1018, Centre de recherche en Epidémiologie et Santé des Populations, Le Kremlin-
13 Bicêtre, France

14 ⁶: Assistance Publique – Hôpitaux de Paris, Département d'épidémiologie, Groupe Hospitalier
15 Universitaire Paris Sud, Hôpital Bicêtre, Le Kremlin-Bicêtre, France

16 ⁷: Institut Pasteur, Unité HIV inflammation et persistance, Paris, France

17 *: co-first authors, equal contribution

18

19 **§: Corresponding author**

20 Pr Olivier Lambotte, Service de Médecine Interne et Immunologie Clinique

21 CHU Bicêtre ; 78, rue du Général Leclerc, 94275 Le Kremlin-Bicêtre CEDEX, FRANCE

22 Phone: +33 145 212 783, Fax: +33 145 212 733

23 olivier.lambotte@aphp.fr

24

25 Count: Main document 1451 words, 1 figure, 20 references, 1 supplemental figure, 1
26 supplemental table

27 **Running head: *ifitm1* expression in HIV controllers**

28 **Conflicts of Interest and Source of Funding: none declared**

1 **Key words:** Type 1 Interferon, Interferon stimulated genes, IFITM1, HIV controllers,

2 HIV.

3

1 INTRODUCTION

2 During HIV infection, the Interferon (IFN)-stimulated genes (ISGs) have been
3 associated with HIV pathophysiology by restricting HIV replication [1], or enhancing
4 the immune activation [2,3]. The expression levels of ISGs depend on their function,
5 as well as their regulation mechanisms and the viral progression [4,5]. Indeed, most
6 ISGs are upregulated in CD4 and CD8 T cells from viremic untreated patients [5–7],
7 and tend to diminish on antiretroviral therapy (ART). However, whereas monocytes
8 are important in HIV pathogenesis [8], few studies have focused on ISGs regulation
9 in monocytes from HIV-infected patients [9].

10 Little is known regarding the expression of ISGs in the rare patients who
11 spontaneously control HIV replication (HIV controller patients (HICs)). In PBMCs from
12 HICs, *ifit1* and *mxA* seem overexpressed as compared with HDs and ART treated
13 patients [10]. We have recently showed that plasma IP10 levels are higher in HICs
14 than HDs and associated with low CD4 T cell counts [11]. Other studies reported that
15 some ISGs known to restrict HIV, such as *samhd1*, *schlafen 11*, *apobec3c* and
16 *apobec3d* might be overexpressed in CD4 T cells or PBMCs from HICs [12–14].

17 Here, we studied the expression of four ISGs in three sorted cell subtypes (CD4 and
18 CD8 T lymphocytes, and monocytes). We compared their expressions between
19 HICs, HIV-1 infected ART-treated (ARTs), viremic treatment-naïve patients (VIRs),
20 and healthy donors (HD). The expression of these ISGs according to the levels of T
21 cell- and monocyte activation was analyzed. A strong *ifitm1* expression in CD4 T
22 cells in HIV controllers was correlated with immune activation.

23

24

1 PATIENTS AND METHODS

2 HICs (n=11) were enrolled in the ANRS CO21 CODEX cohort [11] with the following
3 definition: known HIV diagnosis for at least 5 years prior to enrolment, with HIV VL <
4 400 copies/mL in the last five consecutive measurements. ART-treated patients (n=8)
5 and VIRs (n=8) were defined as previously reported [11]. HIV – negative healthy
6 donors (HDs) (n=8) were recruited as controls. All patients gave their informed
7 consent for this study, in accordance with the Helsinki declaration.

8 Freshly collected PBMCs were stained with the following conjugated monoclonal
9 antibodies: PE-CD14, PerCP Cy5.5-CD8, APC-CD3, and APC-H7-CD4, before the
10 FACS-based sorting procedure on BD ARIA (BD Biosciences™). CD4+ T cells,
11 CD8+ T cells and monocytes (CD3-CD14+) were sorted and immediately lysed in
12 RNeasy Lysis Buffer (QIAGEN*) and cryopreserved at -80°C. RNA isolation and
13 reverse transcription were performed according to the manufacturers' protocols
14 (RNeasy mini kit, Qiagen™, and enhanced avian RT-PCR Kit, SIGMA™,
15 respectively). qRT-PCR analysis was performed using a SyBR Green RT PCR
16 method (LightCycler system, Roche™). All results were normalized for *gapdh*
17 expression.

18 Levels of soluble CD14 (sCD14) were determined by ELISA (R&D Systems,
19 Minneapolis, Minnesota, USA), and the surface expression of HLA-DR and CD38 on
20 T cells was analyzed by flow cytometry, as described previously [11].

21 For statistical analyses, continuous variables were compared using the non-
22 parametric Kruskal-Wallis test followed by Dunn's test for multiple analysis.
23 Categorical variables were compared using chi² test, and Spearman's coefficient was
24 used for correlation analyses. The threshold for statistical significance was set to *p*

1 less than 0.05. Data were stored and analysed using PRISM software (version 5,
2 GraphPad software, La Jolla, California, USA).

3

1 RESULTS

2 The characteristics of the study population are summarized in the **Supplemental**
3 **Table 1**. There was no difference in terms of age and gender between each group of
4 patients.

5 ***Ifitm1* expression is elevated in CD4 T cells from HICs**

6 The analysis of ISGs expression in the 4 groups and the 3 cell subtypes is depicted
7 in **Figure 1**. In VIRs, the expression of *ifitm1* in monocytes, *pkr* in all cell types, *mxA*
8 in CD4 and CD8 T cells, and *ifit1* in CD4 were higher than in ARTs, suggesting a
9 strong role of the virus in ISGs induction. Similar results were seen when comparing
10 ISGs expression between VIRs and HICs, except for *mxA* in all cell types and *ifit1* in
11 CD8 T cells and monocytes, for which the expression were similar in the two groups.

12 Importantly, we found that the expression of *ifitm1* expression in CD4 T cells from
13 HICs ($p=0.01$) and ARTs ($p=0.001$) was similar to VIRs and higher than HDs. In
14 contrast, the expression of *ifitm1* in CD8 T cells and monocytes and those of the
15 other genes in the three cell subtypes were similar in HICs and in HDs.

16

17 **The expression of most ISGs is correlated positively with HIV viral load**

18 In view of these observations, we hypothesized that the expression of most ISGs was
19 associated with the HIV viral load (VL). Indeed, pooling the data of all HIV infected
20 patients, the HIV VL was significantly correlated with the expression of *pkr* in CD4 T
21 cells, CD8 T cells, and monocytes, as well as *mxA* in CD4 and CD8 T cells, and with
22 *ifit1* in CD4 T cells. Of note, similar results were obtained when classifying the HIV
23 VL into categorical variables (i.e., detectable or undetectable). Also, when

1 considering all HIV infected patients, the CD4 T cell count was negatively correlated
2 with the expression of *pkr* in CD4 T cells and monocytes (with a strong trend for CD8
3 T cells), and *ifit1* only in CD4 T cells, but not with *mxA* expression.

4 Interestingly, the expression of *ifitm1* was associated with the HIV VL in monocytes
5 only, but was negatively correlated with CD4 count in every cell subtypes. These
6 results suggested that *ifitm1* expression could depend on other conditions than the
7 amount of virus itself, especially in T cells.

8

9 ***ifitm1* expression is associated with immune activation**

10 Thus, we lastly investigated the relationships between ISGs expression and immune
11 activation. When all HIV-1 infected patients were considered, positive correlations
12 were observed between *ifitm1* expression in each cellular subtype and the
13 corresponding activation markers (*i.e.* % of HLA-DR+CD38+ CD4+ T cells ($r= 0.4$, $p=$
14 0.03) and CD8+ T cells ($r= 0.4$, $p= 0.04$), and sCD14 for monocyte activation ($r=0.48$,
15 $p= 0.02$)) (**Supplemental Figure 1a**). Such relationships were not observed with the
16 other ISGs, except for *ifit1* in CD4 T cells. Moreover, the correlation between *ifitm1*
17 expression in CD4 T cells and the frequency of circulating HLADR+CD38+ CD4+ T
18 cells was still observed when HICs were analyzed separately ($r=0.8$, $p= 0.003$)
19 (**Supplemental Figure 1b**).

20

1 **DISCUSSION**

2 We selected 4 ISGs with distinct production mechanisms and reported their
3 expression in three cell subsets in four groups of individuals. *pkr* and *ifit1* expressions
4 depend on the engagement of type I IFN, but can be also directly stimulated by HIV
5 RNA or particles [15,16]. *mxA* seems to be directly correlated with type I IFN, with a
6 strong link with HIV viral load [17]. Conversely, *ifitm1* expression depends on type I,
7 but also type II IFN, with strong links to the proinflammatory milieu [16].
8 Consequently, viremic patients had higher levels of *pkr* in each cell subtypes, *mxA* in
9 CD4 and CD8 T cells, and *ifit1* in CD4 T cells than the other groups. The correlation
10 between these ISGs and the HIV viral load support the role of viral particles in their
11 production [5–7,18].

12 The most striking point was the finding that in HICs' CD4 T cells, *ifitm1* was
13 overexpressed compared with HDs, whereas no differences were found between
14 HICs and HD for the other ISGs. This observation extends previous results from
15 gene expression profiling of CD4 T cells from HICs but had never been quantified by
16 RT-PCR in sorted cells [13,14]. *ifitm1* has been recently described as part of a HIV-1
17 restriction factor family [19,20]. Moreover, it has been suggested that *ifitm1* might be
18 induced by IL-6 or other acute-phases cytokines [16]. This is consistent with our
19 observations that *ifitm1* was globally linked with immune activation but barely with the
20 viral load, the correlation being still present only when HICs are studied.

21 It seems important to separately analyze the expression of ISGs in purified CD4 T
22 cells and monocytes, since the role of monocytes in the pathophysiology of HIV and
23 inflammation is increasing. The analysis of the ex vivo purified monocytic
24 compartment has seldom been reported in the field of ISG expression. Indeed, we
25 observed that the expression profile of most ISGs was different between CD4 T cells

1 and monocytes, suggesting that the individualization of these cell types might better
2 decipher the mechanisms of ISGs regulation in the setting of HIV infection.

3 Our results show that the expression profile of ISGs in HIV infected patients varies
4 according to the cell type, the disease status and the gene considered, with multiple
5 regulation mechanisms beyond type I/II Interferons. Importantly, the gene expression
6 profiles in monocytes should be distinguished from that in CD4 T cells. In HICs, *ifitm1*
7 seems to have a unique expression profile and is associated with immune CD4 T cell
8 activation. Its putative role in HIV pathogenesis and antiviral control deserves further
9 investigations.

1 REFERENCES

- 2 1 Fenton-May AE, Dibben O, Emmerich T, Ding H, Pfafferott K, Aasa-Chapman MM, *et al.*
3 Relative resistance of HIV-1 founder viruses to control by interferon-alpha. *Retrovirology*
4 2013; **10**:146.
- 5 2 Fernandez S, Tanaskovic S, Helbig K, Rajasuriar R, Kramski M, Murray JM, *et al.* CD4+ T-
6 cell deficiency in HIV patients responding to antiretroviral therapy is associated with
7 increased expression of interferon-stimulated genes in CD4+ T cells. *J Infect Dis* 2011;
8 **204**:1927–1935.
- 9 3 Lederman MM, Calabrese L, Funderburg NT, Clagett B, Medvik K, Bonilla H, *et al.*
10 Immunologic failure despite suppressive antiretroviral therapy is related to activation
11 and turnover of memory CD4 cells. *J Infect Dis* 2011; **204**:1217–1226.
- 12 4 Liu S-Y, Sanchez DJ, Aliyari R, Lu S, Cheng G. Systematic identification of type I and type II
13 interferon-induced antiviral factors. *Proc Natl Acad Sci U S A* 2012; **109**:4239–4244.
- 14 5 Hycza MD, Kovacs C, Loutfy M, Halpenny R, Heisler L, Yang S, *et al.* Distinct
15 Transcriptional Profiles in Ex Vivo CD4+ and CD8+ T Cells Are Established Early in Human
16 Immunodeficiency Virus Type 1 Infection and Are Characterized by a Chronic Interferon
17 Response as Well as Extensive Transcriptional Changes in CD8+ T Cells. *J Virol* 2007;
18 **81**:3477–3486.
- 19 6 Sedaghat AR, German J, Teslovich TM, Cofrancesco J Jr, Jie CC, Talbot CC Jr, *et al.* Chronic
20 CD4+ T-cell activation and depletion in human immunodeficiency virus type 1 infection:
21 type I interferon-mediated disruption of T-cell dynamics. *J Virol* 2008; **82**:1870–1883.
- 22 7 Rotger M, Dang KK, Fellay J, Heinzen EL, Feng S, Descombes P, *et al.* Genome-wide mRNA
23 expression correlates of viral control in CD4+ T-cells from HIV-1-infected individuals.
24 *PLoS Pathog* 2010; **6**:e1000781.
- 25 8 Campbell JH, Hearps AC, Martin GE, Williams KC, Crowe SM. The importance of
26 monocytes and macrophages in HIV pathogenesis, treatment, and cure. *AIDS*
27 2014;**28**:2175-87
- 28 9 Wu JQ, Sassé TR, Wolkenstein G, Conceicao V, Saksena MM, Soedjono M, *et al.*
29 Transcriptome analysis of primary monocytes shows global down-regulation of genetic
30 networks in HIV viremic patients versus long-term non-progressors. *Virology* 2013;
31 **435**:308–319.
- 32 10 Krishnan S, Wilson EMP, Sheikh V, Rupert A, Mendoza D, Yang J, *et al.* Evidence for
33 innate immune system activation in HIV type 1-infected elite controllers. *J Infect Dis*
34 2014; **209**:931–939.
- 35 11 Noel N, Boufassa F, Lécuroux C, Saez-Cirion A, Bourgeois C, Dunyach-Remy C, *et al.*
36 Elevated IP10 levels are associated with immune activation and low CD4⁺ T-cell counts in
37 HIV controller patients. *AIDS* 2014; **28**:467–476.

- 1 12 Riveira-Muñoz E, Ruiz A, Pauls E, Permanyer M, Badia R, Mothe B, *et al.* Increased
2 expression of SAMHD1 in a subset of HIV-1 elite controllers. *J Antimicrob Chemother*
3 2014; **69**:3057–3060.
- 4 13 Abdel-Mohsen M, Raposo RAS, Deng X, Li M, Liegler T, Sinclair E, *et al.* Expression profile
5 of host restriction factors in HIV-1 elite controllers. *Retrovirology* 2013; **10**:106.
- 6 14 de Masson A, Kirilovsky A, Zoorob R, Avettand-Fenoel V, Morin V, Oudin A, *et al.* Blimp-1
7 overexpression is associated with low HIV-1 reservoir and transcription levels in central
8 memory CD4+ T cells from elite controllers. *AIDS* 2014;**28**:1567-77
- 9 15 Clerzius G, Gélinas J-F, Gatignol A. Multiple levels of PKR inhibition during HIV-1
10 replication. *Rev Med Virol* 2011; **21**:42–53.
- 11 16 Diamond MS, Farzan M. The broad-spectrum antiviral functions of IFIT and IFITM
12 proteins. *Nat Rev Immunol* 2013; **13**:46–57.
- 13 17 Haller O, Kochs G. Human MxA protein: an interferon-induced dynamin-like GTPase with
14 broad antiviral activity. *J Interferon Cytokine Res Off J Int Soc Interferon Cytokine Res*
15 2011; **31**:79–87.
- 16 18 Jacquelin B, Mayau V, Targat B, Liovat A-S, Kunkel D, Petitjean G, *et al.* Nonpathogenic
17 SIV infection of African green monkeys induces a strong but rapidly controlled type I IFN
18 response. *J Clin Invest* 2009; **119**:3544–3555.
- 19 19 Lu J, Pan Q, Rong L, He W, Liu S-L, Liang C. The IFITM proteins inhibit HIV-1 infection. *J*
20 *Virology* 2011; **85**:2126–2137.
- 21 20 Compton AA, Bruel T, Porrot F, Mallet A, Sachse M, Euvrard M, *et al.* IFITM Proteins
22 Incorporated into HIV-1 Virions Impair Viral Fusion and Spread. *Cell Host Microbe* 2014;
23 **16**:736–747.

24

25


1 **FIGURE LEGENDS**

2

3 **Fig. 1.** Expression of ISGs between patient groups and cell types. (a) IFN-induced
4 transmembrane protein 1 (IFITM1). (b) RNA-activated Protein Kinase (PKR). (c) IFN-induced
5 protein with tetratricopeptide repeats (IFIT1). (d) Myxoma virus Resistance protein A (MxA).
6 HDs, healthy donors; HICs, HIV controllers; ARTs, aviremic ART-treated patients; VIRs,
7 viremic treatment naïve patients.

8

1 **FIGURE**
 2 **Figure 1.**



3

1 **SUPPLEMENTAL MATERIAL**
2

1 **FIGURE LEGEND**


2 **Supplemental Figure 1.** Relationships with the *ifitm1* expression and immune activation
3 marker (HLA-DR and CD38 for CD4 and CD8 T cells and level of sCD14 for the monocytes)
4 (a) Considering all HIV-1 infected patients. (b) Considering HICs separately. IFITM1, IFN-
5 induced transmembrane protein 1; HICs, HIV controllers; ARTs, aviremic ART-treated
6 patients; VIRs, viremic treatment naïve patients.

7


1 Supplemental Figure 1

2

a)



b)



3

4

1
2
3
4
5

SUPPLEMENTAL FIGURES/ TABLES

Supplemental Table 1. Characteristics of study participants

1 **Supplemental Table 1**

Supplemental Table 1. Characteristics of study participants					
Characteristics	HDs (n=8)	HICs (n=11)	ARTs (n=8)	VIRs (n=8)	p
Age (years)	46 [43.75-48.5]	46.5 [42.25-50.75]	47 [43-53]	37 [32-54]	0.6
Male, <i>n</i> (%)	3 (37)	4 (36)	4 (50)	3 (37)	0.7
CD4+ T-cell count at enrollment (cells/ μ L)	-	875 [768-1001]	467 [430-550]	100 [46-184]	<0.01
Lowest CD4+ T-cell count since HIV diagnosis (nadir) (cells/ μ L)	-	523 [464.5-637]	285 [272-330]	100 [46-184]	<0.01
Plasma HIV RNA VL at enrollment (log ₁₀ copies/ml)	-	1.6 [<1.6 -1.6]	<1.6 [<1.6 - <1.6]	4.9 [4.4-5.35]	<0.01
HCV coinfection, <i>n</i> (%)	0 (0)	0 (0)	0 (0)	1 (14)	0.3
HBV coinfection, <i>n</i> (%)	0 (0)	0 (0)	0 (0)	2 (28)	0.06
Active opportunistic infection at enrollment, <i>n</i> (%)	-	0	0	5 (62)	<0.01
Results are shown as median (IQR) or <i>n</i> (%). HDs, healthy donors; HICs, HIV controllers; ARTs, aviremic ART-treated patients; VIRs, viremic treatment naïve patients ; VL, viral load					