

HIV controllers : a genetically determined or inducible phenotype?

Asier Saez-Cirion, Gianfranco Pancino

► **To cite this version:**

Asier Saez-Cirion, Gianfranco Pancino. HIV controllers : a genetically determined or inducible phenotype?. Immunological Reviews, Wiley, 2013, 254 (1), pp.281–294. <10.1111/imr.12076>. <pasteur-01420535>

HAL Id: pasteur-01420535

<https://hal-pasteur.archives-ouvertes.fr/pasteur-01420535>

Submitted on 16 May 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HIV CONTROLLERS: A GENETICALLY DETERMINED

OR INDUCIBLE PHENOTYPE?

Asier Sáez-Cirión¹ and Gianfranco Pancino^{1,2*}

1. Institut Pasteur, Unité de Régulation des Infections Rétrovirales, Paris, France, 2.

INSERM, Paris, France

- *Corresponding author : Gianfranco Pancino, Unité Régulation des Infections Rétrovirales, Institut Pasteur, 28, rue du Docteur Roux, 75724 Paris cedex 15
Tel : +33145688768, Fax : +33145688957, e-mail : gianfranco.pancino@pasteur.fr*

Running title : HIV Controllers**Summary**

1 Less than 0.5% of HIV-1 infected patients maintain viral load below the threshold of
2 detection by conventional assays for many years without antiretroviral therapy. These
3 patients are known as HIV controllers (HICs) and offer a unique opportunity to explore and
4 unravel efficient mechanisms of defense against HIV. An intense research effort has led to
5 identify viral and host factors that contribute to the control of viral replication. Although the
6 host factors appear to be the primary determinants for HIV control, different genomic,
7 transcriptomic and immunological profiles have been described in HICs. This supports the
8 existence of different mechanisms of control. We review here the most significant data in
9 the field, including our own work, in the light of a crucial question: whether the ability to
10 achieve and maintain the control of HIV replication is linked to genetic or immunological
11 features specific to HIC or can be induced by vaccine or therapeutic approaches in the
12 general population.

13

14 Keywords: HIV-1, HIV-controllers, Post-treatment controllers, antiviral responses, viral
15 reservoirs

16

1 **Introduction**

2 The pathogenesis of HIV infection in humans depends on a balance of viral and host factors.
3 The fitness of the virus and the type of cell that it is able to infect may have an enormous
4 impact on the rate of progression to disease (1-3). The virus relies on numerous cellular
5 factors to complete its replication cycle, and it needs to altogether overcome intrinsic cell
6 factors that are devised to hinder viral replication (4-6). Moreover, the host mobilizes a
7 panoply of immune responses, both innate and adaptive, to counteract infection. A better
8 understanding of the intimate interactions between the virus and the host will undoubtedly
9 provide important clues in the search for an HIV vaccine and cure. Interestingly, in a few
10 patients, the host factors appear to have prevailed over the virus, as these patients are able
11 to naturally control viral replication, maintaining extremely low virus levels (7-11). These
12 patients are known as HIV controllers (HICs) and offer a unique opportunity to explore and
13 unveil efficient mechanisms of defense against HIV.

14

15 **Clinical characteristics**

16 Transitory control of infection may occur in up to ~7% of all HIV-infected patients (12).
17 However, in most of these patients, control of the infection does not last long and is lost
18 within less than 12 months. In contrast, once control of viremia has been maintained for at
19 least one year, the probability of these patients maintaining undetectable viral replication
20 for numerous years is increased. Overall, the frequency of these long-term HIV controllers is
21 low. Several epidemiological studies have estimated that these patients may represent
22 around 0.5% of the total population of HIV-infected patients (9, 13, 14). The definition of
23 HICs varies worldwide depending on the threshold of maximal viral replication that is used or

1 the length of control (15). In the French ANRS CO21 cohort, HICs are defined as patients
2 naïve to antiretroviral therapy (except for transient treatments during pregnancy) who have
3 been infected for at least five years and have maintained their five last viral loads below 400
4 HIV RNA copies per ml of plasma. The 400 RNA copy threshold was chosen for historical
5 reasons, as this was the threshold of the first viral load tests largely available in French
6 clinical centers in the 90s. Other cohorts, such as the international HIV controller
7 consortium, have defined these patients as HIV-infected individuals who have maintained at
8 least two consecutive plasma viral loads below 50 copies/ml within the last 12 months
9 without therapy (10).

10

11 Patients who will become HICs typically have a non-symptomatic primary infection and have
12 higher CD4+ T cell levels and lower plasma viral loads and reservoir levels in the blood
13 compared to those who will not achieve control of infection (12, 16). This suggests that
14 efficient mechanisms of control start acting soon after infection in most of these patients (9).
15 However, the control of infection in some particular cases is a progressive phenomenon that
16 may require several years to accomplish (9). What differentiates these late controllers from
17 early controllers remains unknown, but comparative studies may provide interesting insights
18 into the development and relative contribution of the different mechanisms that may
19 participate in establishing control of infection in HICs. Little information is available
20 regarding co-infections by other pathogens in HICs, but at least some of these patients
21 appear to be able to control or even clear hepatitis C infection (17, 18).

22

1 Although HICs maintain a remarkable control of infection, blips in viral load levels have been
2 observed for many of these individuals during long-term follow-up (13). Very low levels of
3 plasma virus can be detected by ultrasensitive RT-PCR assays in HICs, revealing a persistence
4 of viral replication despite maintaining viremia below the limit of detection by standard RT-
5 PCR (19-21). Interestingly, these patients may have been able to control infection of the
6 central nervous system even more effectively, as detection of viral replication is infrequent
7 in cerebrospinal fluid from these patients, even when using ultrasensitive techniques (22,
8 23). Many HICs also experience a decline of their CD4+ T cell levels over time (13, 24). It is
9 likely that the loss of CD4+ T cells in HICs is related to low-level viremia; we and others have
10 recently reported that declining CD4+ T cell levels were observed in patients who had
11 experienced blips in their viral loads or had higher low-level viremia (13, 21). In contrast, for
12 some HICs, a viral blip has never been recorded over very long follow-up periods (13). These
13 patients maintain stable CD4+ T cell levels, and many have undetectable viral loads even
14 when using ultrasensitive techniques able to detect 1 RNA copy/ml of plasma (our
15 unpublished observations). Interestingly, these HICs are close to healthy individuals from a
16 transcriptomic point of view (25).

17

18 The slopes of CD4+ loss in HICs are typically very modest (13). However, AIDS-related events
19 have been reported for a couple of HICs (24), and CD4+ T cell losses have prompted
20 therapeutic intervention in a few others (26, 27). Although data on such patients and their
21 clinical evolution are scarce, the reaction of HICs to antiretroviral treatment appears to be
22 unequal. Some regain their CD4+ T cell counts, although their gains are smaller than those of
23 patients with uncontrolled viremia who start therapy (26). Other patients do not increase

1 their CD4+ T cell numbers despite exhibiting a reduction in their low-level viremia upon
2 therapy initiation (27). This would suggest that additional factors underlie the evolution of
3 CD4+ T cell counts in HICs, and their identification may help to devise additional therapeutic
4 approaches to implement in these HICs and other HIV-infected patients as a complement to
5 antiretroviral therapy.

6

7 CD4+ T cell loss in HICs has been associated with elevated levels of T cell activation (24).
8 Although lower compared to viremic patients, most HICs have abnormally high levels of T
9 cell activation and inflammatory markers and signs of microbial translocation when
10 compared to the healthy population or, for some markers, HIV-infected patients receiving
11 antiretroviral therapy (24, 28, 29). Indeed, high levels of immune activation may well be as
12 important a determinant of CD4+ T cell loss in HICs as they are in uncontrolled infection.
13 However, high levels of immune activation are a rather common characteristic observed in
14 HICs and do not entirely explain the more profound loss of CD4+ T cell counts in the rare
15 cases presented above.

16

17 As in other HIV-infected patients, HICs have higher proportions of differentiated T cell
18 subsets compared to healthy donors (30). Two recent reports suggest that HICs can
19 regenerate T cells through optimal thymic output, homeostatic proliferation and preserved
20 lymphopoiesis (30, 31). In contrast, HICs experiencing significant progression have reduced
21 levels of hematopoietic progenitor cells (31), which likely compromises thymic function in
22 these patients (30). Thus, there may be some cost for HICs to maintain an optimally tuned

1 immune response against HIV (see below), but the benefits overwhelm the disadvantages
2 for the vast majority of HICs.

3

4 **Host-virus interactions involved in HIV control**

5 **Viral determinants**

6 The low levels of viremia in HICs are similar to those found in HAART-treated patients (19,
7 21, 32), and a median value of 2 copies/ml in a cohort of HICs has been reported (33). In
8 contrast, HIV-1 reservoirs are much smaller in HICs than in patients on HAART (7, 34, 35).
9 This discrepancy raises the question of whether plasma viruses in HICs originate from
10 reactivations of this decreased latent reservoir or correspond to ongoing viral replication.
11 Cross-sectional studies of plasma and cell-associated virus sequencing from HLA-B*57-
12 positive HICs have shown discordance between *gag* plasma viral sequences and sequences
13 from archived provirus in resting CD4+ T cells (32). All the amplified plasma viral clones had
14 mutations in HLA-B*57-restricted epitopes, which is consistent with selective pressure from
15 the CTL response. In contrast, almost all proviral clones were wild type, suggesting that the
16 infecting founder viruses did not contain escape mutations. Strong evidence indicating that
17 persistent viremia in HICs is sustained by continuous cycles of viral replication comes from
18 longitudinal studies in which gene sequencing was performed in sequential samples. The
19 analysis of *gag* and *nef* sequences from the plasma of HLA-B*57- and/or HLA-B*27-positive
20 HICs showed a temporal evolution of viral genes (36, 37). Interestingly, genetic evolution
21 was primarily restricted to the plasma viruses and was rare in cell-associated viral
22 sequences. *Gag* and *nef* proviral clones appeared to be ancestral to the plasma clones and
23 likely represent archived viruses. This suggests that viral reseeding and expansion of the

1 latent reservoir is not allowed by the very low rate of HIV-1 replication in the cellular
2 compartments of HICs. While mutations found in HLA-B*57/5801- and HLA-B*27-restricted
3 epitopes persisted over time, suggesting a continuous CTL-driven pressure, the majority of
4 the evolution observed in the *gag* and *nef* genes in the patients included in these studies
5 was driven by synonymous mutations and not by evolution in CTL-targeted epitopes (36). A
6 lower rate of viral replication after an initial control of viral replication by immune responses
7 may explain the lower frequency of nonsynonymous mutations and CTL escape variants in
8 HICs compared to chronic viremic patients. Single-genome sequencing followed by a
9 phylogenetic analysis of the *pro-rt* and *env* genes from plasma samples also showed that
10 sequence divergence in these genes increases with time in many HICs (20). In this study,
11 evolution of plasma viruses was found both in patients who did not have protective HLA
12 alleles and in patients carrying the HLA-B*57/27 alleles, indicating ongoing virus replication
13 despite the presence of these protective HLA alleles. However, in accordance with the
14 results reported for *gag* sequences (36), significantly fewer mutations within known CTL
15 epitopes in Pro and RT were found in HICs than in noncontroller patients (20).

16 Strong cellular response-driven immune selection has been associated with a reduced
17 replication capacity of mutant viruses (38-40). Accordingly, chimeric HIV-1 viruses containing
18 the *gag-pol*, *rev*, *vif*, and *env* genes isolated from the plasma of HICs showed reduced
19 replicative capacity compared with viruses containing the corresponding genes isolated from
20 patients with progressive disease (41, 42), (43, 44). In addition, the reduced viral fitness of
21 *gag-protease* clones derived from HICs was correlated with mutations in epitopes restricted
22 by protective HLA alleles (40, 42). A recent study documented a relative impairment of Nef
23 functions in Nef clones isolated from HICs compared to those from chronic progressors (45).
24 Noncanonical B57*-associated polymorphisms in CTL epitopes, different from the common

1 polymorphisms described in progressors, have been associated with Gag-Pol and Nef
2 attenuations in HICs (40, 45). These results suggest that pressure by HLA-B57* CTLs selects
3 for rare variants in HICs leading to lower viral replication capacity. This is well illustrated by a
4 study that analyzed the common HLA-B57/B*5801 associated T242N mutation in the Gag
5 TW10 epitope found in B57/B*5801-positive HICs and HIV progressors. Rare variants that
6 reduced the viral replicative capacity were found exclusively in HICs within and flanking the
7 TW10 epitope; these variants were recognized by variant-specific CTL responses (40). These
8 studies support a reduction of viral fitness associated with CTL escape mutations in chronic
9 infection. The open question is whether infection with a mutated attenuated virus may be at
10 the origin of the control of viremia in HICs or whether viral attenuation is caused by the
11 strong immune response that leads to the control of viremia. The evolution in HLA-
12 B*57/B*27-restricted epitopes found in plasma viruses but not in the cell-associated viruses
13 of HICs, as mentioned above, is in favor of the second hypothesis. Obvious genetic defects,
14 such as deletions, insertions and stop codons, have not been found in viruses isolated from
15 HICs, except in sporadic cases ((46) and references herein). However, one study has
16 described an impaired replicative capacity of plasma gag-protease clones obtained during
17 the acute/early phase of infection from patients who subsequently become controllers (47).
18 In some cases, HICs were recipients of drug-resistant strains or of strains containing B57-
19 associated escape mutations, suggesting the transmission of attenuated viruses; others
20 carried HLA-protective alleles that could have forced the selection of escape mutations in
21 early phases of infection. These data suggest that the characteristics of the infecting viruses
22 and an early selection of fitness cost mutations by protective HLA alleles are conditions
23 surrounding viral control in HICs. However, many HICs do not carry protective alleles, and
24 mechanisms of viral control that are dependent on immune selection cannot be generalized

1 to all the patients. Moreover, all the studies performed with chimeric viruses focused on
2 isolated genes introduced into a reference heterologous backbone strain and did not
3 address the replicative potential of whole viral isolates. Therefore, if the weakened function
4 of viral proteins appears to be frequent in HICs, this does not exclude the possibility that
5 viruses infecting HICs can exhibit replicative fitness. Indeed, replication-competent viruses
6 were isolated from the plasma of HICs from different cohorts, including the French ANRS
7 cohort, and showed replication levels similar to those of reference HIV-1 strains (34, 48-50).

8 Overall, some important points can be inferred from these studies: infection with an
9 attenuated/defective virus can make achieving control of viral replication easier in some
10 HICs, but it alone does not explain the HIC phenotype. Indeed, control of infection has been
11 reported in patients receiving virus from pairs with progressive infection, suggesting that
12 host factors are the most important determinant for HIV control (51). When present,
13 protective HLA alleles help to achieve and/or maintain control of viremia but still require the
14 presence of other factors. Indeed, HICs and progressors carrying the same HLA allele display
15 different viral polymorphisms and sequence evolution. In addition, different outcomes of
16 transmitted viruses have been reported in HLA-B*57-positive HICs and progressors; viruses
17 from progressors showed compensatory mutations and became progressively more fit over
18 time, whereas viruses from HICs did not (52),(53).

19

20 **Genetic determinants of the innate and adaptive immune response**

21 A number of studies have attempted to understand the contribution of genetic factors and
22 of immune responses to viral control. Genome wide association studies (GWAS) have located
23 the genetic determinants with the strongest association with HIV-1 control in the HLA class I

1 region in the major histocompatibility complex (MHC) on chromosome 6 (54, 55). The single
2 nucleotide polymorphisms (SNPs) with the highest significant association were found in
3 close proximity to the HLA-B and HLA-C genes, which is consistent with earlier reports
4 showing an association between specific HLA class I alleles and disease progression (56). The
5 strongest association was found with a SNP located in the HCP5 gene that is in linkage
6 disequilibrium with the HLA-B*5701 allele in whites and with HLA-B*5703 in African-
7 Americans (57, 58). A deeper analysis by inferred sequencing of the HLA class I alleles in
8 conjunction with GWAS showed that the specific amino acid composition in the HLA B
9 peptide-binding groove underlies the associations between protection or disease
10 progression and protective alleles such as B*27, B*57 and B*14 and deleterious alleles such
11 as B*35, respectively (59). The second association with HIV-1 control was found with a SNP
12 35 kb upstream from the HLA-C gene (54, 59). This SNP was also associated with higher
13 expression of HLA C alleles on the surface of cells (60). This effect on HLA C expression has
14 been subsequently explained by a differential binding of microRNA Hsa-miR-148a to its
15 target site within the 3'untranslated region of HLA-C, which leads to a specific post-
16 transcriptional regulation of different HLA-C alleles (61).

17 It is important to note that the HLA genetic associations together explain 19% of the
18 variance of host control (59), indicating that other factors are involved in determining the
19 HIC phenotype. However, it is not totally surprising that GWAS have identified only the
20 genetic determinants located in the MHC region, considering the low frequency of HICs (~
21 0.5% of HIV-1-infected individuals) and their heterogeneity. Indeed, only common variants
22 can be detected by linkage in a population, and standard GWAS do not have the power to
23 detect rare variants (62).

1 The strong genetic associations in the MHC region point to the role of immune responses in
2 HIV-1 control at least in some HICs. The associations between specific HLA class I alleles, HIV-
3 1 control and, in particular, the impact of the peptide composition of the class I binding
4 groove on viral peptide-binding capacity suggest a direct role for the specific cytotoxic T
5 lymphocyte (CTL) response. However, HLA class I molecules are also ligands for NK cell killer
6 cell immunoglobulin-like receptors (KIRs). Certain KIRs, namely KIR3DL1 and KIR3DS1, have
7 been found to be associated with better control of HIV-1 when found in patients that also
8 have HLA-B alleles with Bw4 epitope specificity (Bw4-80I) (63). In line with this observation,
9 NK cells from individuals carrying KIR3DL1 receptor-HLA-Bw4 ligand pairs perform multiple
10 functions in response to stimulation (64). The HLA-B alleles that have been associated with
11 HIV-1 control (HLA-B*57/27/58) are of the Bw4-80I group, suggesting that the mechanisms
12 of viral control may not only be due to CTL responses but also to the induction of an efficient
13 NK cell response through their interaction with NK cell receptors.

14 However, data regarding the impact of NK cells on HIV-1 control in HICs are scarce. The
15 combination of the KIR3DL1*h/*y genotype and HLA-B*57 has been associated with
16 protection from HIV-1 disease progression (65). NK cells from subjects carrying the
17 *h/*y+B*57 genotypes exhibit the highest polyfunctional potential, but this is dependent on
18 cocarriage of the NK receptor and its HLA-Bw4*80I ligand (64). One study reported increased
19 target cell-induced degranulation and cytokine secretion following IFN α stimulation in NK
20 cells from HICs carrying the inhibitory KIR3DL1*h/*y receptor genotype and the
21 corresponding HLA-Bw4*80I ligand (66). However, NK cells from HICs show a modest
22 capacity to suppress viral replication in autologous CD4+ T cells *in vitro*, much lower than
23 that mediated by CD8(+) T cells ((67), and our unpublished results).

24

1 **Adaptive responses**

2 Cytotoxic CD8+ T lymphocytes

3 In contrast to the NK response, the HIV-1-specific CD8+ T cell response in HICs has been the
4 focus of intense investigation. Despite the extremely low level of viral antigen, many HICs,
5 whom we have termed “strong responders”, present a high frequency of HIV-1-specific CD8+
6 T cells that is much higher than in HAART patients, despite having comparable viral loads (33,
7 49, 68). However, the magnitude of this response, measured by IFN gamma ELISPOT assay, is
8 similar or even lower than in viremic progressors (10, 68), suggesting that viral control
9 implies qualitative differences not revealed by this assay. Several phenotypic markers are
10 differentially expressed on HIV-1-specific CD8+ T cells from HICs in comparison with viremic
11 progressors, including markers of T cell activation/exhaustion, such as PD-1 and CD57 (69,
12 70). A peculiar trait of HIV-1-specific CD8+ T cells from HICs is the discordant expression of
13 the activation markers HLA-DR and CD38. Whereas HIV-1-specific CD8+ T cells from viremic
14 progressors express high levels of the two markers and those from HAART patients express
15 low levels of both, HIC CD8+ T cells express high levels of HLA-DR and low levels of CD38
16 (68). This profile was previously described for the entire CD8⁺ T cell population in
17 asymptomatic HIV-infected patients with stable CD4⁺ T cell counts (71). This differential
18 expression might be linked to a superior capacity to respond to antigenic stimulation in the
19 absence of nonspecific activation driven by the inflammatory context present in viremic
20 patients. Accordingly, HLA-DR expression was correlated with the percentage of proliferating
21 cells among HIV-specific CD8⁺ T cells (68). A higher ability of HIV-specific CD8 T cells from
22 HICs to proliferate in response to HIV-infected CD4+ T cells or HIV-1 peptides in comparison
23 with viremic patients has been consistently found and associated in many studies with the
24 protective HLA-B*57 allele (8, 68, 72). The level of polyfunctional HIV-specific CD8+ T cells,

1 defined by the capacity to exert multiple effector functions, including the secretion of
2 several cytokines, such as IFN gamma, TNF- α , MIP-1 β , CD107a and IL-2, following antigenic
3 stimulation, is inversely correlated with viral load (73). A polyfunctional response is found in
4 HICs and is scarce or absent in viremic progressors, most likely as a consequence of
5 persistent exposure to antigen during viremic infection (73, 74). The functional property that
6 most clearly differentiates the HIV-specific CD8+ T cells of HICs from those of viremic
7 progressors and HAART-treated patients is their capacity to directly kill infected cells *in vitro*.
8 HIV-specific CD8+ T cells from HICs have been shown to effectively suppress viral replication
9 in autologous CD4+ T cells by eliminating HIV-infected cells without the need for exogenous
10 stimulation (68). This suppressive capacity is not observed in either viremic progressors or in
11 HAART-treated patients and has been associated with an effective loading of perforin and
12 granzyme B in CD8+ T cells from controllers upon contact with HIV-infected cells (75, 76). Of
13 note, whereas more efficient loading of lytic granules has been documented in CD8+ T cells
14 after 6 days exposure to antigen (76), the killing of infected cells is detected a few hours
15 after adding CD8+ T cells to infected CD4+ T cells (68). This may be related to the ability of
16 CD8+ T cells from HICs to rapidly upregulate perforin and granzyme B following exposure to
17 antigen, an ability that has been associated with high levels of expression of the T-box
18 transcription factor T-bet after the *in vitro* expansion of HIV-specific CD8+ T cells (77). The
19 rapid elimination of infected cell targets by *ex vivo* CD8+ T cells implies the presence of CTLs
20 with immediate effector functions. Consistent with these results, differentiated effector
21 CD8+ T cells directed against HIV-1 are found more frequently in HICs compared to HIV-1
22 progressors (68, 78).

23 CD8+ T cell suppressive capacity in HICs is highly correlated with the anti-Gag response, not
24 with Nef- or Env-specific responses (49, 50, 79, 80), which is consistent with earlier studies

1 reporting an association between the anti-Gag CD8⁺ T cell response and viral control (81,
2 82). A direct contribution to the control of the viral reservoir by the Gag-specific CD8⁺ T cell
3 response in HICs is suggested by their association with a low HIV reservoir in the central
4 memory CD4⁺ T cells of HICs carrying HLA-B* 57/27 alleles (83).

5 An increased magnitude and polyfunctionality of CD8⁺ T cell responses to HIV Gag has also
6 been found in the rectal mucosae of HICs and was more strongly associated with controller
7 status than peripheral blood responses (84, 85). It has been suggested that a superior T cell
8 response in the intestinal mucosae may contribute to limit CD4⁺ T cell depletion and
9 preserve mucosal integrity (84). The superior antiviral activity of Gag-specific CTLs may be
10 related not only to their lytic function but also to the selection of immune escape variants
11 with lower replication capacity (40-42, 86, 87), as discussed above.

12 Several studies suggest that Gag-specific CTLs from HICs mediate effective viral control
13 because of their higher functional avidity, i.e., the ability to react to lower antigen
14 concentrations and to recognize more epitope variants than those from HIV progressors (88,
15 89). Accordingly, control of HIV infection has been associated with the selection of HLA-
16 B*27-restricted KK10 epitope-specific CD8⁺ T cell clonotypes with high avidity,
17 polyfunctionality and suppressive capacity (90). A major step in our understanding of the
18 mechanisms underlying the superior quality of the CD8⁺ T cell response in HICs came from
19 two recent studies that compared CD8⁺ T cell responses to Gag-immunodominant viral
20 epitopes in HICs and chronic progressors bearing HLA-B*27 or B*57 (91, 92). The ability of
21 CD8⁺ T cells targeted to the same epitopes to inhibit viral replication was higher in HICs than
22 in HIV progressors and was associated with a different selection of epitope-specific TCR
23 clonotypes during infection. The TCR clonotypes of HICs displayed a better capacity and
24 higher avidity to cross-recognize naturally occurring viral variants (92) and upregulate

1 perforin and granzyme B more than the TCR clonotypes of progressors (91). Interestingly,
2 the selection of these clonotypes occurs very rapidly after infection in controller patients
3 and in the context of low-level viremia (92). These studies provide an explanation for why
4 only a few individuals bearing the HLA-B* 27 and B*57 alleles control HIV-1 infection and
5 links the protective effect of these alleles to peculiar TCR rearrangements that lead to the
6 selection of the most effective clonotypes.

7

8 How favorable TCR clonotypes are selected in HICs remains unknown. The peculiar amino
9 acid composition of the protective HLA B peptide-binding groove (59) may favor an optimal
10 presentation of the viral peptide. It has also reported that the myeloid dendritic cells of HICs
11 efficiently generate T cell responses and that this capacity is associated with increased
12 expression of the leukocyte immunoglobulin-like receptors LILRB1 and LILRB3 (93). However,
13 additional studies will be needed to clarify these issues.

14 Polyfunctional and effective HIV-specific CD8+ T cells in strong-responder controllers may
15 result from the maintenance of an efficient response developed during the primary infection
16 that is lost in HIV progressors due to immune activation-driven T cell exhaustion. We
17 addressed this question in a study on a large number of patients during primary infection
18 who were included in a clinical assay (Optiprim) before the onset of HAART (94). We found
19 that most patients exhibited high frequencies of polyfunctional HIV-specific CD8+ T cells, but
20 these cells lacked strong HIV-suppressive capacities *ex vivo*. This indicates that HIV-
21 suppressive capacity is not a common characteristic of the CD8+ T cell response during
22 primary infection and suggests that it is an intrinsic feature found in HICs. This also implies
23 that this quality of the T cell response is a cause and not a consequence of viral control.

24

1 If the presence of protective alleles and an effective CTL response certainly play a major role
2 in viral control in many HICs, particular MHC alleles or a strong CD8+ T cell response are not
3 necessary to achieve controller status. Indeed, as discussed above, HICs are a heterogeneous
4 population: many HICs lack protective HLA alleles and/or have weak HIV-specific CD8⁺ T-cell
5 responses (10, 49, 95). A potential role of CD8+ T cells in viral control that is also observed in
6 HICs who have a low frequency of specific CD8+ T cells has been suggested by recent reports
7 showing that specific CD8+ T cells with viral suppressive capacity can be expanded upon
8 stimulation with Gag peptides, consistent with the presence of HIV-specific CD8+ T cells with
9 a memory phenotype (96, 97). In an *in vitro* suppressive assay, sorted effector memory and
10 terminal effector HIV-specific CD8+ T cells exhibited the strongest suppressive capacity in the
11 first days of CD8+/CD4+ T cell co-cultures. However, central memory CD8 T cells that
12 expanded after 5 days of incubation with antigen also acquired an effective suppressive
13 capacity (96). If these studies suggest that the presence of a few specific central memory
14 CD8+ T cells may be sufficient to maintain the control of eventual reactivations of HIV-1
15 replication in weak-controller HICs, this conclusion should be moderated by the following
16 considerations: the study of Buckheit included only HLA-B* 57-positive patients and
17 therefore may concern qualitative features of the CTL response linked to this protective
18 allele; other studies have reported that pre-stimulation of CD8+ T cells from HAART patients
19 with HIV antigen also leads to the efficient killing of infected cells and that the primary
20 difference with CD8+ T cells from HICs is their rapid viral suppressive activity (98).

21

22 Regulatory T cells

23 Several reports, if not all, have reported that the frequency of regulatory T cells (Tregs) is not
24 increased in HICs and is similar or lower compared to uninfected individuals (99-102). This

1 may contribute to the maintenance of high-quality HIV-1-specific CD8+ T cell responses in
2 HICs. The cost may be the persistent immune activation found in most HICs despite
3 controlled viremia (see previous section) (24).

4

5 Humoral response

6 As in the case of the CD8+ T cell response, HICs also have a heterogeneous humoral
7 response. As could be expected due to their low levels of viremia, neutralizing antibodies are
8 present at low levels in HICs and are unlikely to play a major role in the control of infection
9 (10, 103). In contrast, ADCC potential, both related to the quality of non-neutralizing
10 antibodies and the cell surface expression of Fcγ receptors, appears to be greater in
11 controllers (103-105). Nevertheless, the weakest antibody levels are found among HICs with
12 the best control of infection (no viral blips, no CD4+ T cell loss), and weakly reactive/partial
13 western blots have been reported for some of these HICs (106).

14

15 CD4+ T cell response

16 CD4+ T cell help is required to sustain an effective CD8+ T cell response in chronic viral
17 infections, and numerous studies have addressed the characteristics and the potential anti-
18 viral role of CD4+ T cells in HICs. GWA and transcriptomic analyses of CD4+ T cells,
19 comparing elite controllers with patients successfully treated with HAART and healthy
20 donors, did not find a transcriptome profile characteristic of viral control (107). This study
21 demonstrated that viral replication was the major driver of RNA expression changes in HIV-
22 1-infected individuals. Consistent with that report, another study focusing on HLA-DR⁻ CD4+
23 T cell subsets showed that the transcriptional profiles of most elite controllers were similar
24 to those of successfully HAART-treated patients but different from those of HIV-1-negative

1 persons (25). Although it was not possible to identify transcriptional patterns specific of
2 HICs, phenotypic and functional profiles of HIV-specific CD4⁺ T cells indicated a higher
3 quality and superior survival in HICs compared to viremic or HAART-controlled patients, as
4 described for CD8⁺ T cells. A comparison between HICs and HAART-treated patients with
5 controlled viremia showed equivalent levels of expression of activation markers on the CD4⁺
6 T cells of HICs, but they were lower compared to the CD4⁺ T cells of viremic patients and
7 higher compared to those of healthy donors (24, 29, 108, 109). The CD4⁺ T cells of HICs have
8 lower levels of expression of the negative immunoregulatory molecule cytotoxic-T-
9 lymphocyte-associated antigen 4 (CTLA-4) compared to those of HAART patients, which is
10 consistent with the increased functionality of these cells (110). In contrast to viremic
11 patients, the CD4⁺ T cells from HICs maintain the ability to secrete IL-2 and proliferate upon
12 HIV-1 Gag stimulation, which is associated with a preserved central memory population (29,
13 111). Compared to viremic patients and healthy individuals, a lower level of FoxO3a-
14 mediated pro-apoptotic transcriptional activity in memory T cells from HICs has been
15 proposed as a molecular basis for the preservation of memory CD4⁺ T cells in HICs (112).
16 Memory T cells from HICs show high avidity for immunodominant Gag peptides and are
17 capable of maintaining robust responses despite the presence of only a low level of viral
18 antigen (113). Like CD8⁺ T cells, CD4⁺ T cells from HICs can secrete several
19 cytokines/chemokines simultaneously in response to antigen (29, 108). Polyfunctional CD4⁺
20 T cell responses associated with the HLA-DRB1*13 and HLA-DQB1*06 alleles were also found
21 in the gut mucosa of HICs (114). Consistent with high levels of IL-2⁺/IFN- γ ⁺ production by
22 HIV-specific CD4⁺ T cells, a high frequency of CD4⁺ Th1 effectors in response to
23 immunodominant Gag peptides has been reported in HICs (109). Taken together, these data
24 are consistent with the presence of HIV-specific, long-lived, highly functional memory CD4⁺ T

1 cells with the capacity to be activated and rapidly proliferate in response to viral reactivation
2 in HICs. These CD4+ T cells would provide a helper function for driving and sustaining the
3 antiviral CD8+ T cell response.

4 However, memory CD4+ T cells are the preferred target of HIV-1, and infected central and
5 transitional memory CD4+ T cells have been reported to be the major cellular compartments
6 of the latent HIV reservoir (115).

7

8 **Intrinsic immunity**

9 Some studies have reported that CD4+ T cells from HICs are relatively resistant to HIV-1
10 infection *in vitro* compared to CD4+ T cells from healthy donors or viremic patients (116-
11 118). Interestingly, the decreased susceptibility of CD4+ T cells correlates with the level of
12 viral DNA that these CD4+ T cells carry *in vivo*, suggesting that it may contribute to the
13 decrease of the viral reservoir in HICs (117). Decreased permissiveness to HIV-1 replication
14 was also found in the monocyte-derived macrophages of HICs and was related to the
15 inhibition of the accumulation of viral reverse transcripts in both CD4+ T cells and
16 macrophages (117). An additional inhibition at the level of HIV-1 transcription has been
17 described in CD4+ T cells (116). Viral restriction does not depend on the upregulation of
18 known retrovirus restriction factors, including *TRIM5 α* , *APOBEC3 family*, *BST2/Tetherin*, and
19 accordingly, transcriptomic data show that the expression of these genes is driven by an
20 increasing viral load (107) and our unpublished results). We and others have reported that
21 CD4+ T cells from HICs have increased *ex vivo* expression of the CDK inhibitor p21 compared
22 to uninfected individuals (116, 117). p21 has been previously shown to inhibit the
23 preintegrative steps of HIV-1 replication *in vitro* in macrophages and hematopoietic stem
24 cells (119, 120). One study demonstrated that p21 overexpression was involved in the

1 restriction of HIV-1 replication in activated CD4+ T cells (116); by contrast, we did not find a
2 direct role for p21 in HIV-1 restriction in CD4+ T cells or macrophages from HICs (117). The
3 permissiveness of target cells to HIV-1 remains a debated question because other authors
4 have found that CD4+ T cells from HICs and uninfected individuals have similar susceptibility
5 to HIV-1 (121). Interestingly, however, the study by O'Connell reported that CD4+ T cells
6 from HICs produced less virus per infected cell than CD4+ T cells from viremic progressors
7 (121), suggesting a further defect that could affect viral spread. A study conducted in a few
8 HICs reported hypermethylation of the HIV-1 proviral 5'-LTR in comparison with HIV
9 progressors (122), suggesting that epigenetic modifications may contribute to the control of
10 viral replication. This hypothesis is consistent with the observation that despite an effective
11 control of viremia and an extremely low cell-associated HIV-1 DNA level, infecting virus can
12 be isolated from CD4+ T cells from many HICs upon activation with mitogens, cytokines or
13 TCR stimulation (34, 48-50), suggesting that viral expression from latently infected cells may
14 be repressed *in vivo*.

15
16 In conclusion, virologic and/or genetic factors likely account for viral control in some HICs.
17 Adaptive immune responses, specifically polyfunctional CD8⁺ and/or CD4⁺ T-cell responses,
18 may limit viral replication, particularly in strong-responder HICs. HICs lacking protective MHC
19 alleles and/or having weak T cell responses may rely on other mechanisms, such as the
20 cellular restriction of viral replication and/or innate immune responses that have not been
21 clearly identified.

22

23 **Post-treatment controllers**

1 HICs are accepted as an appealing example of what a functional HIV cure may represent
2 (123). Other than to better understand how they control infection, the challenge now is to
3 devise ways to induce a similar phenotype of control in a larger group of patients and
4 irrespective of their MHC background. Some HIV patients diagnosed early in primary
5 infection who start antiretroviral treatment almost immediately and maintain treatment for
6 long periods of time (more than 1 year) are able to durably control viremia after treatment
7 discontinuation (124-127). Because of the rapid therapeutic initiation, one of the difficulties
8 when studying these post-treatment controllers (PTCs) is to exclude the possibility that they
9 may have also been able to control infection without any intervention. Although data on
10 these patients are still scarce, many differences have been observed between PTCs and
11 natural HICs. Whereas natural control of viremia is generally associated with non-
12 symptomatic primary infection, lower viral loads and high CD4+ T cell numbers (12, 16), most
13 PTCs have a highly symptomatic primary infection with high viral loads and low CD4+ T cell
14 numbers at levels comparable to patients who will not control the infection (127). Moreover,
15 protective HLA alleles are not overrepresented among PTCs, who show a significant
16 frequency of some HLAs that are rarely found in HICs and are associated with higher viral
17 loads (127). Overall, PTCs have lower levels of T cell activation and much weaker and less
18 efficient CD8+ T cell responses than HICs. Additionally, the probability of controlling infection
19 for at least one year after the interruption of treatment initiated close to the primary
20 infection and maintained for at least one year is estimated to be 8-15% (124-127). This
21 estimation is much higher than that for > 1 year of natural control of infection in patients
22 diagnosed during the primary infection (12, 16). There are some cases in which PTC status
23 was achieved after the interruption of an antiretroviral treatment initiated (in the context of
24 detectable viremia) during chronic infection (128), but these cases are much more

1 infrequent. Therefore, early treatment appears to enhance the conditions that allow control
2 of infection in some patients. Early treatment initiation limits the establishment of viral
3 reservoirs (129) and the diversity of the virus (130), which may be easier to control later by
4 undamaged immune responses (131-133). However, it is still unknown why only a fraction of
5 patients treated early are able to control infection after treatment cessation. The precocity
6 and duration of the treatment and the decrease of viral reservoirs to extremely low levels at
7 the time of treatment interruption distinguished PTCs from non-controllers in two studies
8 (124, 125). Similarly, the 14 PTCs described in the VISCONTI study, who have been
9 controlling infection off therapy for a median of 7.5 years, initiated their treatment
10 approximately 35 days after their presumed infection and were treated for a median of
11 three years before interrupting treatment (127). The PTCs from the VISCONTI study also
12 have very low viral reservoirs, similar to those of HICs, in which there is a small contribution
13 of long-lived cells. Interestingly, these viral reservoirs have appeared to shrink in some PTCs
14 in the absence of therapy (127). Nevertheless, further studies will be necessary to
15 understand the factors allowing the control of infection in these appealing patients.

16

17 **Conclusions**

18 From this review, it appears clear that HICs are not a homogeneous population; they can be
19 distinguished by specific transcriptomic, genomic and immunological profiles. They are
20 heterogeneous in terms of the viral loads determined by ultrasensitive single-copy PCR
21 assays, HLA class I alleles, the frequency and quality of CD8+ and CD4+ T cell responses and
22 levels of immune activation. It is yet not possible to associate a multiparametric profile that
23 distinguishes different subpopulations of HICs. It is possible that the key parameters that
24 determine eventual viral control may only be detectable early in infection and have thus

1 been partially missed in recent studies. Indeed, as mentioned, viral control is achieved early
2 in infection in most patients. It is urgent to perform longitudinal studies on cohorts of
3 acutely infected patients to identify viral and host factors that determine viral control.
4 However, the data suggest that some HICs maintain high CD4+ T cell counts and
5 transcriptional profiles and levels of immune activation similar to those in non-infected
6 individuals. A specific effort should be dedicated to perform a more in-depth
7 characterization of these individuals who may provide the most significant keys to
8 understanding the mechanisms of viral control and to evaluate whether it is possible to
9 induce viral control using preventive or therapeutic strategies. Many questions remain about
10 the relative contribution of viral factors, innate host mechanisms of defense and specific
11 anti-HIV T cell responses in the control of viral replication and the containment of viral
12 reservoirs. One critical question is what viral reservoirs in the blood and tissue
13 compartments fuel low-level viral replication in most HICs. Recently, it has been reported
14 that the CD4+ T follicular helper (TFH) cell subset is expanded in lymph node germinal
15 centers during HIV and SIV infection (134, 135), (136, 137). Replication-competent HIV was
16 isolated from TFH cells not only from patients with high viremia but also from patients with
17 low viremia (< 1000 copies/ml) after *in vitro* activation (137), suggesting that they may also
18 be a source of viral replication in HICs. It would be informative to explore the frequency of
19 HIV-infected TFH cells in the lymphatic tissues of HICs and their permissiveness to HIV-1
20 replication.

21 In contrast with HIC, who maintain a functional immune system because of a tight control of
22 HIV replication, a natural control of pathogenesis in SIV-infected African monkeys is achieved
23 in spite of high levels of virus replication. Sooty Mangabeys and African Green Monkeys, the
24 best studied models of apathogenic infection, are protected against disease mainly because

1 of their capacity to rapidly down-regulate innate immune responses, including type I
2 interferon production and IFN-related genes activation, after the burst that follows acute
3 infection. The absence of chronic immune activation preserves lymphoid tissue integrity and
4 immune functions, including the memory T cell pool. A thorough discussion on the
5 mechanisms of control of viral pathogenesis in SIV-infected African monkeys can be found in
6 (138).

7 As for the mechanisms of viral control in HIC, the data reviewed here support a complex
8 model that involves different components of the immune system as well as those of the
9 infecting virus (Figure 1). The different mechanisms may act together, or a specific
10 mechanism may prevail in different individuals. It could be speculated that at the moment of
11 viral transmission, an attenuated virus may result in slower replication kinetics and viral
12 spread in some individuals. This would allow a better preservation of the immune system
13 and the development of an adequate innate and specific cell response. In other individuals
14 infected with fit viruses, either a lower permissiveness of the target cells or a rapid down-
15 regulation of immune activation may hinder viral replication and the establishment of
16 reservoirs. Efficient innate antiviral responses may also contribute to early viral containment.
17 It is possible that in the case of PTCs, treatment during acute infection may reach the same
18 goal by reducing the early dynamics of viral replication and limiting viral reservoir formation.
19 In HICs, the genetic HLA background and/or unknown factors regulating T cell responses,
20 including the selection of high CD8+ T cell receptor avidity, may take over during a second
21 phase to maintain control of viral replication following the earliest responses in individuals
22 who experience viral blips. Studies on HICs have highlighted the role of effective CTL
23 responses, specifically those directed to HIV-1-Gag epitopes, in viral control. As discussed
24 above, this CTL functional capacity can predict effective antiviral function and is measured by

1 *in vitro* suppressive assays. The protective role for CTLs *in vivo* has been recently supported
2 by vaccine studies in a rhesus macaque model. The protective capacity of MHC-driven CTL
3 responses has been demonstrated by the vaccination of Indian rhesus macaques that
4 express the protective MHC Mamu-B*08 allele with Mamu-B*08-restricted CD8+ T-cell
5 epitopes (139). The vaccinated animals controlled the replication of a pathogenic SIV clone
6 after challenge. Perhaps even more promising, because the protection was not directly
7 associated with the MHC background, the induction of SIV-specific effector memory T cells
8 by vaccination with rhesus cytomegalovirus vectors efficiently controlled viral replication in
9 macaques following mucosal challenge (140).

10 Although much work remains to be done to translate these vaccine approaches from the
11 monkey model to humans, these data, together with the promising results arisen from the
12 PTC studies, nurture hope that induction of HIV control may be possible, well beyond the
13 small population of HIV controllers, in a not so far future.

14

15 **Acknowledgments**

16 Our researches on HIV controllers are funded by the ANRS, the French National Agency
17 for Research on AIDS and Viral Hepatitis.

18

19 **References**

- 20 1. Deacon NJ, et al. Genomic structure of an attenuated quasi species of HIV-1 from a blood
21 transfusion donor and recipients. *Science*.1995;270:988-991.
- 22 2. Hirsch VM, et al. Vpx is required for dissemination and pathogenesis of SIV(SM) PBj: evidence
23 of macrophage-dependent viral amplification. *Nature medicine*.1998;4:1401-1408.
- 24 3. Verhofstede C, Nijhuis M, Vandekerckhove L. Correlation of coreceptor usage and disease
25 progression. *Curr Opin HIV AIDS*.2012;7:432-439.

- 1 4. Brass AL, et al. Identification of host proteins required for HIV infection through a functional
2 genomic screen. *Science*.2008;319:921-926.
- 3 5. Konig R, et al. Global analysis of host-pathogen interactions that regulate early-stage HIV-1
4 replication. *Cell*.2008;135:49-60.
- 5 6. Yan N, Chen ZJ. Intrinsic antiviral immunity. *Nature immunology*.2012;13:214-222.
- 6 7. Lambotte O, et al. HIV controllers: a homogeneous group of HIV-1-infected patients with
7 spontaneous control of viral replication. *Clin Infect Dis*.2005;41:1053-1056.
- 8 8. Migueles SA, et al. HLA B*5701 is highly associated with restriction of virus replication in a
9 subgroup of HIV-infected long term nonprogressors. *Proc Natl Acad Sci U S A*.2000;97:2709-2714.
- 10 9. Okulicz JF, et al. Clinical outcomes of elite controllers, viremic controllers, and long-term
11 nonprogressors in the US Department of Defense HIV natural history study. *J Infect*
12 *Dis*.2009;200:1714-1723.
- 13 10. Pereyra F, et al. Genetic and immunologic heterogeneity among persons who control HIV
14 infection in the absence of therapy. *J Infect Dis*.2008;197:563-571.
- 15 11. Saez-Cirion A, Pancino G, Sinet M, Venet A, Lambotte O. HIV controllers: how do they tame
16 the virus? *Trends Immunol*.2007;28:532-540.
- 17 12. Madec Y, Boufassa F, Porter K, Meyer L. Spontaneous control of viral load and CD4 cell count
18 progression among HIV-1 seroconverters. *AIDS*.2005;19:2001-2007.
- 19 13. Boufassa F, et al. CD4 dynamics over a 15 year-period among HIV controllers enrolled in the
20 ANRS French observatory. *PLoS ONE*.2011;6:e18726.
- 21 14. Grabar S, Selinger-Leneman H, Abgrall S, Pialoux G, Weiss L, Costagliola D. Prevalence and
22 comparative characteristics of long-term nonprogressors and HIV controller patients in the French
23 Hospital Database on HIV. *AIDS*.2009;23:1163-1169.
- 24 15. Saez-Cirion A, Pancino G, Lambotte O. Definition, Natural History and Heterogeneity of HIV
25 controllers. In: Pancino G, Silvestri G, Fowke KR, eds. *Models of Protection Against HIV/SIV Avoiding*
26 *AIDS in humans and monkeys: Academic Press, 2012:233-252.*
- 27 16. Goujard C, et al. Spontaneous control of viral replication during primary HIV infection: when
28 is "HIV controller" status established? *Clin Infect Dis*.2009;49:982-986.
- 29 17. Kim AY, et al. Spontaneous control of HCV is associated with expression of HLA-B 57 and
30 preservation of targeted epitopes. *Gastroenterology*.2011;140:686-696 e681.
- 31 18. Ruiz-Mateos E, et al. Hepatitis C virus replication in Caucasian HIV controllers. *J Viral*
32 *Hepat*.2011;18:e350-357.
- 33 19. Hatano H, et al. Evidence for persistent low-level viremia in individuals who control human
34 immunodeficiency virus in the absence of antiretroviral therapy. *J Virol*.2009;83:329-335.
- 35 20. Mens H, et al. HIV-1 continues to replicate and evolve in patients with natural control of HIV
36 infection. *J Virol*.2010;84:12971-12981.
- 37 21. Pereyra F, et al. Persistent low-level viremia in HIV-1 elite controllers and relationship to
38 immunologic parameters. *J Infect Dis*.2009;200:984-990.
- 39 22. Dahl V, et al. Single-copy assay quantification of HIV-1 RNA in paired cerebrospinal fluid and
40 plasma samples from elite controllers. *AIDS*.2012.
- 41 23. Probasco JC, et al. Cerebrospinal fluid in HIV-1 systemic viral controllers: absence of HIV-1
42 RNA and intrathecal inflammation. *AIDS*.2010;24:1001-1005.
- 43 24. Hunt PW, et al. Relationship between T cell activation and CD4+ T cell count in HIV-
44 seropositive individuals with undetectable plasma HIV RNA levels in the absence of therapy. *J Infect*
45 *Dis*.2008;197:126-133.
- 46 25. Vigneault F, et al. Transcriptional profiling of CD4 T cells identifies distinct subgroups of HIV-1
47 elite controllers. *J Virol*.2011;85:3015-3019.
- 48 26. Okulicz JF, et al. CD4 T cell count reconstitution in HIV controllers after highly active
49 antiretroviral therapy. *Clin Infect Dis*.2010;50:1187-1191.
- 50 27. Sedaghat AR, Rastegar DA, O'Connell KA, Dinoso JB, Wilke CO, Blankson JN. T cell dynamics
51 and the response to HAART in a cohort of HIV-1-infected elite suppressors. *Clin Infect*
52 *Dis*.2009;49:1763-1766.

- 1 28. Brenchley JM, et al. Microbial translocation is a cause of systemic immune activation in
2 chronic HIV infection. *Nat Med.*2006;12:1365-1371.
- 3 29. Potter SJ, et al. Preserved central memory and activated effector memory CD4+ T-cell subsets
4 in human immunodeficiency virus controllers: an ANRS EP36 study. *J Virol.*2007;81:13904-13915.
- 5 30. Yang Y, et al. CD4 T-cell regeneration in HIV-1 elite controllers. *AIDS.*2012;26:701-706.
- 6 31. Sauce D, et al. HIV disease progression despite suppression of viral replication is associated
7 with exhaustion of lymphopoiesis. *Blood.*2011;117:5142-5151.
- 8 32. Bailey JR, Williams TM, Siliciano RF, Blankson JN. Maintenance of viral suppression in HIV-1-
9 infected HLA-B*57+ elite suppressors despite CTL escape mutations. *J Exp Med.*2006;203:1357-1369.
- 10 33. Dinoso JB, Kim SY, Siliciano RF, Blankson JN. A comparison of viral loads between HIV-1-
11 infected elite suppressors and individuals who receive suppressive highly active antiretroviral
12 therapy. *Clin Infect Dis.*2008;47:102-104.
- 13 34. Blankson JN, et al. Isolation and Characterization of Replication-Competent HIV-1 from a
14 Subset of Elite Suppressors. *J Virol.*2007;81:2508-2518.
- 15 35. Lewin SR, Rouzioux C. HIV cure and eradication: how will we get from the laboratory to
16 effective clinical trials? *Aids.*2011;25:885-897.
- 17 36. O'Connell KA, Brennan TP, Bailey JR, Ray SC, Siliciano RF, Blankson JN. Control of HIV-1 in elite
18 suppressors despite ongoing replication and evolution in plasma virus. *J Virol.*2010;84:7018-7028.
- 19 37. Salgado M, et al. Evolution of the HIV-1 nef gene in HLA-B*57 positive elite suppressors.
20 *Retrovirology.*2010;7:94.
- 21 38. Schneidewind A, et al. Escape from the dominant HLA-B27-restricted cytotoxic T-lymphocyte
22 response in Gag is associated with a dramatic reduction in human immunodeficiency virus type 1
23 replication. *J Virol.*2007;81:12382-12393.
- 24 39. Brockman MA, et al. Escape and compensation from early HLA-B57-mediated cytotoxic T-
25 lymphocyte pressure on human immunodeficiency virus type 1 Gag alter capsid interactions with
26 cyclophilin A. *J Virol.*2007;81:12608-12618.
- 27 40. Miura T, et al. HLA-B57/B*5801 human immunodeficiency virus type 1 elite controllers select
28 for rare gag variants associated with reduced viral replication capacity and strong cytotoxic T-
29 lymphocyte [corrected] recognition. *J Virol.*2009;83:2743-2755.
- 30 41. Martinez-Picado J, et al. Fitness cost of escape mutations in p24 Gag in association with
31 control of human immunodeficiency virus type 1. *J Virol.*2006;80:3617-3623.
- 32 42. Miura T, et al. HLA-associated alterations in replication capacity of chimeric NL4-3 viruses
33 carrying gag-protease from elite controllers of human immunodeficiency virus type 1. *J*
34 *Virol.*2009;83:140-149.
- 35 43. Lassen KG, et al. Elite suppressor-derived HIV-1 envelope glycoproteins exhibit reduced entry
36 efficiency and kinetics. *PLoS Pathog.*2009;5:e1000377.
- 37 44. Brumme ZL, et al. Reduced replication capacity of NL4-3 recombinant viruses encoding
38 reverse transcriptase-integrase sequences from HIV-1 elite controllers. *J Acquir Immune Defic*
39 *Syndr.*2011;56:100-108.
- 40 45. Mwimanzi P, et al. Attenuation of multiple Nef functions in HIV-1 elite controllers.
41 *Retrovirology.*2012;10:1.
- 42 46. Miura T, et al. Genetic characterization of human immunodeficiency virus type 1 in elite
43 controllers: lack of gross genetic defects or common amino acid changes. *J Virol.*2008;82:8422-8430.
- 44 47. Miura T, et al. Impaired replication capacity of acute/early viruses in persons who become
45 HIV controllers. *J Virol.*2010;84:7581-7591.
- 46 48. Lamine A, et al. Replication-competent HIV strains infect HIV controllers despite
47 undetectable viremia (ANRS EP36 study). *Aids.*2007;21:1043-1045.
- 48 49. Saez-Cirion A, et al. Heterogeneity in HIV suppression by CD8 T cells from HIV controllers:
49 association with Gag-specific CD8 T cell responses. *J Immunol.*2009;182:7828-7837.
- 50 50. Julg B, et al. Infrequent recovery of HIV from but robust exogenous infection of activated
51 CD4(+) T cells in HIV elite controllers. *Clin Infect Dis.*2010;51:233-238.

- 1 51. Buckheit RW, 3rd, et al. Host factors dictate control of viral replication in two HIV-1
2 controller/chronic progressor transmission pairs. *Nature communications*.2012;3:716.
- 3 52. Navis M, et al. Viral replication capacity as a correlate of HLA B57/B5801-associated
4 nonprogressive HIV-1 infection. *J Immunol*.2007;179:3133-3143.
- 5 53. Bailey JR, et al. Transmission of human immunodeficiency virus type 1 from a patient who
6 developed AIDS to an elite suppressor. *J Virol*.2008;82:7395-7410.
- 7 54. Fellay J, et al. A whole-genome association study of major determinants for host control of
8 HIV-1. *Science*.2007;317:944-947.
- 9 55. Limou S, et al. Genomewide association study of an AIDS-nonprogression cohort emphasizes
10 the role played by HLA genes (ANRS Genomewide Association Study 02). *J Infect Dis*.2009;199:419-
11 426.
- 12 56. Kaslow RA, et al. Influence of combinations of human major histocompatibility complex
13 genes on the course of HIV-1 infection. *Nat Med*.1996;2:405-411.
- 14 57. Pelak K, et al. Host determinants of HIV-1 control in African Americans. *J Infect*
15 *Dis*.2010;201:1141-1149.
- 16 58. McLaren PJ, et al. Fine-mapping classical HLA variation associated with durable host control
17 of HIV-1 infection in African Americans. *Human molecular genetics*.2012;21:4334-4347.
- 18 59. Pereyra F, et al. The major genetic determinants of HIV-1 control affect HLA class I peptide
19 presentation. *Science*.2010;330:1551-1557.
- 20 60. Thomas R, et al. HLA-C cell surface expression and control of HIV/AIDS correlate with a
21 variant upstream of HLA-C. *Nature genetics*.2009;41:1290-1294.
- 22 61. Kulkarni S, et al. Differential microRNA regulation of HLA-C expression and its association
23 with HIV control. *Nature*.2011;472:495-498.
- 24 62. Goldstein DB. Common genetic variation and human traits. *The New England journal of*
25 *medicine*.2009;360:1696-1698.
- 26 63. Martin MP, et al. Epistatic interaction between KIR3DS1 and HLA-B delays the progression to
27 AIDS. *Nature genetics*.2002;31:429-434.
- 28 64. Kanya P, et al. Receptor-ligand requirements for increased NK cell polyfunctional potential in
29 slow progressors infected with HIV-1 coexpressing KIR3DL1*h/*y and HLA-B*57. *J*
30 *Virol*.2011;85:5949-5960.
- 31 65. Martin MP, et al. Innate partnership of HLA-B and KIR3DL1 subtypes against HIV-1. *Nature*
32 *genetics*.2007;39:733-740.
- 33 66. Tomescu C, et al. Impact of protective killer inhibitory receptor/human leukocyte antigen
34 genotypes on natural killer cell and T-cell function in HIV-1-infected controllers. *Aids*.2012;26:1869-
35 1878.
- 36 67. O'Connell KA, Han Y, Williams TM, Siliciano RF, Blankson JN. Role of natural killer cells in a
37 cohort of elite suppressors: low frequency of the protective KIR3DS1 allele and limited inhibition of
38 human immunodeficiency virus type 1 replication in vitro. *J Virol*.2009;83:5028-5034.
- 39 68. Sáez-Cirión A, et al. HIV controllers exhibit potent CD8 T cell capacity to suppress HIV
40 infection ex vivo and peculiar CTL activation phenotype. *Proc Natl Acad Sci U S A*.2007;104:6776-
41 6781.
- 42 69. Day CL, et al. PD-1 expression on HIV-specific T cells is associated with T-cell exhaustion and
43 disease progression. *Nature*.2006;443:350-354.
- 44 70. Petrovas C, et al. Differential association of programmed death-1 and CD57 with ex vivo
45 survival of CD8+ T cells in HIV infection. *J Immunol*.2009;183:1120-1132.
- 46 71. Giorgi JV, et al. Shorter survival in advanced human immunodeficiency virus type 1 infection
47 is more closely associated with T lymphocyte activation than with plasma virus burden or virus
48 chemokine coreceptor usage. *The Journal of infectious diseases*.1999;179:859-870.
- 49 72. Horton H, et al. Preservation of T cell proliferation restricted by protective HLA alleles is
50 critical for immune control of HIV-1 infection. *J Immunol*.2006;177:7406-7415.
- 51 73. Betts MR, et al. HIV nonprogressors preferentially maintain highly functional HIV-specific
52 CD8+ T cells. *Blood*.2006;107:4781-4789.

- 1 74. Streeck H, et al. Antigen load and viral sequence diversification determine the functional
2 profile of HIV-1-specific CD8+ T cells. *PLoS medicine*.2008;5:e100.
- 3 75. Migueles SA, et al. HIV-specific CD8+ T cell proliferation is coupled to perforin expression and
4 is maintained in nonprogressors. *Nat Immunol*.2002;3:1061-1068.
- 5 76. Migueles SA, et al. Lytic granule loading of CD8+ T cells is required for HIV-infected cell
6 elimination associated with immune control. *Immunity*.2008;29:1009-1021.
- 7 77. Hersperger AR, et al. Perforin expression directly ex vivo by HIV-specific CD8 T-cells is a
8 correlate of HIV elite control. *PLoS Pathog*.2010;6:e1000917.
- 9 78. Addo MM, et al. Fully differentiated HIV-1 specific CD8+ T effector cells are more frequently
10 detectable in controlled than in progressive HIV-1 infection. *PLoS one*.2007;2:e321.
- 11 79. Chen H, et al. Differential neutralization of human immunodeficiency virus (HIV) replication
12 in autologous CD4 T cells by HIV-specific cytotoxic T lymphocytes. *J Virol*.2009;83:3138-3149.
- 13 80. Kiepiela P, et al. CD8+ T-cell responses to different HIV proteins have discordant associations
14 with viral load. *Nat Med*.2007;13:46-53.
- 15 81. Pontesilli O, et al. Longitudinal analysis of human immunodeficiency virus type 1-specific
16 cytotoxic T lymphocyte responses: a predominant gag-specific response is associated with
17 nonprogressive infection. *J Infect Dis*.1998;178:1008-1018.
- 18 82. Edwards BH, Bansal A, Sabbaj S, Bakari J, Mulligan MJ, Goepfert PA. Magnitude of functional
19 CD8+ T-cell responses to the gag protein of human immunodeficiency virus type 1 correlates
20 inversely with viral load in plasma. *J Virol*.2002;76:2298-2305.
- 21 83. Descours B, et al. Immune responses driven by protective human leukocyte antigen alleles
22 from long-term nonprogressors are associated with low HIV reservoir in central memory CD4 T cells.
23 *Clin Infect Dis*.2012;54:1495-1503.
- 24 84. Ferre AL, et al. Mucosal immune responses to HIV-1 in elite controllers: a potential correlate
25 of immune control. *Blood*.2009;113:3978-3989.
- 26 85. Ferre AL, et al. Immunodominant HIV-specific CD8+ T-cell responses are common to blood
27 and gastrointestinal mucosa, and Gag-specific responses dominate in rectal mucosa of HIV
28 controllers. *J Virol*.2010;84:10354-10365.
- 29 86. Miura T, et al. HLA-associated viral mutations are common in human immunodeficiency virus
30 type 1 elite controllers. *J Virol*.2009;83:3407-3412.
- 31 87. Wang YE, et al. Protective HLA class I alleles that restrict acute-phase CD8+ T-cell responses
32 are associated with viral escape mutations located in highly conserved regions of human
33 immunodeficiency virus type 1. *J Virol*.2009;83:1845-1855.
- 34 88. Berger CT, et al. High-functional-avidity cytotoxic T lymphocyte responses to HLA-B-restricted
35 Gag-derived epitopes associated with relative HIV control. *J Virol*.2011;85:9334-9345.
- 36 89. Mothe B, et al. CTL responses of high functional avidity and broad variant cross-reactivity are
37 associated with HIV control. *PLoS One*.2012;7:e29717.
- 38 90. Almeida JR, et al. Antigen sensitivity is a major determinant of CD8+ T-cell polyfunctionality
39 and HIV-suppressive activity. *Blood*.2009;113:6351-6360.
- 40 91. Chen H, et al. TCR clonotypes modulate the protective effect of HLA class I molecules in HIV-1
41 infection. *Nat Immunol*.2012;13:691-700.
- 42 92. Ladell K, et al. A molecular basis for the control of pre-immune escape variants by HIV-
43 specific CD8+ T-cells. *Immunity*.2013;In press.
- 44 93. Huang J, et al. Leukocyte immunoglobulin-like receptors maintain unique antigen-presenting
45 properties of circulating myeloid dendritic cells in HIV-1-infected elite controllers. *J*
46 *Virol*.2010;84:9463-9471.
- 47 94. Lecuroux C, et al. CD8 T-cells from most HIV-infected patients lack HIV-suppressive capacity
48 during acute and early infection. *PLoS One*.2013;In press.
- 49 95. Emu B, et al. HLA class I-restricted T-cell responses may contribute to the control of human
50 immunodeficiency virus infection, but such responses are not always necessary for long-term virus
51 control. *J Virol*.2008;82:5398-5407.

- 1 96. Buckheit RW, 3rd, Salgado M, Silciano RF, Blankson JN. Inhibitory Potential of Subpopulations
2 of CD8+ T Cells in HIV-1 Infected Elite Suppressors. *J Virol.*2012.
- 3 97. Ndhlovu ZM, et al. Elite controllers with low to absent effector CD8+ T cell responses
4 maintain highly functional, broadly directed central memory responses. *J Virol.*2012;86:6959-6969.
- 5 98. Shan L, et al. Stimulation of HIV-1-specific cytolytic T lymphocytes facilitates elimination of
6 latent viral reservoir after virus reactivation. *Immunity.*2012;36:491-501.
- 7 99. Schulze Zur Wiesch J, et al. Comprehensive analysis of frequency and phenotype of T
8 regulatory cells in HIV infection: CD39 expression of FoxP3+ T regulatory cells correlates with
9 progressive disease. *J Virol.*2011;85:1287-1297.
- 10 100. Angin M, et al. Preserved function of regulatory T cells in chronic HIV-1 infection despite
11 decreased numbers in blood and tissue. *J Infect Dis.*2012;205:1495-1500.
- 12 101. Shaw JM, et al. Increased frequency of regulatory T cells accompanies increased immune
13 activation in rectal mucosae of HIV-positive noncontrollers. *J Virol.*2011;85:11422-11434.
- 14 102. Hunt PW, et al. A low T regulatory cell response may contribute to both viral control and
15 generalized immune activation in HIV controllers. *PLoS ONE.*2011;6:e15924.
- 16 103. Lambotte O, et al. Heterogeneous neutralizing antibody and antibody-dependent cell
17 cytotoxicity responses in HIV-1 elite controllers. *Aids.*2009;23:897-906.
- 18 104. Deepe RN, Kistner-Griffin E, Martin JN, Deeks SG, Pandey JP. Epistatic interactions between
19 Fc (GM) and FcgammaR genes and the host control of human immunodeficiency virus replication.
20 *Hum Immunol.*2012;73:263-266.
- 21 105. Liu Q, et al. Matrix metalloprotease inhibitors restore impaired NK cell-mediated antibody-
22 dependent cellular cytotoxicity in human immunodeficiency virus type 1 infection. *Journal of*
23 *virology.*2009;83:8705-8712.
- 24 106. Mendoza D, et al. Comprehensive analysis of unique cases with extraordinary control over
25 HIV replication. *Blood.*2012;119:4645-4655.
- 26 107. Rotger M, et al. Genome-wide mRNA expression correlates of viral control in CD4+ T-cells
27 from HIV-1-infected individuals. *PLoS Pathog.*2010;6:e1000781.
- 28 108. Owen RE, et al. HIV+ elite controllers have low HIV-specific T-cell activation yet maintain
29 strong, polyfunctional T-cell responses. *Aids.*2010;24:1095-1105.
- 30 109. Vingert B, et al. HIV controllers maintain a population of highly efficient Th1 effector cells in
31 contrast to patients treated in the long term. *J Virol.*2012;86:10661-10674.
- 32 110. Kaufmann DE, et al. Upregulation of CTLA-4 by HIV-specific CD4+ T cells correlates with
33 disease progression and defines a reversible immune dysfunction. *Nat Immunol.*2007;8:1246-1254.
- 34 111. Emu B, et al. Phenotypic, functional, and kinetic parameters associated with apparent T-cell
35 control of human immunodeficiency virus replication in individuals with and without antiretroviral
36 treatment. *J Virol.*2005;79:14169-14178.
- 37 112. van Grevenynghe J, et al. Transcription factor FOXO3a controls the persistence of memory
38 CD4(+) T cells during HIV infection. *Nat Med.*2008;14:266-274.
- 39 113. Vingert B, et al. HIV controller CD4+ T cells respond to minimal amounts of Gag antigen due
40 to high TCR avidity. *PLoS Pathog.*2010;6:e1000780.
- 41 114. Ferre AL, et al. HIV controllers with HLA-DRB1*13 and HLA-DQB1*06 alleles have strong,
42 polyfunctional mucosal CD4+ T-cell responses. *J Virol.*2010;84:11020-11029.
- 43 115. Chomont N, et al. HIV reservoir size and persistence are driven by T cell survival and
44 homeostatic proliferation. *Nat Med.*2009;15:893-900.
- 45 116. Chen H, et al. CD4+ T cells from elite controllers resist HIV-1 infection by selective
46 upregulation of p21. *J Clin Invest.*2011;121:1549-1560.
- 47 117. Saez-Cirion A, et al. Restriction of HIV-1 replication in macrophages and CD4+ T cells from HIV
48 controllers. *Blood.*2011;118:955-964.
- 49 118. Buzon MJ, et al. Inhibition of HIV-1 integration in ex vivo-infected CD4 T cells from elite
50 controllers. *J Virol.*2011;85:9646-9650.
- 51 119. Zhang J, Scadden DT, Crumpacker CS. Primitive hematopoietic cells resist HIV-1 infection via
52 p21. *J Clin Invest.*2007;117:473-481.

- 1 120. Bergamaschi A, David A, Le Rouzic E, Nisole S, Barre-Sinoussi F, Pancino G. The CDK inhibitor
2 p21Cip1/WAF1 is induced by FcγR activation and restricts the replication of human
3 immunodeficiency virus type 1 and related primate lentiviruses in human macrophages. *J*
4 *Virol.*2009;83:12253-12265.
- 5 121. O'Connell KA, Rabi SA, Siliciano RF, Blankson JN. CD4+ T cells from elite suppressors are more
6 susceptible to HIV-1 but produce fewer virions than cells from chronic progressors. *Proc Natl Acad Sci*
7 *U S A.*2011;108:E689-698.
- 8 122. Palacios JA, et al. Long-term nonprogressor and elite controller patients who control viremia
9 have a higher percentage of methylation in their HIV-1 proviral promoters than aviremic patients
10 receiving highly active antiretroviral therapy. *J Virol.*2012;86:13081-13084.
- 11 123. Deeks SG, et al. Towards an HIV cure: a global scientific strategy. *Nat Rev*
12 *Immunol.*2012;12:607-614.
- 13 124. Goujard C, et al. HIV-1 control after transient antiretroviral treatment initiated in primary
14 infection: role of patient characteristics and effect of therapy. *Antivir Ther.*2012;17:1001-1009.
- 15 125. Hocqueloux L, et al. Long-term immunovirologic control following antiretroviral therapy
16 interruption in patients treated at the time of primary HIV-1 infection. *Aids.*2010;24:1598-1601.
- 17 126. Lodi S, et al. Immunovirologic control 24 months after interruption of antiretroviral therapy
18 initiated close to HIV seroconversion. *Arch Intern Med.*2012;172:1252-1255.
- 19 127. Saez-Cirion A, et al. Post-treatment HIV-1 controllers with a long-term virological remission
20 after the interruption of early initiated antiretroviral therapy. ANRS VISCONTI study. *PLoS*
21 *Pathog.*2013.
- 22 128. Van Gulck E, et al. Immune and viral correlates of "secondary viral control" after treatment
23 interruption in chronically HIV-1 infected patients. *PLoS ONE.*2012;7:e37792.
- 24 129. Ngo-Giang-Huong N, et al. Proviral HIV-1 DNA in subjects followed since primary HIV-1
25 infection who suppress plasma viral load after one year of highly active antiretroviral therapy.
26 *AIDS.*2001;15:665-673.
- 27 130. Delwart E, et al. Homogeneous quasispecies in 16 out of 17 individuals during very early HIV-
28 1 primary infection. *AIDS.*2002;16:189-195.
- 29 131. Alter G, et al. Sequential deregulation of NK cell subset distribution and function starting in
30 acute HIV-1 infection. *Blood.*2005;106:3366-3369.
- 31 132. Moir S, et al. B cells in early and chronic HIV infection: evidence for preservation of immune
32 function associated with early initiation of antiretroviral therapy. *Blood.*2010;116:5571-5579.
- 33 133. Oxenius A, et al. Early highly active antiretroviral therapy for acute HIV-1 infection preserves
34 immune function of CD8+ and CD4+ T lymphocytes. *Proc Natl Acad Sci U S A.*2000;97:3382-3387.
- 35 134. Petrovas C, et al. CD4 T follicular helper cell dynamics during SIV infection. *J Clin*
36 *Invest.*2012;122:3281-3294.
- 37 135. Lindqvist M, et al. Expansion of HIV-specific T follicular helper cells in chronic HIV infection. *J*
38 *Clin Invest.*2012;122:3271-3280.
- 39 136. Hong JJ, Amancha PK, Rogers K, Ansari AA, Villinger F. Spatial alterations between CD4(+) T
40 follicular helper, B, and CD8(+) T cells during simian immunodeficiency virus infection: T/B cell
41 homeostasis, activation, and potential mechanism for viral escape. *J Immunol.*2012;188:3247-3256.
- 42 137. Perreau M, et al. Follicular helper T cells serve as the major CD4 T cell compartment for HIV-1
43 infection, replication, and production. *J Exp Med.*2012;210:143-156.
- 44 138. Pancino G, Silvestri G, Fowke K. Models of Protection Against HIV/SIV. Avoiding AIDS in
45 humans and monkeys. Academic Press, 2012.
- 46 139. Mudd PA, et al. Vaccine-induced CD8+ T cells control AIDS virus replication.
47 *Nature.*2012;491:129-133.
- 48 140. Hansen SG, et al. Profound early control of highly pathogenic SIV by an effector memory T-
49 cell vaccine. *Nature.*2011;473:523-527.
- 50
51
52

1 **Figure legends**

2

3 **Figure 1A**

4 In the acute phase of infection of the majority of individuals, HIV-1 exponentially replicates
5 in target cells, including CD4+ T cells and monocytes/macrophages, reaching levels of several
6 million copies of plasma viral RNA/ml. Viral load then declines with the development of host
7 immune response and reaches a steady state that is a predictor of long-term disease
8 progression. A massive destruction of CD4+ T lymphocytes occurs in the acute/early phase,
9 especially within the lymphoid tissues of the gut. In the meantime, permanent viral
10 reservoirs are established in the tissues. A dramatic cytokine cascade is activated in the
11 acute phase, along with an abnormal activation of all the compartments of immune system,
12 including NK cells, B and T lymphocytes. The levels of immune activation strongly correlate
13 with the progression of disease. The impairment of HIV-1-specific CD4 T and CD8 T cell
14 functions occurs early in acute infection and the gradual exhaustion of HIV-specific T cell
15 response leads then to the loss of the control of viral replication at later times.

16 **Figure 1B**

17 In HIC, during acute infection, several factors may contribute, together or separately, to
18 reduce the dynamics of HIV replication: the transmission of an attenuated virus, a lower
19 permissiveness of the target cells (CD4+ T cells, dendritic cells and macrophages), efficient
20 innate antiviral responses and a rapid down regulation of immune activation. This would
21 limit the size of viral reservoirs and allow a better preservation of the immune system. A
22 favorable genetic HLA background and the selection of high avidity CD8+ T cell receptor may
23 then contribute to suppress viral replication and control viral blips in many HICs. Other still
24 unknown factors may act to maintain viral control in other HIC.



