

## Natural resistance to HIV infection: lessons learned from HIV-exposed uninfected individuals

Gianfranco Pancino, Asier Saez-Cirion, Daniel Scott-Algara, Pascale Paul

► **To cite this version:**

Gianfranco Pancino, Asier Saez-Cirion, Daniel Scott-Algara, Pascale Paul. Natural resistance to HIV infection: lessons learned from HIV-exposed uninfected individuals. *Journal of Infectious Diseases*, Oxford University Press (OUP), 2010, 202 Suppl 3, pp.S345–350. <10.1086/655973>. <pasteur-01420585>

**HAL Id: pasteur-01420585**

**<https://hal-pasteur.archives-ouvertes.fr/pasteur-01420585>**

Submitted on 16 May 2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



1                   **Natural resistance to HIV infection: lessons learned from**  
2                   **HIV-exposed uninfected individuals**

3  
4    *Gianfranco Pancino<sup>1</sup>, Asier Saez-Cirion<sup>1</sup>, Daniel Scott-Algara<sup>1</sup> and Pascale Paul<sup>2</sup>*

5    <sup>1</sup>Institut Pasteur, Unité de Régulation des Infections Rétrovirales, Paris, France, <sup>2</sup> Laboratoire  
6    Hématologie Spécialisée, Laboratoire d'Hématologie, Hôpital de la Conception, and UMR-S  
7    608, Faculté de Pharmacie, Université de la Méditerranée, Marseille, France

8  
9  
10  
11  
12   Running head: Lessons learned from EU

13  
14   Word counts

15   Abstract : 100

16   Text: 2291

17

1 **Footnote page**

2

3 Authors do not have conflicts of interest

4

5 The researches have been funded by the French Agency of Research on AIDS and viral  
6 hepatitis (ANRS) and Sidaction, France

7

8 This paper has been presented at the Symposium on Natural Immunity to HIV, Winnipeg,  
9 Manitoba, Canada, November 15-17, 2009

10

11 Corresponding author

12 Gianfranco Pancino, Institut Pasteur, Unité de Régulation des Infections Rétrovirales, 25, rue  
13 du Docteur Roux, Paris, France. email : [gianfranco.pancino@pasteur.fr](mailto:gianfranco.pancino@pasteur.fr)

14

1 **Abstract**

2 We explored potential mechanisms of resistance to HIV-1 infection in different groups of  
3 uninfected individuals exposed by systemic or mucosal routes: intravascular drug-users in  
4 Vietnam, spouses of HIV-infected individuals in Cambodia and Central African Republic.  
5 Our main findings were: reduced susceptibility of PBMC to HIV-1 infection *in vitro*,  
6 associated with low levels of CD4+T cell activation *in vivo* and/or cell restriction of viral  
7 replication; enhanced NK cell activity associated with increased activating/inhibitory NK cell  
8 receptor ratios. These results support a contribution of innate responses to resistance against  
9 HIV-1 infection. Scientific and ethical issues encountered during research on EU are  
10 discussed.

11

12 Key words:

13 HIV, Exposed uninfected, risk of infection, natural resistance, viral restriction, NK cells,  
14 immune activation, Africa, South-East Asia, ethics.

15

## 1 INTRODUCTION

2 A small percentage of individuals who have been repeatedly exposed to HIV-1 through sexual  
3 or systemic routes over long periods of time remain free of any detectable sign of infection.  
4 These naturally resistant individuals have been referred to as “Exposed Seronegative” (ESN)  
5 or “Exposed Uninfected” (EU). Since the overwhelming majority of them are HIV-negative  
6 by both serology and PCR we will use here the EU definition.

7 While some EU individuals may remain free of infection by chance (for example, the  
8 probability of transmission by sexual intercourse is low [1]), there is strong evidence that host  
9 mechanisms are involved in resistance to HIV-1 infection [2]. Both genetic polymorphisms,  
10 including the CCR5 $\Delta$ 32 mutation in Caucasian populations, and HIV-specific T and B cell  
11 responses have been detected in EU subjects [3-6] (reviewed in [7]) raising the question of  
12 whether EU are resistant to HIV-1 infection or whether they are protected by specific  
13 immunity. This question is obviously relevant to the possibility that helpful clues for vaccine  
14 research could be inferred from studies of EU.

15 Acquired immune responses to infectious agents require time to develop, and these may be  
16 not able to block primary HIV infection and the consequent integration of the viral DNA into  
17 host genome. Innate mechanisms of resistance and/or immediate/early immune defenses may  
18 be needed to fight the incoming virus and to help to establish efficient adaptive responses. We  
19 explored these potential mechanisms in EU from South-East Asia and Central Africa, notably  
20 through analyzing target cell susceptibility to HIV-1 infection and NK cell function. We will  
21 briefly summarize our main findings and draw conclusions from the whole body of our  
22 studies.

23

## 1 **Study populations**

2 We studied three groups of EU exposed to HIV-1 by either systemic or mucosal routes:  
3 intravascular drug users from Hô Chi Minh City (HCMC), Vietnam, (N: 45; 38 M, 7 F; drug  
4 use time: 13-31 yr, risk factor: shared needles); heterosexual partners of HIV+ individuals  
5 from Phnom Penh, Cambodia (N: 48; 7 M, 41 F; risk factor: unprotected sexual intercourse >  
6 1.5 yr beyond seroconversion), and Bangui, Central African Republic (CAR) (N: 45; 20 M,  
7 25 F; risk factor: unprotected sexual intercourse, communal life: med 8 yr). The populations  
8 studied have been described in detail elsewhere [8-10]. Informed consent was obtained by all  
9 the participants. The exposure to HIV-1 and the risk of infection differed for the three groups.  
10 Vietnamese EU were recruited from a group of IDU, amongst whom 86% were HIV-  
11 seropositive. High seroprevalences of HBV (82%), HCV (100%) and HTLV (80%) in the  
12 EU IDU supported a high risk of infection by HIV-1. The cumulative risk of being HIV-  
13 infected by their infected spouse for EU from Cambodia and CAR was estimated by the  
14 equation  $1-(1-p)^n$ , where  $p$  designates the probability of infection per unprotected  
15 heterosexual intercourse with an infected individual (0.0008 for men, and 0.0012 for women;  
16 adapted from [11] to take into account the 50% increased risk in women compared to men),  
17 and  $n$  the number of reported unprotected sexual intercourses during the period of communal  
18 life. In CAR (2 to 3 intercourses per week), the risk of infection per year was 9.1% for men  
19 and 13.4% for women, and the cumulative risk over 8 years of communal life was 53.5% for  
20 men and 68.3% for women. In Cambodia (1-2 intercourses per week), the cumulative risk of  
21 infection was 6.1% for men and 8.9% for women during one and a half years, and 22.1% for  
22 men and 31.2% for women over 3 years. These data suggest that the probability of remaining  
23 HIV-uninfected owing to some mechanism of resistance or protection, and not by chance, was  
24 highest in the Vietnamese EU. Accordingly, although some mechanisms of resistance

1 appeared to be shared by the three groups of EU, the clearest results were obtained from  
2 studies on Vietnamese EU.

### 3 **Reduced cell susceptibility to HIV-1 infection in vitro**

4 To assess whether the absence of infection could be associated with cell resistance to HIV-1,  
5 we first compared the susceptibility to HIV-1 infection *in vitro* of PBMC from EU to those  
6 from local low risk populations (healthy blood donors). In each of the three groups, PBMC  
7 from EU were more resistant to infection, by one or more viral isolates, than were control  
8 PBMC [8-10]. While infectivity assays were generally performed on PHA-activated PBMC,  
9 in CAR we also evaluated the susceptibility of PBMC to HIV-1 by exposing unstimulated  
10 PBMC to the virus inoculum and activating the cells after infection, to allow the virus to  
11 replicate. This experiment was done to evaluate PBMC susceptibility *ex vivo*, at a baseline  
12 activation state. In this condition, both the R5 and X4 HIV-1 isolates tested replicated to a  
13 significantly lesser extent in PBMC from EU than in control PBMC [10]. Interestingly, the  
14 expression of the activation marker HLA-DR was lower *in vivo* on CD4+ T cells from Central  
15 African EU than of those from controls (p= 0.0001), suggesting decreased activation of  
16 peripheral CD4+ T cells [10]. Vietnamese EU also exhibited a decreased expression of HLA-  
17 DR on their CD4+ T cells compared to controls (absolute number and percentage  
18 <0.001)[12]. Taken together these observation suggest that lowered levels of activated CD4+  
19 T cells may contribute to protection in EU, decreasing the size of the pool of HIV-1 target  
20 cells able to replicate the virus efficiently. It is tempting to speculate that a low level of  
21 CD4+ T cell activation may result in a reduced permissivity to HIV-1 replication *in vivo*, as  
22 suggested by the *ex vivo* PBMC infectivity assays. The causes of the reduced CD4+ T cell  
23 activation in EU have not been elucidated. In another study, a low CD4+ T cell activation in  
24 EU sex workers from Kenya has been associated with an increased frequency of regulatory T  
25 cells [13].

1 Additional factors, including specific mechanisms of restriction of viral replication, may  
2 contribute to the PHA-activated PBMC resistance to HIV-1 infection in EU [14]. We  
3 investigated a potential intrinsic resistance of CD4<sup>+</sup> T cells in Vietnamese EU. In several  
4 cases (7 out of 14 EU analyzed), CD4<sup>+</sup> T cells were resistant to HIV-1 infection [8]. We  
5 followed 5 of these EU for 2-5 years, and found that the CD4 T cell resistance persisted over  
6 time [15]. Different restriction phenotypes were identified in the CD4<sup>+</sup> T cells from these 5  
7 EU [15] (Fig. 1A). In four EU the restriction affected R5 tropic HIV-1 inhibiting viral entry  
8 and was linked either to a low/absent expression of CCR5 co-receptor variants (G106R,  
9 C178R) on cell surface in two individuals or to an enhanced sensitivity to the inhibitory  
10 action of endogenous produced  $\beta$ -chemokines on viral entry in the other two [15].  
11 Remarkably, in one EU the restriction was independent of HIV-1 tropism and suppressed  
12 early reverse transcription products. The restriction also affected other retroviruses such as  
13 SIVmac, SIVagm (Fig 1B) and MoMLV (not shown), and was not overcome at high HIV-1  
14 doses, suggesting that a saturable species-specific inhibitory factor such as TRIM5 $\alpha$  was not  
15 involved [15]. Some CD4<sup>+</sup> T cell clones derived from this EU retained the restriction  
16 phenotype observed in the parental cells (Fig 1B). Infectivity experiments with heterokaryons  
17 obtained by fusion of a resistant T cell clone with a replication competent T cell line  
18 suggested that the restriction was due to the lack of a cellular co-factor needed for viral  
19 replication (Fig. 1C).

## 20 **Enhanced NK cell activity**

21 NK cells may exert antiviral activity during the earliest phases of HIV-1 infection before the  
22 induction of adaptive immune responses, and may influence the efficiency of the adaptive  
23 immune responses. NK cell activity is regulated by the integration of target cell signals  
24 transmitted through inhibitory or activating NK-cell receptors (NKR), including the  
25 polymorphic Killer Immunoglobulin-like Receptors (KIR), the c-type lectin NKG2 receptors,



1 the natural cytotoxicity receptors (NCR) and Fc $\gamma$ IIIa (CD16). KIR expression is controlled  
2 both at the genetic level and by regulatory mechanisms that lead to a stochastic expression of  
3 KIR in a single NK cell. Several studies have unraveled the impact of particular HLA and  
4 KIR genotypes with regard to HIV infection [16, 17](reviewed in [18]). We investigated NK  
5 cell receptor repertoire and function in Vietnamese EU [19, 20]. NK cell cytotoxic activity  
6 against K562 target cells was strongly enhanced in EU, as compared to exposed infected  
7 individuals who eventually seroconverted or to low-risk controls (Fig 2A). Interestingly, in  
8 contrast to the control groups, significant cytolytic activity was observed against the NK-  
9 resistant Daudi cell line, suggesting that EU NK cells are activated *in vivo*. NK cell from EU  
10 were also able to produce cytokines (IFN- $\gamma$  and TNF- $\alpha$ ) and chemokines (CCL3-5), even in  
11 the absence of any exogenous stimulation (TNF- $\alpha$  and CCL3 are shown in Fig. 2B) [19].  
12 Accordingly, NK cytotoxic granule exocytosis, as evaluated by the measurement of CD107a  
13 cell surface expression, showed that NK cell activation could be detected in EU even in the  
14 absence of target cell stimulation (Fig 2C) [20]. Remarkably, the kinetics of unstimulated NK  
15 cell degranulation was more rapid in EU than in controls, indicating an ability of EU to exert  
16 NK effector functions within a very short delay (Fig 2D).

17 Analysis of KIR transcript in EU showed enhanced KIR3DS1/KIR3DL1 and  
18 NKG2C/NKG2A ratios in favor of an activated profile of NK cells [20]. We also evidenced a  
19 higher representation of EU individuals expressing the inhibitory KIR2DL3 receptor in the  
20 absence of its KIR2DL2 allelic counterpart in individuals bearing the HLA class I C1 allele  
21 [20]. Since KIR2DL3 has lower affinity than KIR2DL2 for the HLA-C1 ligand, this should  
22 confer lower HLA inhibitory signaling. Interestingly, homozygosis for KIR2DL3 in  
23 combination with HLA-C1 has been reported to be protective in hepatitis C infection [21].  
24 The balance of expression patterns in favor of activating NKR expression and of KIR  
25 combinations with lower inhibitory potential may underlie the enhanced NK cell activity

1 observed in EU. These NKR expression patterns in EU may be the result of a selection of NK  
2 receptor-ligand combination repertoires driven by other viral challenges including HCV and  
3 HBV to which EU were highly exposed.

#### 4 **Concluding remarks**

5 The reduced susceptibility to HIV-1 infection *in vitro*, associated with low levels of CD4+ T  
6 cell activation and to genuine mechanisms of viral restriction, as well as the enhanced NK cell  
7 activity that we found in different groups of EU sustains the concept of a contribution of  
8 innate anti-viral responses to the resistance against HIV infection. The intrinsic resistance of  
9 CD4+ T cells in some EU, associated or not to mutations of the CCR5 receptor genes,  
10 indicates a prominent role of the genetic background in these individuals. This is also  
11 suggested by the association of the constitutive activation of NK cells in Vietnamese EU with  
12 particular patterns of NKR expression. Although NKR expression may be epigenetically  
13 regulated by environmental factors, including prior infectious challenges, different repertoires  
14 were found in EU and in HIV-1 infected IDU, with similar prevalence of viral infections,  
15 pointing out the role of genetic differences in providing EU with a better armed immune  
16 response to resist encounters with HIV. Other studies have also reported increased ratios of  
17 activating vs inhibitory NKR expression in EU or KIR profiles consistent with enhanced NK  
18 cell function, including higher representation of KIR3DS1 and KIR2DL3 homozygosity [20-  
19 23]. It should be underlined, however, that the heterogeneity of NKR repertoires and of the  
20 mechanisms of resistance to HIV-1 in CD4+ T cells, as well as of other immune mechanisms  
21 of protection observed among EU [7] are not in favor of a single, general mechanism of  
22 resistance.

23 Our researches carried out on limited numbers of EU from Africa and South East Asia  
24 havebrought to light some informative issues concerning the mechanisms that may contribute  
25 to resistance in such high-risk individuals. However such studies remain difficult to conduct

1 and may even raise ethical concerns associated with the reduction of risk behaviors that  
2 hopefully result from better medical counseling linked with such protocols in developing  
3 countries. In Cambodia, for example, 93% of serodiscordant couples declared as having safe  
4 sex at their second visit 6 months after recruitment, which is an achievement in terms of  
5 research protocol impact [9]. IDU in HCMC finally declared that they used disposable  
6 syringes or stopped drug use. Therefore, EU become unexposed, and, unless genetically  
7 determined, the mechanisms of resistance may vanish with time [22, 23]. Some groups of EU,  
8 such as sex workers or drug users, are vulnerable populations and particular ethical care has  
9 to be taken at recruitment and during follow-up, because these individuals may be easily  
10 influenced and may be the object of various environmental pressures that may lead to their  
11 loss during follow-up. Problems related to specific socio-political conditions of the country in  
12 which the study is carried out may also jeopardize the study itself. For example, political  
13 instability in CAR (especially a political coup in March 2003) caused the flight of urban  
14 populations, including many families recruited in our study, and hindered and finally blocked  
15 the research.

16 Nevertheless, the study of cohorts of individuals who develop mechanisms to resist HIV  
17 infection remains a major challenge to understand the basis of « natural resistance » and may  
18 orient the design of novel therapeutic or vaccine approaches, based on manipulating host  
19 defenses to provide better protection against HIV. Vaccine strategies that induce target cell  
20 resistance to HIV by stimulating  $\beta$ -chemokine production and/or increasing intracellular  
21 levels of restriction factors have been proposed [24]. NK activating receptor triggering or  
22 inhibitory ligand/receptor blockade may constitute potential ways to modulate NK cell  
23 activity. Alternatively, vectors that have an impact on the NK cell repertoire or cytokines,  
24 such as IL-21 or IL-15, as adjuvant may be used to induce or improve NK cell response [25,  
25 26].

## 1 **Acknowledgements**

2 We thank Arnaud Fontanet, Institut Pasteur, for the estimation of the cumulative risk of  
3 infection among EU from Cambodia and CAR, and Katherine Kean for English editing of the  
4 manuscript.

## 1 **References**

- 2 1. Royce RA, Sena A, Cates W, Jr. and Cohen MS. Sexual transmission of HIV. *N Engl J*  
3 *Med* **1997**;336:1072-8.
- 4 2. Fowke KR, Nagelkerke NJ, Kimani J, et al. Resistance to HIV-1 infection among  
5 persistently seronegative prostitutes in Nairobi, Kenya. *Lancet* **1996**;348:1347-51
- 6 3. Samson M, Libert F, Doranz BJ, et al. Resistance to HIV-1 infection in caucasian  
7 individuals bearing mutant alleles of the CCR-5 chemokine receptor gene. *Nature*  
8 **1996**;382:722-725
- 9 4. Langlade-Demoyen P, Ngo-Giang-Huong N, Ferchal F and Oksenhendler E. Human  
10 immunodeficiency virus (HIV) nef-specific cytotoxic T lymphocytes in noninfected  
11 heterosexual contact of HIV-infected patients [see comments]. *Journal of Clinical*  
12 *Investigation* **1994**;93:1293-7
- 13 5. Clerici M, Giorgi JV and Chou CC. Cell-mediated immune response to human  
14 immunodeficiency virus (HIV) type 1 in seronegative homosexual men with recent sexual  
15 exposure to HIV-1. *J Infect Dis* **1992**;165:1012-1019
- 16 6. Mazzoli S, Trabattoni D, Lo Caputo S, et al. HIV-specific mucosal and cellular immunity  
17 in HIV-seronegative partners of HIV-seropositive individuals. *Nat Med* **1997**;3:1250-7
- 18 7. Miyazawa M, Lopalco L, Mazzotta F, Lo Caputo S, Veas F and Clerici M. The  
19 'immunologic advantage' of HIV-exposed seronegative individuals. *Aids* **2009**;23:161-75
- 20 8. Truong TXL, Luong TT, Scott-Algara D, et al. CD4+ cell and CD8+ cell-mediated  
21 resistance to HIV-1 infection in exposed uninfected intravascular drug users in Vietnam.  
22 *AIDS* **2003**;17:1-10
- 23 9. Nguyen M, Pean P, Lopalco L, et al. HIV-specific antibodies but not T cell responses are  
24 associated to protection in seronegative partners of HIV-1 infected individuals in Cambodia. *J*  
25 *Acquir Immune Defic Syndr* **2006**;42:412-19

- 1 10. Begaud E, Chartier L, Marechal V, et al. Reduced CD4 T cell activation and in vitro  
2 susceptibility to HIV-1 infection in exposed uninfected Central Africans. *Retrovirology*  
3 **2006**;3:35
- 4 11. Wawer MJ, Gray RH, Sewankambo NK, et al. Rates of HIV-1 transmission per coital act,  
5 by stage of HIV-1 infection, in Rakai, Uganda. *J Infect Dis* **2005**;191:1403-9
- 6 12. Tran HK, Chartier L, Troung LX, et al. Systemic immune activation in HIV-1-exposed  
7 uninfected Vietnamese intravascular drug users. *AIDS Res Hum Retroviruses* **2006**;22:255-  
8 61
- 9 13. Card CM, McLaren PJ, Wachihi C, Kimani J, Plummer FA and Fowke KR. Decreased  
10 immune activation in resistance to HIV-1 infection is associated with an elevated frequency  
11 of CD4(+)CD25(+)FOXP3(+) regulatory T cells. *J Infect Dis* **2009**;199:1318-22
- 12 14. Liu R, Paxton WA, Choe S, et al. Homozygous defect in HIV-1 coreceptor accounts for  
13 resistance of some multiply-exposed individuals to HIV-1 infection. *Cell* **1996**;86:367-377
- 14 15. Saez-Cirion A, Versmisse P, Truong LX, et al. Persistent resistance to HIV-1 infection in  
15 CD4 T cells from exposed uninfected Vietnamese individuals is mediated by entry and post-  
16 entry blocks. *Retrovirology* **2006**;3:81
- 17 16. Martin MP, Gao X, Lee JH, et al. Epistatic interaction between KIR3DS1 and HLA-B  
18 delays the progression to AIDS. *Nat Genet* **2002**;31:429-34
- 19 17. Martin MP, Qi Y, Gao X, et al. Innate partnership of HLA-B and KIR3DL1 subtypes  
20 against HIV-1. *Nat Genet* **2007**;39:733-40
- 21 18. Carrington M, Martin MP and van Bergen J. KIR-HLA intercourse in HIV disease.  
22 *Trends Microbiol* **2008**;16:620-7
- 23 19. Scott-Algara D, Truong LX, Versmisse P, et al. Cutting edge: increased NK cell activity  
24 in HIV-1-exposed but uninfected Vietnamese intravascular drug users. *J Immunol*  
25 **2003**;171:5663-7

- 1 20. Ravet S, Scott-Algara D, Bonnet E, et al. Distinctive NK-cell receptor repertoires sustain  
2 high-level constitutive NK-cell activation in HIV-exposed uninfected individuals. *Blood*  
3 **2007**;109:4296-305
- 4 21. Knapp S, Warshow U, Hegazy D, et al. Consistent beneficial effects of killer cell  
5 immunoglobulin-like receptor 2DL3 and group 1 human leukocyte antigen-C following  
6 exposure to hepatitis C virus. *Hepatology* **2009**, [Epub ahead of print]
- 7 22. Mazzoli S, Lopalco L, Salvi A, et al. Human immunodeficiency virus (HIV)-specific IgA  
8 and HIV neutralizing activity in the serum of exposed seronegative partners of HIV-  
9 seropositive persons. *Journal of Infectious Diseases* **1999**;180:871-875
- 10 23. Kaul R, Rowland-Jones SL, Kimani J, et al. Late seroconversion in HIV-resistant Nairobi  
11 prostitutes despite pre-existing HIV-specific CD8(+) responses. *J Clin Invest* **2001**;107:341-  
12 349.
- 13 24. Lehner T, Wang Y, Pido-Lopez J, Whittall T, Bergmeier LA and Babaahmady K. The  
14 emerging role of innate immunity in protection against HIV-1 infection. *Vaccine*  
15 **2008**;26:2997-3001
- 16 25. Capitini CM, Fry TJ and Mackall CL. Cytokines as Adjuvants for Vaccine and Cellular  
17 Therapies for Cancer. *Am J Immunol* **2009**;5:65-83
- 18 26. Scott-Algara D, Mancini-Bourgine M, Fontaine H, Pol S and Michel ML. Changes to the  
19 natural killer cell repertoire after therapeutic hepatitis B DNA vaccination. *PLoS One*; **2010**,  
20 5:e8761

21

22


## Figure Legends

**Figure 1.** **A.** We identified 3 phenotypes of HIV-1 restriction in CD4<sup>+</sup> T cells from Vietnamese EU: in three cases the restriction affected entry of R5 HIV-1 and was associated with low surface expression of heterozygous CCR5 co-receptor variants (G106R, C178R and C269F) (phenotype 1); in two other cases, R5 HIV-1 infection was inhibited due to an enhanced sensitivity to the blocking effect of endogenously produced  $\beta$ -chemokines (phenotype 2). In one case (W276) the restriction affected not only HIV-1 but also other retroviruses, and was due to a post-entry block of viral replication (phenotype 3). **B.** Primary CD4<sup>+</sup> T cells from the W276 EU and a CD4<sup>+</sup> T cell clone derived from these cells were highly resistant to infection with HIV-1 particles, pseudotyped with BaL or VSV-G fusion proteins, or VSV-G-pseudotyped SIVmac particles. **C.** The formation of heterokaryons (indicated by the appearance of double positive CD4<sup>+</sup> CXCR4<sup>+</sup> cells, circles) between a resistant T cell clone not expressing CXCR4 and a replication competent CD4<sup>neg</sup> T cell line was accompanied by a significant increase of HIV-1 infection (bars), suggesting that a cellular co-factor needed for viral replication was lacking in the resistant T cell clone and was provided by the T cell line.

**Figure 2 :** **A.** EU individuals have a higher cytotoxic potential to lyse K562 and DAUDI target cells compared to both healthy donors (HD) who are at low risk of HIV exposure and seroconverted IDU individuals tested before (EbSC) or after seroconversion (EaSC) . PBMCs from EU, HD or IDU were incubated with K562 or Daudi cell lines at an effector to target cell ratio of 50:1 and lysis was tested by chromium release assays. Significant differences are indicated. **B.** Spontaneous production of TNF- $\alpha$  and CCL33 by NK cells from EU, HD or IDU tested before seroconversion evaluated in the absence of an extrinsic stimulus are shown. Intracellular staining of CD3-CD16<sup>+</sup>CD56<sup>+</sup> NK cells for TNF- $\alpha$  and CCL3 is shown for each group. Significant differences are indicated. **C.** CD107a cell surface expression was analyzed




within the CD3-CD56<sup>+</sup> NK cell subset both in the absence of stimulation or after 3 hours of K562 target cell stimulation (effector to target ratio =2). The percentage of CD3-CD56<sup>+</sup> NK cells expressing the CD107a cell surface marker were used to compare NK cell granule exocytosis potential in HD, EU, and EI individuals. In the absence of exogenous stimulation, granule exocytosis could only be observed in NK cells from EU, and not from low-risk healthy donors (HD) or IDU seroconverters (exposed infected, EI). D. Kinetics of NK cell degranulation reveals early constitutive activation in EU. NK cell CD107a degranulating activity was measured at different time points (10 min, 60 min and 120 min) using unstimulated NK cells from EU , healthy donors (HD) and an HIV-infected IDU (exposed infected, EI).




1  
2

**Figure 1**


A



C



B



D




Figure 2