

**Genetic analysis of the compatibility between
polymerase proteins from human and avian strains of
influenza A viruses**

Nadia Naffakh, Pascale Massin, Nicolas Escriou, Bernadette
Crescenzo-Chaigne, Sylvie Van Der Werf

► **To cite this version:**

Nadia Naffakh, Pascale Massin, Nicolas Escriou, Bernadette Crescenzo-Chaigne, Sylvie Van Der Werf. Genetic analysis of the compatibility between polymerase proteins from human and avian strains of influenza A viruses. *Journal of General Virology, Microbiology Society*, 2000, 81 (5), pp.1283 - 1291. <10.1099/0022-1317-81-5-1283>. <pasteur-01611108>

HAL Id: pasteur-01611108

<https://hal-pasteur.archives-ouvertes.fr/pasteur-01611108>

Submitted on 5 Oct 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Genetic analysis of the compatibility between polymerase proteins from human and avian strains of influenza A viruses

Nadia Naffakh, Pascale Massin, Nicolas Escriou, Bernadette Crescenzo-Chaigne and Sylvie van der Werf

Unité de Génétique Moléculaire des Virus Respiratoires, URA CNRS 1966, Institut Pasteur, 25 rue du Dr Roux, 75724 Paris Cedex 15, France

In order to determine how efficiently the polymerase proteins derived from human and avian influenza A viruses can interact with each other in the context of a mammalian cell, a genetic system that allows the *in vivo* reconstitution of active ribonucleoproteins was used. The ability to achieve replication of a viral-like reporter RNA in COS-1 cells was examined with heterospecific mixtures of the core proteins (PB1, PB2, PA and NP) from two strains of human viruses (A/Puerto Rico/8/34 and A/Victoria/3/75), two strains of avian viruses (A/Mallard/NY/6750/78 and A/FPV/Rostock/34), and a strain of avian origin (A/Hong Kong/156/97) that was isolated from the first human case of H5N1 influenza in Hong Kong in 1997. In accordance with published observations on reassortant viruses, PB2 amino acid 627 was identified as a major determinant of the replication efficiency of heterospecific complexes in COS-1 cells. Moreover, the results showed that replication of the viral-like reporter RNA was more efficient when PB2 and NP were both derived from the same avian or human virus or when PB1 was derived from an avian virus, whatever the origin of the other proteins. Furthermore, the PB1 and PB2 proteins from the A/Hong Kong/156/97 virus exhibited intermediate properties with respect to the corresponding proteins from avian or human influenza viruses, suggesting that some molecular characteristics of PB1 and PB2 proteins might at least partially account for the ability of the A/Hong Kong/156/97 virus to replicate in humans.

Introduction

Influenza A viruses have been isolated from a wide range of animal species. Aquatic birds, in which all 15 haemagglutinin (HA) subtypes have been found, are believed to be the reservoir of influenza A viruses from which new virus subtypes can episodically be transmitted to new hosts (Webster & Bean, 1998). Generally, avian influenza viruses (AIVs) replicate poorly in humans (Beare & Webster, 1991) and in other primates (Murphy *et al.*, 1982). In contrast, reassortants that derive their gene segments from both human and avian viruses are known to be potentially threatening to humans, as such viruses were responsible for the influenza pandemics in 1957 and 1968 (Scholtissek *et al.*, 1978). Segments of avian origin

not only encoded novel neuraminidase (NA) and/or HA molecules to which the human population had no immunity, but also the RNA polymerase PB1 (Kawaoka *et al.*, 1989). Several observations suggest that avian and human viruses may have reassorted in pigs and that the resulting reassortants may have been transmitted from pigs to humans (Scholtissek *et al.*, 1998). In 1997, a highly pathogenic H5N1 virus that circulated in chickens in Hong Kong infected at least 18 persons and caused six deaths (Centers for Disease Control & Prevention, 1998). Each case appeared to result from independent transmission of the virus from poultry to humans. There was no convincing evidence for human-to-human transmission and since the mass culling of poultry in Hong Kong, no new human cases of H5N1 virus infection have been recorded. However, related viruses of the H9N2 subtype, with internal genes that showed great similarity with those of the H5N1 Hong Kong viruses, were responsible for two cases of human infection in Hong Kong in 1999, and are still circulating in poultry in Hong Kong (Guan *et al.*, 1999). Therefore, the

Author for correspondence: Sylvie van der Werf.

Fax +33 1 4061 3241. e-mail svdwerf@pasteur.fr

The EMBL accession numbers of the sequences reported in this paper are AJ243990–AJ243994.

possibility of a productive reassortment between these AIVs and human influenza viruses, which may yield a pandemic reassortant, is a matter of concern. This situation highlights the need for a better understanding of the molecular basis of infectivity of avian/human reassortant influenza A viruses.

For all viral proteins, distinct evolutionary lineages of viral genes are associated with human and avian influenza A viruses (Webster & Bean, 1998). A number of reassortant viruses which contain human influenza HA and NA genes and various combinations of avian or human influenza internal genes have been generated by coinfection of cultured cells and evaluated for their ability to replicate *in vitro* and/or *in vivo* (Tian *et al.*, 1985; Buckler-White *et al.*, 1986; Clements *et al.*, 1986, 1992; Snyder *et al.*, 1987). This has led to the idea that specific constellations of gene segments encoding the internal proteins could be a determinant for the ability of reassortant viruses to replicate efficiently in mammalian cells and/or in the respiratory epithelium of primates. As far as the core proteins (PB1, PB2, PA and NP) are concerned, studies on reassortant viruses have shown that efficient virus replication is observed for some, but not all, constellations of human and avian virus-derived proteins, suggesting that specific interactions between the core proteins are required. On the other hand, co-immunoprecipitation and two-hybrid experiments have indicated that molecular assembly of the polymerase complex relies on PB1–PB2 and PB1–PA interactions, whereas no direct PB2–PA interaction has been detected (Digard *et al.*, 1989; Gonzalez *et al.*, 1996; Toyoda *et al.*, 1996; Zürcher *et al.*, 1996). Biochemical and genetic data have established that the PB1 subunit displays RNA-dependent RNA polymerase activity (Braam *et al.*, 1983; Biswas & Nayak, 1994) and that the PB2 subunit is involved in the cap-snatching mechanism by which viral mRNA synthesis is initiated (Blaas *et al.*, 1982; Braam *et al.*, 1983; Blok *et al.*, 1996; Shi *et al.*, 1996), while the only biochemical activity found to be associated with PA is the induction of proteolysis (Sanz-Ezquerro *et al.*, 1995). Studies on reassortant viruses have clearly shown that the origin of PB2 is a determinant for their efficiency of replication in mammalian cells (Almond, 1977; Clements *et al.*, 1992). A reassortant virus that derived its PB2 gene from the AIV A/Mallard/NY/6750/78 and its remaining genes from a human virus failed to produce plaques in mammalian cells and its replication was restricted in squirrel monkeys and humans (Clements *et al.*, 1992). Analysis of phenotypic revertants of the reassortant virus indicated that amino acid 627 of PB2 was a major determinant of the phenotype (Subbarao *et al.*, 1993). However, a major limitation from studies on reassortant viruses stems from the fact that all possible constellations among internal genes could not be examined as viruses with certain reassortant genotypes were not isolated, possibly because of a very low replication efficiency.

To address the question as to how efficiently the polymerase proteins derived from a human and an avian virus can interact with each other in the context of a mammalian cell, we

made use of the genetic system described by Pleschka *et al.* (1996) for the *in vivo* reconstitution of functional ribonucleoproteins (RNPs). This was achieved by transfecting into COS-1 cells a plasmid that directed the expression of a viral-like reporter RNA together with plasmids from which the four core proteins were expressed. The synthetic reporter RNA contained the CAT reporter gene sequence in antisense orientation flanked by the non-coding sequences that serve as promoters for the viral polymerase, which are highly conserved among influenza A viruses (Desselberger *et al.*, 1980). Homospecific as well as heterospecific RNPs were reconstituted by expressing proteins derived (i) from two strains of human viruses, A/Puerto Rico/8/34 (which is anterior to the 1957 reassortment event) and A/Victoria/3/75 (which derives from the 1968 reassortant virus); (ii) from two strains of AIVs, the A/Mallard/NY/6750/78 strain, which was used in the reassortment experiments mentioned above (Clements *et al.*, 1992; Subbarao *et al.*, 1993), and A/FPV/Rostock/34, which is known to be highly pathogenic in chickens; (iii) from A/Hong Kong/156/97, the H5N1 strain of avian origin that was isolated from the index case in Hong Kong (Claas *et al.*, 1998; Subbarao *et al.*, 1998). This experimental system allowed us to examine how efficiently all possible heterospecific mixtures of the four core proteins permitted transcription/replication of the viral-like reporter RNA.

Methods

■ **Plasmids.** Plasmid pHMG contains a mouse hydroxymethylglutaryl coenzyme A reductase (HMG) promoter and a simian virus 40 polyadenylation signal. It was kindly provided by J. Pavlovic (Institut für Medizinische Virologie, Zurich, Switzerland) together with plasmids pHMG-PR8-PB1, pHMG-PR8-PB2, pHMG-PR8-PA and pHMG-PR8-NP, which were used to express the PB1, PB2, PA and NP of influenza virus A/Puerto Rico/8/34 (PR8) under the control of the HMG promoter. Plasmids pGEM-VIC-PB1, pGEM-VIC-PB2, pGEM-VIC-PA and pGEM-VIC-NP were kindly provided by A. Portela (Instituto de Salud Carlos III, Madrid, Spain). The cDNAs encoding the PB1, PB2, PA and NP of influenza virus A/Victoria/3/75 (VIC) were transferred from the pGEM-3-derived plasmids to the *EcoRV* site of plasmid pHMG. Plasmid pPoll-CAT-RT (Pleschka *et al.*, 1996) was kindly provided by P. Palese (Mount Sinai Medical Center, New York, USA). It contains the CAT gene open reading frame in negative orientation, flanked by the 5' and 3' extremities of the NS segment of influenza virus A/WSN/33. Expression of the influenza virus-like RNA is driven by a truncated human RNA Pol I promoter. The correct 3' end is ensured by the use of the hepatitis delta virus ribozyme sequence.

■ **Molecular cloning and sequencing of cDNAs.** Virus isolates A/FPV/Rostock/34 (FPV), A/Mallard/NY/6750/78 (MAL) and A/Hong Kong/156/97 (HK) were grown in 11-day-old embryonated chicken eggs (FPV, MAL) or on MDCK cells (HK). Viral genomic RNA was extracted using 5 M guanidium isothiocyanate and phenol. Full-length cDNAs were prepared by reverse-transcription of the viral RNA using a primer complementary to the 12 bp conserved at the 3' end of each viral segment and 20 U avian myeloblastosis virus reverse transcriptase with incubation at 42 °C for 1 h followed by incubation at 55 °C for 15 min. Amplification was performed next using primers

specific for the coding sequences of PB1, PB2, PA or NP, with additional nucleotides providing an *EcoRV* site at their 5' end. The exact sequence of the primers can be obtained from the authors upon request. The PCR conditions were optimized for each sequence to be amplified: either *Pwo* or Expand High Fidelity polymerase (Boehringer Mannheim) were used, for 30 cycles, each consisting of 1 min at 94 °C, 1 min at 50–60 °C (depending on the sequence), 2–9 min (depending on the sequence) at 72 °C. The PCR products were digested with *EcoRV* and cloned into the *EcoRV* site of plasmid pGEM5z(f+). At least three positive clones were sequenced using a dRhod or a Big Dye terminator sequencing kit (Perkin Elmer) and run on a 377 automated sequencer (Perkin Elmer). Clones without PCR-induced mutations were selected, and the *EcoRV* fragments derived from these were sub-cloned into the *EcoRV* site of plasmid pHMG.

■ **Construction of mutants.** The PCR-based protocol from Tomic *et al.* (1990) was used to introduce point mutations at residue 627 (Gln → Lys) of MAL- and FPV-PB2. Primers were designed to provide two intermediate PCR products with a region of common sequence containing the mutation at codon 627 (GAA or GAG → AAA) and a site for *Bsal*. The exact sequence of the primers can be obtained from the authors upon request. Plasmids pGEM5z-MAL-PB2 or pGEM5z-FPV-PB2 were used as the starting template. Proof-reading *Pwo* polymerase (Boehringer Mannheim) was used for PCR amplification, for 20 cycles, each consisting of 1 min at 94 °C, 1 min at 50 °C, 90 s at 72 °C. The PCR products were digested with *Bsal* and ligated to each other. The ligation products were sub-cloned either in pGEM5z-MAL-PB2 using *MfeI* and *EcoRI* restriction sites or in pGEM5z-FPV-PB2 using two *EcoRI* sites. Positive clones were sequenced as previously indicated. Clones bearing the site-directed mutation but no unanticipated nucleotide change were selected, and the *EcoRV* fragments derived from these were sub-cloned into the *EcoRV* site of plasmid pHMG.

■ **Transfections and CAT assays.** COS-1 cells were grown in Dulbecco's modified Eagle medium (DMEM) containing 10% foetal calf serum (FCS). Subconfluent monolayers of COS-1 cells (3×10^5 cells in 35 mm dishes) were transfected using the Fugene 6-mediated method (Boehringer Mannheim). For the transient replication assay, a mixture of plasmids pHMG-PB1, -PB2, -PA, -NP (1; 1; 1; 2 µg) and pPoll-CAT-RT (1 µg) was transfected. Cells were incubated at 37 °C in DMEM containing 10% FCS until harvest. At 48 h post-transfection, cell extracts were prepared in 500 µl of lysis buffer and tested for CAT levels using the CAT ELISA kit (Boehringer Mannheim), which allowed detection of 0.05 ng/ml CAT. All experiments were done in duplicate.

■ **Western blotting.** For Western blot assays, total cell extracts were used. Transfected cells were resuspended directly in 300 µl of sample loading buffer, the proteins were resolved by electrophoresis in SDS-polyacrylamide gels and transferred onto a nitrocellulose membrane (Biorad). The filters were incubated for 1 h at room temperature or overnight at 4 °C with primary antibodies diluted in PBS containing 0.1% BSA (Sigma) and 0.25% Tween 20. Rabbit antisera directed against the PB1, PB2, and PA proteins of VIC virus were kindly provided by J. Ortin (Universidad Autonoma de Madrid, Spain) and were diluted 1/1000 (PB1, PA) or 1/5000 (PB2). For NP, a mouse antiserum was prepared by repeated intramuscular injection of plasmid pHMG-PR8-NP in mice, and used at a 1/500 dilution. After washing, the filters were incubated for 1 h at room temperature with peroxidase-conjugated secondary antibodies (donkey anti-rabbit-peroxidase or goat anti-mouse-peroxidase, Amersham) diluted 1/10000 in PBS containing 0.1% BSA and 0.25% Tween 20. After washing, the filters were incubated with peroxidase substrate (ECL Plus System, Amersham), scanned using a

STORM820 optical scanner (Molecular Dynamics) and analysed using the Image Quant program (Molecular Dynamics).

Results

Functional expression of the core proteins of human and avian influenza viruses

The cDNAs of the core proteins (PB1, PB2, PA and NP) of the AIV strains FPV, MAL and HK were cloned into the pHMG expression vector as described in Methods, and their sequences were determined. Few or no amino acid differences were found with previously published sequences for MAL proteins (MAL-PB2: Pro → Ser 409, Ala → Gly 590; MAL-NP: Pro → Ala 336) and HK proteins (HK-PB1: Arg → Lys 386; HK-PB2: Met → Val 202). In contrast, FPV-PB2, -PA and -NP sequences determined by us showed, respectively, five (out of 759), 15 (out of 716) and four (out of 498) amino acid differences as compared to the previously published sequences; for this reason they were submitted to the EMBL database (accession number for influenza A/FPV/Rostock/34 PB2, AJ243991; influenza A/FPV/Rostock/34 PA, AJ243992; influenza A/FPV/Rostock/34 NP, AJ243993). Sequences of FPV-PB1 and MAL-PA had not been published before (EMBL accession number for influenza A/FPV/Rostock/34 PB1, AJ243990; influenza A/Mallard/NY/6750/78 PA, AJ243994). They respectively showed great similarity with available PB1 sequences (at least 87.6% at the nucleotide level and 97.3% at the amino acid level) and PA sequences (at least 89.5% at the nucleotide level and 98.2% at the amino acid level) from other avian virus strains. Alignment of all available sequences for PB1, PB2, PA or NP revealed that at specific positions some amino acids appeared to be characteristic of the human or avian origin of the sequences (Fig. 1). Interestingly, in the HK-derived sequence, human-type residues were observed at some of these specific positions (PB2 Ser 199, PB1 Arg 54, PA Leu 400 and Asn 409, NP Met 136). However, only a limited number of avian PB1, PB2 and PA sequences were available for alignment.

In COS-1 cells transfected with the pHMG-PR8-PB1, -PB2, -PA and -NP plasmids together with plasmid pPoll-CAT-RT, high levels of CAT expression were measured (2441 ± 2 ng/ml), as expected from previously published work (Pleschka *et al.*, 1996). Comparable levels of CAT expression were achieved when VIC or FPV replication complexes were reconstituted (2439 ± 101 and 2203 ± 89 ng/ml, respectively), whereas reduced levels were repeatedly observed for MAL and HK complexes (834 ± 152 and 1084 ± 161 ng/ml, respectively). When cell extracts were analysed by Western blot, using PB1-, PB2-, PA- or NP-specific antisera, all the recombinant polypeptides could be detected with their expected size (not shown). Furthermore, comparable levels of expression were observed irrespective of the virus from which the proteins were derived. The observed variations in CAT production might therefore reflect a lower specific activity of the MAL and HK complexes.

PB2 aa	9	81	199	475	588	627	702		
Human	N	M	S	M	I/V	K	R		
HK156	D	T	S	L	A	E	K		
Avian	D	T	A	L	A	E	K		
PB1 aa	54	473	576	645	741				
Human anterior to 1957	R	L/H	I	I/M	T				
Human posterior to 1957	K	V	L	V	A/S				
HK156	R	V	L	V	A				
Avian	K	V	L	V	A				
PA aa	57	65	100	312	337	382	400	409	552
Human	Q	L	A	R	S	D	L	N/I	S
HK156	R	S	V	K	A	E	L	N	T
Avian	R	S	V	K	A/T	E	PQTS	S	T
NP aa	16	31	61	136	283	313			
Human	D	K	L	M	P	Y			
HK156	G	R	I	M	L	F			
Avian	G	R	I/M	L/I	L	F			

Fig. 1. Amino acids characteristic of the human or avian origin of influenza A virus core proteins. Number of strains used in this analysis: PB1, four human anterior to 1957, 15 human posterior to 1957, 12 avian; PB2, 20 human, 14 avian; PA, 15 human, 13 avian; NP, 45 human, 53 avian. HK156, A/Hong Kong/156/97 strain. Avian sequences do not include the A/Chicken/Hong Kong/-220/97, -G23/97, -G9/97, A/Chicken/Korea/-25232/96, -38349/96, A/Duck/Hong Kong/-Y280/97, -Y439/97, A/Pigeon/Hong Kong/Y233/97 and A/Quail/Hong Kong/G1/97 sequences, which were found to be very close to HK156 sequences (Suarez *et al.*, 1998; Guan *et al.*, 1999) and could therefore be considered as possibly intermediate between pure avian and pure human viruses.




Fig. 2. Effect of the human or avian origin of PB2 on the expression of the viral-like reporter RNA. COS-1 cells were transfected in duplicate with the 16 possible combinations between plasmids encoding the four PR8- and FPV-derived core proteins, together with the pPoll-CAT-RT plasmid. At 48 h post-transfection, cell extracts were prepared and tested for the levels of CAT as described in Methods. For a given combination of FPV (F)- and PR8 (P)-derived PB1, PA and NP, the CAT levels measured when PB2 was derived from FPV (grey bars) were compared to the CAT levels measured when PB2 was derived from PR8 (black bars). The results are expressed as the average \pm SD of two independent experiments.

Functional analysis of heterospecific replication complexes

In order to determine how efficiently heterospecific mixtures of PB1, PB2, PA and NP allowed transcription/replication of the viral-like reporter RNA, heterospecific RNPs were reconstituted with proteins derived from either human (PR8, VIC), avian (FPV, MAL) or HK viruses. For a given pair of viruses, all 16 possible combinations between the two sets of pHMG-PB1, -PB2, -PA and -NP plasmids were transfected together with plasmid pPoll-CAT-RT into COS-1 cells. The efficiency with which the various reconstituted heterospecific complexes permitted transcription/replication of the viral-like reporter RNA was determined by measuring the CAT levels in

extracts prepared from transfected cells. In order to analyse separately the effects of the origin of PB2, PB1, PA or NP, we studied successively the variations of CAT levels that resulted from the permutation of one of the four proteins from one strain to another.

The human or avian origin of PB2 appeared to be a determinant for the efficiency with which heterospecific complexes allowed transcription/replication of the viral-like reporter RNA. Indeed, whatever the combination of PB1, PA and NP between PR8 and FPV viruses, high levels of CAT expression (> 600 ng/ml) were measured in cell extracts when PB2 was derived from the human virus (PR8). Furthermore, except when PB1, PA and NP were of avian origin, CAT expression was systematically higher (5–100-fold) when PB2 was derived from the human (PR8) rather than from the avian (FPV) virus (Fig. 2). Very similar results were obtained when PR8/MAL or VIC/FPV protein combinations were tested (not shown). In contrast, PB2 permutations had little effect among the MAL/FPV combinations (not shown). As amino acid 627 of PB2 (Glu in PB2 proteins of avian origin, Lys in PB2 proteins of human origin) has been identified as an important determinant of the host range of human/avian reassortant viruses (Subbarao *et al.*, 1993), we generated cDNAs encoding PB2 from MAL or FPV viruses with a Glu \rightarrow Lys substitution at residue 627 (E627K), by site-directed mutagenesis. Western blot analysis showed that the wild-type and mutant proteins were expressed at similar levels (Table 1). When the mutant MAL- or FPV-PB2 proteins were used in the replication assay, the levels of CAT expression were drastically increased as compared to the levels observed with the corresponding avian PB2 proteins (Table 1). In the case of the combination where avian PB1, human PA and avian NP were used, the levels of CAT expression reached with the E627K mutant PB2 proteins were similar to those observed with PR8-PB2 (Table 1). This confirmed that PB2 amino acid 627 is an important determinant of the ability of avian/human hetero-

Table 1. Effect of the human or avian origin of PB2 on expression of the viral-like reporter RNA

PB1	PA	NP	PB2	Western blot*	CAT (ng/ml)†	CAT (%)‡
–	–	–	–		<0.05	
PR8	PR8	PR8	PR8		2207 ± 175	100
PR8	PR8	PR8	MAL		16 ± 3	0.7 ± 0.05
PR8	PR8	PR8	MAL-E627K		618 ± 42	28 ± 2
PR8	PR8	PR8	FPV		21 ± 0.5	0.9 ± 0.05
PR8	PR8	PR8	FPV-E627K		959 ± 140	43 ± 18
PR8	PR8	PR8	HK		6 ± 1	0.3 ± 0.02
MAL	PR8	MAL	PR8		1518 ± 76	100
MAL	PR8	MAL	MAL		235 ± 22	15 ± 1
MAL	PR8	MAL	MAL-E627K		1634 ± 28	108 ± 2
FPV	PR8	FPV	PR8		886 ± 69	100
FPV	PR8	FPV	FPV		303 ± 2	34 ± 0.05
FPV	PR8	FPV	FPV-E627K		1219 ± 162	138 ± 18
HK	PR8	HK	PR8		845 ± 51	100
HK	PR8	HK	HK		434 ± 43	50 ± 5

* For Western blot analysis, COS-1 cells were transfected in duplicate with the pHMG control plasmid (first line) or the pHMG plasmids encoding either PR8-, MAL-, MAL-E627K- FPV-, FPV-E627K-, or HK-PB2.

At 48 h post-transfection, cell extracts were prepared and analysed by Western blot using a polyclonal serum specific for PB2 as described in Methods. The image obtained by scanning the blot with a STORM820 optical scanner is shown.

† CAT levels are expressed as the mean ± SD values from one representative experiment in duplicate, out of two.

‡ For a given combination of PB1, PA and NP, CAT levels measured when PB2 was of avian origin are expressed as the percentage of the CAT level measured when PB2 was derived from PR8 virus.

specific RNPs to undergo transcription/replication in mammalian cells. When PR8 and HK proteins were mixed, substitutions of PR8-PB2 by HK-PB2 generally had the same effects as substitution of PR8-PB2 by MAL- or FPV-PB2 (not shown). However, in a number of combinations, such as the avian- or HK-PB1, PR8-PA, avian- or HK-NP combination shown in Table 1, we repeatedly observed that CAT expression was reduced to a lesser extent when PR8-PB2 was replaced by HK-PB2 rather than by MAL- or FPV-PB2 (50% reduction compared to 85 and 66% reduction respectively, $P < 0.1$).

The human or avian origin of PB1 did not appear to be as strong a determinant as the origin of PB2 for the efficiency with which heterospecific complexes permitted transcription/replication of the viral-like reporter RNA. The differences between the CAT levels measured in the presence of PB1 derived from either the human PR8 or the avian FPV virus were generally less than 5-fold (Fig. 3). However, they were systematically in favour of FPV-PB1. The CAT levels observed with PR8-PB1 were especially low as compared to FPV-PB1 in the combinations in which PB2 was derived from FPV, which

could possibly result from inefficient PR8-PB1/FPV-PB2 interactions. However, the CAT levels observed with PR8-PB1 were also lower than the levels obtained with FPV-PB1 in combinations in which most of the proteins were derived from PR8 (Fig. 3). Similarly, for heterospecific replication complexes derived from the human PR8 and avian MAL viruses, the CAT levels were consistently higher with the avian MAL-PB1 relative to the human PR8-PB1 (not shown). Interestingly, permutations between VIC- and FPV-PB1, as well as between MAL- and FPV-PB1, in heterospecific VIC/FPV or MAL/FPV replication complexes, respectively, had no significant effect on CAT levels (not shown). This observation is consistent with the fact that, unlike PR8, the human VIC strain is derived from the 1968 reassortant virus whose PB1 segment was acquired from an avian virus. When proteins from PR8 and HK were mixed, for several combinations of PB2, PA and NP, the augmentation of CAT levels observed when HK-PB1 was used instead of PR8-PB1 was not as marked as the increase observed with FPV-PB1 relative to PR8-PB1 (Fig. 3*b*).

The origin of NP had only a slight effect on the functional efficiency of PR8/FPV heterospecific complexes for the




Fig. 3. Effect of the human or avian origin of PB1 on the expression of the viral-like reporter RNA. COS-1 cells were transfected in duplicate with the 16 possible combinations between plasmids encoding the four PR8- and FPV-derived core proteins, or between plasmids encoding the four PR8- and HK-derived core proteins, together with the pPoll-CAT-RT plasmid. At 48 h post-transfection, cell extracts were prepared and tested for the levels of CAT as described in Methods. (a) For a given combination of FPV (F)- and PR8 (P)-derived PB2, PA and NP, the CAT levels measured when PB1 was derived from FPV (grey bars) were compared to the CAT levels measured when PB1 was derived from PR8 (black bars). The results are expressed as the average \pm SD of one representative out of two independent experiments. (b) For a given combination of FPV (F)- and PR8 (P)-derived PB2, PA and NP, the CAT levels measured when PB1 was derived from FPV (grey bars) were compared to the CAT levels measured when PB1 was derived from PR8 (black bars = 100%). For a given combination of HK (H)- and PR8 (P)-derived PB2, PA and NP, the CAT levels measured when PB1 was derived from HK (white bars) were compared to the CAT levels measured when PB1 was derived from PR8 (black bars = 100%). The results are expressed as percentage values, and as the average \pm SD of two independent experiments.

transcription/replication of the viral-like reporter RNA. However, whatever the origin of PB1 and PA, the CAT levels appeared to be higher when NP and PB2 were derived from the same virus (Fig. 4). Homospecificity of NP and PB2 also appeared to result in enhanced CAT levels when proteins from PR8 and MAL (not shown) or PR8 and HK (Fig. 4b) were mixed. In contrast, NP and PB2 homospecificity had no effect on CAT expression when combinations between FPV and MAL proteins were tested (not shown).




Fig. 4. Effect of the human or avian origin of NP on the expression of the viral-like reporter RNA. COS-1 cells were transfected in duplicate with the 16 possible combinations between plasmids encoding the four PR8- and FPV-derived core proteins, or between plasmids encoding the four PR8- and HK-derived core proteins, together with the pPoll-CAT-RT plasmid. At 48 h post-transfection, cell extracts were prepared and tested for the levels of CAT as described in Methods. (a) For a given combination of FPV (F)- and PR8 (P)-derived PB1, PB2 and PA, the CAT levels measured when NP was derived from FPV (grey bars) were compared to the CAT levels measured when NP was derived from PR8 (black bars). The results are expressed as the average \pm SD of one representative out of two independent experiments. (b) For a given combination of FPV (F)- and PR8 (P)-derived PB1, PB2 and PA, the CAT levels measured when NP was derived from FPV (grey bars) were compared to the CAT levels measured when NP was derived from PR8 (black bars = 100%). For a given combination of HK (H)- and PR8 (P)-derived PB1, PB2 and PA, the CAT levels measured when NP was derived from HK (white bars) were compared to the CAT levels measured when NP was derived from PR8 (black bars = 100%). The results are expressed as percentage values, and as the average \pm SD of two independent experiments.

The effect of the origin of PA on the ability of heterospecific complexes to allow transcription/replication of the viral-like reporter RNA could not be analysed because we observed that the level of proteolytic activity associated with PA (Sanz-Ezquerro *et al.*, 1995) varied from one strain to another (not shown). Since this could affect the level of accumulation of coexpressed influenza virus core proteins and reporter CAT protein, it precluded the analysis of PA permutations in the replication assay. However, it was not likely to interfere with

the results concerning the role of the other core proteins, as the origin of PA was kept unchanged when permutations of PB1, PB2 or NP were analysed.

Discussion

It was previously demonstrated that functional influenza A-like RNPs can be reconstituted *in vivo* by expressing the four core proteins (PB1, PB2, PA and NP) from cloned cDNAs together with a plasmid-derived viral-like RNA (Pleschka *et al.*, 1996). In this report we establish that this exclusively plasmid-driven system can be used for the genetic analysis of the compatibility of core proteins from different strains of influenza A viruses. Homologous complexes, derived either from human viruses (PR8 and VIC), AIVs (MAL and FPV) or a strain of avian origin that was responsible for a fatal respiratory illness in a young boy in Hong Kong in 1997 (HK), allowed efficient transcription/replication of the viral-like reporter RNA in COS-1 cells. This observation is consistent with the fact that human and avian viruses have been shown to replicate with similar efficiencies in mammalian cells (Clements *et al.*, 1986, 1992; Snyder *et al.*, 1987). It confirms that both avian and human polymerase complexes can assemble correctly (and possibly, interact efficiently with cellular factors) in mammalian cells. We examined how efficiently heterologous mixtures of PB1, PB2, PA or NP allowed transcription/replication of the viral-like reporter RNA in COS-1 cells. Important variations in the efficiency of transcription/replication were observed among the combinations of proteins derived from an avian and a human strain, which could possibly reflect some level of incompatibility between these proteins.

The origin of PB2, and especially the nature of PB2 amino acid 627, appeared to be an important determinant of the functional efficiency of heterospecific complexes in COS-1 cells. Our observations being in accordance with previous observations on human/avian reassortant viruses (Clements *et al.*, 1992; Subbarao *et al.*, 1993), we believe that they constitute a validation of our experimental approach for the identification of other molecular determinants of the compatibility between core proteins from human and avian strains of influenza A viruses. Moreover, they demonstrate that in order to manifest, the PB2-mediated restriction of transcription/replication in mammalian cells does not need the expression of viral genes other than PB1, PA and NP. Detailed analysis of the RNA species will be needed to determine precisely which transcription and/or replication process is affected by the origin of PB2. The poor efficiency of heterospecific complexes comprising a PB2 subunit of avian origin could result from an incompatibility with other subunits of human origin and/or an altered interaction with a host factor. If protein incompatibility only was involved, poor efficiency would probably characterize both an avian/human complex and its human/avian counterpart, which does not correspond to our observations. Indeed our analysis of all possible constellations among the

polymerase genes (which was not provided by studies on reassortant viruses) shows that in the context of a mammalian cell, PB2 of human origin cooperates efficiently with any combination of PB1, PA or NP proteins. If interaction of PB2 with a host factor was the only mechanism involved, avian homospecific complexes and any heterospecific complex comprising a PB2 protein of avian origin would function with an equally poor efficiency, which does not correspond to our observations either. Indeed avian-derived homospecific complexes achieved similar levels of transcription/replication of the viral-like reporter RNA as compared to human-derived complexes. On the whole, our observations suggest that the mechanism by which differences in the PB2 protein, and especially the nature of amino acid 627, alter the efficiency of transcription/replication is complex, relying on interactions of PB2 with both viral and cellular proteins. Interaction of PB2 residue 627 with a host factor is likely to be the major determinant, as the efficiencies of the MAL and FPV complexes were increased when MAL-E627K-PB2 and FPV-E627K-PB2 were combined with MAL-PB1, -PA, -NP and FPV-PB1, -PA, -NP, respectively (results not shown). Further experiments involving both mammalian and avian cell lines will probably help clarify this point.

We observed that homospecificity between PB2 and NP appeared to enhance the functional efficiency of chimeric complexes. The NP, in addition to its structural role for the formation of the RNPs, is required for efficient elongation of RNA chains and is thought to be involved in the switch from transcription to replication (Beaton & Krug, 1986; Honda *et al.*, 1988). In accordance with previously published results based on genetic (Mandler *et al.*, 1991) or biochemical (Biswas *et al.*, 1998) studies, our observations suggest that PB2–NP interactions could be involved in the transcription/replication process, and could therefore impose some constraints on reassortment events that involve the corresponding segments. Replication assays using human/avian chimeric proteins may help identify the domains of interaction between PB2 and NP. In this study, the human or avian origin of NP per se was not found to be a determinant for the efficiency of transcription/replication of the viral-like reporter RNA in COS-1 cells. In the context of a reassortant virus which derived all remaining genes from a human virus, the NP gene from the MAL avian virus was shown to specify restriction of virus replication in squirrel monkeys (Tian *et al.*, 1985; Clements *et al.*, 1992), but not in humans or in MDCK cells (Clements *et al.*, 1992). Our observations suggest that the attenuation which has been observed in squirrel monkeys does not result from an impairment of the transcription/replication process.

In the context of a mammalian cell, PB1 of avian origin appeared to cooperate with PB2, PA and NP of human origin more efficiently than PB1 of human origin appeared to cooperate with PB2, PA and NP of avian origin. In particular, in our experiments the PB2 human–PB1 avian–PA human constellation (HAH), associated either with a human or avian

NP, permitted efficient replication of the viral-like RNA reporter. This suggests that the phenotype of influenza A/Pintail/119/79 × A/Washington/897/80 reassortant viruses, which possessed the HAH constellation and were restricted in MDCK cells and in monkeys (Snyder *et al.*, 1987), was not determined by the HAH constellation only, but probably by the origin of the NS and M segments also. Our results may also be related to the fact that segments found to be of avian origin in the 1957 and 1968 pandemic viruses encoded not only the surface glycoproteins (NA and/or HA) but also PB1 (Kawaoka *et al.*, 1989). Taken together, these observations suggest that PB1 proteins of avian origin might have a higher specific activity than PB1 proteins of human origin, and that the acquisition of the PB1 segment from an avian virus may have conferred a selective advantage to the 1957 and 1968 reassortant viruses. In an alignment of PB1 sequences of human or avian origin (given the fact that only four sequences, including PR8-PB1, are available for human viruses isolated before 1957), five amino acids appeared as characteristic of the origin of the sequences (Fig. 1). Remarkably, in all PB1 sequences derived from human viruses isolated later than 1957, including six virus strains isolated in Japan in 1993–1996 (Lindstrom *et al.*, 1998), avian-type residues were found at all but one of these specific positions (Fig. 1). Conservation over a period of 28 years (running from the 1968 reassortment event up to 1996) may reflect some selective advantage conferred by these avian-type residues. In addition to the four conserved motifs found among all viral RNA-dependent polymerases (Poch *et al.*, 1989), putative nucleotide-binding domains (Asano & Ishihama, 1997) and RNA-binding sites (Li *et al.*, 1998; Gonzalez & Ortin, 1999*a, b*) have recently been identified on PB1. A number of amino acids that appear to be characteristic of the human or avian origin of the sequences are located within these domains and may therefore affect the specific activity of the PB1 proteins.

Interestingly, PB2 and PB1 proteins from the HK virus were found to behave in an intermediate fashion with respect to the corresponding proteins from human and avian viruses. Indeed, in the replication assay, both the disadvantage of HK-PB2 over PR8-PB2 and the advantage of HK-PB1 over PR8-PB1 were reduced in comparison to FPV or MAL proteins in a number of combinations. As far as PB2 is concerned, the HK strain used here shows a Glu at residue 627 (typical of avian viruses). Our data suggest that some molecular characteristics other than the nature of residue 627 could render HK-PB2 closer to human virus-derived PB2 than FPV or MAL-PB2. A potential candidate is residue 199 that is of the human type (Ser) in HK-PB2 (Fig. 1), while it is of the avian type (Ala) in the sequence of PB2 from an H5N1 virus (A/Chicken/Hong Kong/220/97) circulating in chickens in Hong Kong during the same period (Suarez *et al.*, 1998). Mutagenesis experiments will be needed to confirm this hypothesis. The fact that not only one but all three PB1, PB2 and PA proteins from HK show some characteristics of human viruses at the level of their amino acid

sequence (Fig. 1) is probably related to the fact that proteins associated physically and functionally undergo co-evolution. Our observations suggest that PB1 and PB2 proteins from HK may also have acquired some functional characteristics from human viruses, which could partially account for the ability of this virus strain to replicate in humans.

The authors are very grateful to J. Pavlovic for providing the pHMG recombinant plasmids, to A. Portela for providing the pGEM recombinant plasmids, to P. Palese for providing plasmid pPolI-CAT-RT, and to J. Ortin for providing polyclonal sera specific for PB1, PB2 and PA proteins. They are grateful to A. Hay for providing the A/FPV/Rostock/34 and A/Hong Kong/156/97 viruses, and R. Webster for providing the A/Mallard/NY/6750/78 virus. The technical assistance of Ida Rijks and Claudine Rousseaux for the production of influenza viruses is gratefully acknowledged. We thank Cécile Quemin for contribution to the initial cloning of the A/Hong Kong/156/97 cDNAs, and Marco Vignuzzi for critical reading of the manuscript.

This work was supported in part by the Ministère de l'Éducation Nationale, de la Recherche et de la Technologie (EA 302).

References

- Almond, J. W. (1977).** A single gene determines the host range of influenza virus. *Nature* **270**, 617–618.
- Asano, Y. & Ishihama, A. (1997).** Identification of two nucleotide-binding domains on the PB1 subunit of influenza virus RNA polymerase. *Journal of Biochemistry* **122**, 627–634.
- Beare, A. S. & Webster, R. G. (1991).** Replication of avian influenza viruses in humans. *Archives of Virology* **119**, 37–42.
- Beaton, A. R. & Krug, R. M. (1986).** Transcription antitermination during influenza viral template RNA synthesis requires the nucleocapsid protein and the absence of a 5' capped end. *Proceedings of the National Academy of Sciences, USA* **83**, 6282–6286.
- Biswas, S. K. & Nayak, D. P. (1994).** Mutational analysis of the conserved motifs of influenza A virus polymerase basic protein 1. *Journal of Virology* **68**, 1819–1826.
- Biswas, S. K., Boutz, P. L. & Nayak, D. P. (1998).** Influenza virus nucleoprotein interacts with influenza virus polymerase proteins. *Journal of Virology* **72**, 5493–5501.
- Blaas, D., Patzelt, E. & Kuechler, E. (1982).** Cap-recognizing protein of influenza virus. *Virology* **116**, 339–348.
- Blok, V., Cianci, C., Tibbles, K. W., Inglis, S. C., Krystal, M. & Digard, P. (1996).** Inhibition of the influenza virus RNA-dependent RNA polymerase by antisera directed against the carboxy-terminal region of the PB2 subunit. *Journal of General Virology* **77**, 1025–1033.
- Braam, J., Ulmanen, I. & Krug, R. M. (1983).** Molecular model of a eucaryotic transcription complex: functions and movements of influenza P proteins during capped RNA-primed transcription. *Cell* **34**, 609–618.
- Buckler-White, A. J., Naeve, C. W. & Murphy, B. R. (1986).** Characterization of a gene coding for M proteins which is involved in host range restriction of an avian influenza A virus in monkeys. *Journal of Virology* **57**, 697–700.
- Centers for Disease Control & Prevention (1998).** Update: isolation of avian influenza A (H5N1) viruses from humans – Hong Kong, 1997–1998. *Morbidity & Mortality Weekly Report* **46**, 1245–1247.
- Claas, E. C., Osterhaus, A. D., van Beek, R., De Jong, J. C., Rimmelzwaan, G. F., Senne, D. A., Krauss, S., Shortridge, K. F. & Webster, R. G. (1998).** Human influenza A H5N1 virus related to a

- highly pathogenic avian influenza virus [published erratum appears in *Lancet* (1998) 351, 1292]. *Lancet* **351**, 472–477.
- Clements, M. L., Snyder, M. H., Buckler-White, A. J., Tierney, E. L., London, W. T. & Murphy, B. R. (1986).** Evaluation of avian–human reassortant influenza A/Washington/897/80 × A/Pintail/119/79 virus in monkeys and adult volunteers. *Journal of Clinical Microbiology* **24**, 47–51.
- Clements, M. L., Subbarao, E. K., Fries, L. F., Karron, R. A., London, W. T. & Murphy, B. R. (1992).** Use of single-gene reassortant viruses to study the role of avian influenza A virus genes in attenuation of wild-type human influenza A virus for squirrel monkeys and adult human volunteers. *Journal of Clinical Microbiology* **30**, 655–662.
- Desselberger, U., Racaniello, V. R., Zazra, J. J. & Palese, P. (1980).** The 3' and 5'-terminal sequences of influenza A, B and C virus RNA segments are highly conserved and show partial inverted complementarity. *Gene* **8**, 315–328.
- Digard, P., Block, V. C. & Inglis, S. C. (1989).** Complex formation between influenza virus polymerase proteins expressed in *Xenopus* oocytes. *Virology* **171**, 162–169.
- Gonzalez, S. & Ortin, J. (1999a).** Distinct regions of influenza virus PB1 polymerase subunit recognize vRNA and cRNA templates. *EMBO Journal* **18**, 3767–3775.
- Gonzalez, S. & Ortin, J. (1999b).** Characterization of influenza virus PB1 protein binding to viral RNA: two separate regions of the protein contribute to the interaction domain. *Journal of Virology* **73**, 631–637.
- Gonzalez, S., Zurcher, T. & Ortin, J. (1996).** Identification of two separate domains in the influenza virus PB1 protein involved in the interaction with the PB2 and PA subunits: a model for the viral RNA polymerase structure. *Nucleic Acids Research* **24**, 4456–4463.
- Guan, Y., Shortridge, K. F., Krauss, S. & Webster, R. G. (1999).** Molecular characterization of H9N2 influenza viruses: were they the donors of the 'internal' genes of H5N1 viruses in Hong-Kong? *Proceedings of the National Academy of Sciences, USA* **96**, 9363–9367.
- Honda, A., Ueda, K., Nagata, K. & Ishihama, A. (1988).** RNA polymerase of influenza virus: role of NP in RNA chain elongation. *Journal of Biochemistry* **104**, 1021–1026.
- Kawaoka, Y., Krauss, S. & Webster, R. G. (1989).** Avian-to-human transmission of the PB1 gene of influenza A viruses in the 1957 and 1968 pandemics. *Journal of Virology* **63**, 4603–4608.
- Li, M. L., Ramirez, B. C. & Krug, R. M. (1998).** RNA-dependent activation of primer RNA production by influenza virus polymerase: different regions of the same protein subunit constitute the two required RNA-binding sites. *EMBO Journal* **17**, 5844–5852.
- Lindstrom, S. E., Hiramoto, Y., Nerome, R., Omoe, K., Sugita, S., Yamasaki, Y., Takahashi, T. & Nerome, K. (1998).** Phylogenetic analysis of the entire genome of influenza A (H3N2) viruses from Japan: evidence for genetic reassortment of the six internal genes. *Journal of Virology* **72**, 8021–8031.
- Mandler, J., Muller, K. & Scholtissek, C. (1991).** Mutants and revertants of an avian influenza A virus with temperature-sensitive defects in the nucleoprotein and PB2. *Virology* **181**, 512–519.
- Murphy, B. R., Hinshaw, V. S., Sly, D. L., London, W. T., Hosier, N. T., Wood, F. T., Webster, R. G. & Chanock, R. M. (1982).** Virulence of avian influenza A viruses for squirrel monkeys. *Infection and Immunity* **37**, 1119–1126.
- Pleschka, S., Jaskunas, R., Engelhardt, O. G., Zurcher, T., Palese, P. & Garcia-Sastre, A. (1996).** A plasmid-based reverse genetics system for influenza A virus. *Journal of Virology* **70**, 4188–4192.
- Poch, O., Sauvaget, I., Delarue, M. & Tordo, N. (1989).** Identification of four conserved motifs among the RNA-dependent polymerase encoding elements. *EMBO Journal* **8**, 3867–3874.
- Sanz-Ezquerro, J. J., de la Luna, S., Ortin, J. & Nieto, A. (1995).** Individual expression of influenza virus PA protein induces degradation of coexpressed proteins. *Journal of Virology* **69**, 2420–2426.
- Scholtissek, C., Rohde, W., von Hoyningen, V. & Rott, R. (1978).** On the origin of the human influenza subtype H2N2 and H3N2. *Virology* **87**, 13–20.
- Scholtissek, C., Hinshaw, V. S. & Olsen, C. W. (1998).** Influenza in pigs and their role as the intermediate host. In *Textbook of Influenza*, pp. 137–145. Edited by K. G. Nicholson, R. G. Webster & A. J. Hay. Oxford: Blackwell Science.
- Shi, L., Galarza, J. M. & Summers, D. F. (1996).** Recombinant-baculovirus-expressed PB2 subunit of the influenza A virus RNA polymerase binds cap groups as an isolated subunit. *Virus Research* **42**, 1–9.
- Snyder, M. H., Buckler-White, A. J., London, W. T., Tierney, E. L. & Murphy, B. R. (1987).** The avian influenza virus nucleoprotein gene and a specific constellation of avian and human virus polymerase genes each specify attenuation of avian–human influenza A/Pintail/79 reassortant viruses for monkeys. *Journal of Virology* **61**, 2857–2863.
- Suarez, D. L., Perdue, M. L., Cox, N., Rowe, T., Bender, C., Huang, J. & Swayne, D. E. (1998).** Comparisons of highly virulent H5N1 influenza A viruses isolated from humans and chickens from Hong Kong. *Journal of Virology* **72**, 6678–6688.
- Subbarao, E. K., London, W. & Murphy, B. R. (1993).** A single amino acid in the PB2 gene of influenza A virus is a determinant of host range. *Journal of Virology* **67**, 1761–1764.
- Subbarao, K., Klimov, A., Katz, J., Regnery, H., Lim, W., Hall, H., Perdue, M., Swayne, D., Bender, C., Huang, J., Hemphill, M., Rowe, T., Shaw, M., Xu, X., Fukuda, K. & Cox, N. (1998).** Characterization of an avian influenza A (H5N1) virus isolated from a child with a fatal respiratory illness [see comments]. *Science* **279**, 393–396.
- Tian, S. F., Buckler-White, A. J., London, W. T., Reck, L. J., Chanock, R. M. & Murphy, B. R. (1985).** Nucleoprotein and membrane protein genes are associated with restriction of replication of influenza A/Mallard/NY/78 virus and its reassortants in squirrel monkey respiratory tract. *Journal of Virology* **53**, 771–775.
- Tomic, M., Sunjevaric, I., Savtchenko, E. S. & Blumenberg, M. (1990).** A rapid and simple method for introducing specific mutations into any position of DNA leaving all other positions unaltered. *Nucleic Acids Research* **18**, 1656.
- Toyoda, T., Adyshev, D. M., Kobayashi, M., Iwata, A. & Ishihama, A. (1996).** Molecular assembly of the influenza virus RNA polymerase: determination of the subunit–subunit contact sites. *Journal of General Virology* **77**, 2149–2157.
- Webster, R. G. & Bean, W. J., Jr (1998).** Evolution and ecology of influenza viruses: interspecies transmission. In *Textbook of Influenza*, pp. 109–119. Edited by K. G. Nicholson, R. G. Webster & A. J. Hay. Oxford: Blackwell Science.
- Zürcher, T., de la Luna, S., Sanz-Ezquerro, J. J., Nieto, A. & Ortin, J. (1996).** Mutational analysis of the influenza virus A/Victoria/3/75 PA protein: studies of interaction with PB1 protein and identification of a dominant negative mutant. *Journal of General Virology* **77**, 1745–1749.

Received 8 November 1999; Accepted 26 January 2000