Influence of breeding sites features on genetic differentiation of *Aedes aegypti* populations analyzed on a local scale in Phnom Penh Municipality of Cambodia.

Christophe Paupy, Ngan Chantha, Karine Huber, Nicolas Lecoq, Jean-Marc Reynes, Francois Rodhain, Anna-Bella Failloux

► To cite this version:

Christophe Paupy, Ngan Chantha, Karine Huber, Nicolas Lecoq, Jean-Marc Reynes, et al.. Influence of breeding sites features on genetic differentiation of *Aedes aegypti* populations analyzed on a local scale in Phnom Penh Municipality of Cambodia.. American Journal of Tropical Medicine and Hygiene, 2004, 71 (1), pp.73-81. 10.4269/ajtmh.2004.71.73 . pasteur-01702881

HAL Id: pasteur-01702881

<https://hal-pasteur.archives-ouvertes.fr/pasteur-01702881>

Submitted on 7 Feb 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

INFLUENCE OF BREEDING SITES FEATURES ON GENETIC DIFFERENTIATION OF *Aedes aegypti* POPULATIONS ANALYZED ON A LOCAL SCALE IN PHNOM PENH MUNICIPALITY OF CAMBODIA

CHRISTOPHE PAUPY, NGAN CHANTHA, KARINE HUBER, NICOLAS LECOZ, JEAN-MARC REYNES, FRANCOIS RODHAIN, AND ANNA-BELLA FAILLOUX

Unité Insectes et Maladies Infectieuses et Ecole Pasteurienne d'Infectiologie, Institut Pasteur, Paris, France; National Malaria Center, Phnom Penh, Cambodia; Unité de Virologie, Institut Pasteur du Cambodge, Phnom Penh, Cambodia

Abstract. This study analyzed genetic differentiation of 20 *Aedes aegypti* populations collected along a street in Phnom Penh Municipality of Cambodia. Using allozyme and microsatellite variations, we demonstrated that populations were differentiated and the pattern of differentiation was dependent on the type of breeding sites. Moreover, insecticide treatments with temephos mostly affect the population functioning of discarded containers. Low gene flow detected could limit the natural diffusion of resistant populations that might instead take advantage of human displacements to spread.

INTRODUCTION

Since dengue hemorrhagic fever (DHF) was first described in the mid 1950s, the occurrence of the disease has largely spread. The number of dengue fever (DF) cases has increased worldwide and dengue infection still continues its geographic expansion over Southeast Asia, America, and the Pacific region.¹ Today, dengue is considered the most important arthropod-born viral disease, and dengue virus causes 50–100 million cases of DF and several hundred thousand cases of DHF each year.² A number of complex factors are related to the emergence and the re-emergence of dengue, particularly population growth, unplanned urbanization, and increased travel by airplane which facilitates the expansion of vectors and dispersion of viruses.

Originally, dengue was restricted to urban centers, but it appears that more and more rural areas in which high numbers of domestic habitats can support large populations of *Aedes aegypti* have to face dengue outbreaks.³ In Cambodia, as in most countries in Southeast Asia, transmission of dengue viruses to humans is ensured mainly by *Ae. aegypti* (Linné 1762). This highly anthropophilic species is closely associated with a domestic environment in which both blood sources for female mosquitoes to feed on and breeding sites to oviposit are available.⁴ Females usually disperse no more than 1 km in urban areas.^{5–8} Mosquito dispersal is an epidemiologic concern because it is the mechanism whereby females acquire and disseminate pathogens. In Phnom Penh Municipality of Cambodia, *Ae. aegypti*, which represents at least 40% of indoor resting mosquitoes,⁹ breeds in various containers because running water systems are not always available. In the city center, only low levels of genetic exchange were detected between populations separated by less than 6 km.¹⁰ Household water practices certainly influence larval distribution of the vector.¹¹ Thus, this has a significant effect on dengue risk.¹²

Since 1962, Cambodia has regularly been confronted with more severe and recurrent dengue outbreaks.¹³ After the worst recorded outbreak in 1998, which caused 16,216 DHF cases and 475 deaths,¹⁴ the National Dengue Control Program (known as Mosquitoblitz) set up by the Ministry of Health in 2001 aimed to control dengue vectors using mass larvicide applications (temephos) in drinking water containers (Chantha N, unpublished data). Before dengue vaccines or other measures such as the release of non-competent, ge-

netically modified mosquitoes become available, vector control through the use of insecticides remains the only realistic way to limit dengue outbreaks. Control measures could lead to habitat disappearance and thus affect species diversity. Species extinction in urban environments is caused mostly by insecticide uses. Acting as a powerful selection factor, insecticides lead to rapid development of resistance.¹⁵ Information on genetic variation within and between populations is critical for understanding the evolutionary history of mosquito populations experiencing different environmental constraints.^{16,17}

In previous studies, we have investigated genetic differentiation of *Ae. aegypti* collected in different cities in Cambodia (i.e., the country scale) (Paupy C and others, unpublished data) and within the city of Phnom Penh (i.e., the city scale).¹⁰ Our results showed that genetic differentiation estimated in Phnom Penh was low compared with other Asian cities such as Chiang Mai, Thailand¹⁸ or Ho Chi Minh City, Vietnam.^{16,17} In this study, we have investigated *Ae. aegypti* populations in a more limited geographic area compatible with the natural flight range of the species. We have assessed these populations using both allozyme and microsatellite variations based on genetic differentiation of populations collected in dwellings along a street in Phnom Penh, and its variation according to water storage devices and insecticide applications.

MATERIALS AND METHODS

House prospecting and sampling. Twenty samples of *Ae. aegypti* larvae and pupae were collected in May 2001 in Phnom Penh. The sampling area was restricted to a single street, Lohat Street, a typical street in the southern suburbs of Phnom Penh. Both individual concrete and wooden houses bordering Lohat Street were inspected after consent was obtained from the inhabitants, and all containers with *Ae. aegypti* larvae were recorded and sampled. For each sample, we recorded the type of breeding site and the presence or absence of insecticide (usually temephos). Collected samples were reared in insectaries and resulting adults were stored at –80°C until allozyme and microsatellite assays.

Allozyme polymorphism. Each adult mosquito was ground in 25 µL of distilled water and centrifuged for five minutes at 15,000 rpm at 4°C. The pellet was kept for extraction of DNA and the supernatant was loaded onto a 12.8% starch gel using

the Tris-maleate-EDTA (pH 7.4) buffer system. The following enzyme systems were studied: glutamate oxaloacetate transaminase (Got1 and Got2, EC 2.6.1.1.), glycerol-3-phosphate dehydrogenase (Gpd, EC 1.1.1.8.), hexokinase (Hk1, Hk2, and Hk3, EC 2.7.1.1.), malic enzyme (Me, EC 1.1.40.), malate dehydrogenase (Mdh, EC 1.1.1.37.), phosphoglucoisomerase (Pgi, EC 5.3.1.9.), and phosphoglucosmutase (Pgm, EC 2.7.5.1.).¹⁹ For each sample, 18–48 adults were assayed for seven enzyme systems that provided 11 putative genetic loci. A reference control was included in each gel corresponding to females established in an isofemale lineage of *Ae. aegypti* collected in French Polynesia.¹⁹ For each locus, alleles have been numbered according to the mobility relative to the most common allele (100) in the reference.

Microsatellite polymorphism. Extraction of DNA and amplification of microsatellites were performed as described by Huber and others.¹⁷ Five microsatellite loci were analyzed: C2A8, 34/72, T3A7, AED 19, and 38/38.¹⁷ For each sample studied (Table 1), 28–30 mosquitoes were analyzed.

Statistical analysis. Gene diversity, deviations from Hardy-Weinberg proportions, genotypic linkage disequilibrium, and genetic differentiation were analyzed using GENEPOP (version 3.3) software.²⁰ Gene diversity was calculated using allele identity method (option 5, sub-option 2). Genotypic association between pairs of loci was tested for each sample using Fisher's exact test on rank $\times$ column contingency tables (option 2). Deviations from Hardy-Weinberg proportions in each population and at each locus were investigated (option 1) using an exact approximation proposed by Haldane.²¹ Multilocus estimates of significance for HW equilibrium tests were estimated by Fisher's combined probability test.²² Heterozygote deficits or excess were tested using an exact test procedure.²³ F_{IS} and F_{ST} were calculated using the formula of Weir and Cockerham.²⁴ Genetic differentiation across populations was estimated by calculating the P value associated with the F_{ST} estimate (option 3). The overall significance of multiple tests was estimated by Fisher's combined probability test.²² Critical significance levels for multiple testing were

corrected using sequential Bonferroni procedures.²⁵ Genetic isolation by geographic distance was tested by estimating rank correlations between $F_{ST}/(1 - F_{ST})$ calculated between pairs of samples and Ln distances.²⁶ Analysis of variance (ANOVA) and Kruskal-Wallis tests were performed for mean comparisons using Epi-Info (version 6.04b) software (Centers for Disease Control and Prevention, Atlanta, GA).

RESULTS

House prospecting. Most (25 of 34) houses bordering the street were inspected, which resulted in the collection of 20 samples of *Ae. aegypti* from 14 houses (Table 1 and Figure 1). The type of breeding sites was heterogeneous: of 20 samples, 10 were collected from water storage containers (WSC) (e.g., small and large jars) and 10 from discarded containers (DC) (e.g., dish, tray, bucket, kettle, jar, tire, vase) (Table 1). In WSC, larval density (i.e., estimated number of larvae) was higher than in DC. Moreover, nine samples came from houses where temephos had been distributed during the dengue prevention campaign.

Linkage disequilibrium. When linkage disequilibrium between pairs of loci encoding allozymes was assessed, seven non-random associations were detected in 110 possible tests: Hk1-Hk2, Hk1-Hk3, and Hk2-Hk3 for samples 9 and 10, and Pgi-Mdh for sample 19. Since loci Hk1, Hk2, and Hk3 seemed to be linked statistically, only Hk1 was taken into account for further analysis. Analysis of genotypic disequilibrium between pairs of microsatellite loci showed that all loci were statistically independent from each other.

Hardy-Weinberg equilibrium. Among the 11 allozyme loci investigated, two (Gpd and Me) were monomorphic for the same allele. Loci Hk1, Mdh, Pgi, and Pgm, which segregated in all samples for three or more alleles, were considered. Of 61 tests performed for Hardy-Weinberg equilibrium, only two significant deviations ($P < 0.05$) were detected (Table 2): the first deviation concerned sample 11 for Mdh, which was due to a heterozygote excess ($F_{IS} = -0.506$), while the second

TABLE 1
Features of *Aedes aegypti* samples collected in Phnom Penh, Cambodia in May 2001

Sample no.	House no.	Type of breeding site	Volume of water in container (mL)	Use of temephos in house Yes/no*	Estimated number of larvae	Genetic variation	
						Allozyme	Microsatellite
1	1	Big jar	50	N	>300	+	+
2	3	Abandoned dish	1.5	Y	0–30	+	
3	3	Abandoned tray	1	Y	30–50	+	
4	5	Big jar	300	N	>300	+	
5	7	Big jar	50	N	100–150	+	
6	7	Big jar	20	N	150–300	+	
7	12	Bucket	1	N	0–30	+	+
8	12	Abandoned kettle	1	N	0–30	+	
9	12	Abandoned jar	1	N	30–50	+	
10	11	Tire	0.5	Y	100–150	+	
11	11	Small jar	1.5	Y	100–150	+	
12	13	Tire	1	N	100–150	+	+
13	16	Big jar	30	N	30–50	+	+
14	15	Big jar	70	Y	100–150	+	+
15	22	Big jar	60	N	0–30	+	
16	22	Small jar	15	N	100–150	+	+
17	19	Small jar	10	Y	30–50	+	+
18	25	Abandoned jar	2	Y	150–300	+	+
19	29	Flower vase	0.5	Y	150–300	+	+
20	36	Abandoned jar	5	Y	100–150	+	+

* Treated houses are houses where bags of temephos have been distributed (only jars were treated). If bags were sufficient, all water jars were treated.

FIGURE 1. Map showing the geographic location of the *Aedes aegypti* samples collected in May 2001 in the Mean Chey district of Phnom Penh, Cambodia.

corresponded to sample 3 for Pgm, which was caused by a heterozygote deficit ($F_{IS} = +0.589$). When all loci were considered together, only sample 11 showed a significant deviation corresponding to a heterozygote excess ($P < 10^{-4}$).

For microsatellites, of 34 tests conducted, eight deviations from Hardy-Weinberg proportions were detected: samples 1 and 14 for 34/72, samples 12, 14, and 16 for T3A7, and samples 1, 7, and 12 for AED19 (Table 3). All deviations were due to heterozygote deficits. No significant deviation was detected when global tests (i.e., all loci for each sample) were performed.

Gene diversity. Gene diversity (i.e., heterozygosity or average proportion of heterozygotes in a subpopulation) was estimated for all loci in each sample (Table 2). Mean heterozygosity was calculated when grouping samples according to 1) type of breeding sites and 2) presence or absence of temephos treatments. Although the mean $\pm$ SD heterozygosity seemed to be higher for samples collected in WSC (0.167 ± 0.051) than those from AR (0.148 ± 0.041), no significant difference was detected ($P = 0.37$, by ANOVA). The same tendency was observed when analyzing samples according to temephos treatments. For samples collected in treated sites, the mean $\pm$ SD heterozygosity (0.162 ± 0.021) was slightly higher than in non-treated sites (0.154 ± 0.066). However, no significant difference was found between them ($P = 0.82$, by ANOVA).

Gene diversity estimated from microsatellite markers was similar ($P = 0.53$, by ANOVA) in both types of breeding

sites: 0.228 ± 0.036 for DC and 0.248 ± 0.056 for WSC. When grouping of samples was done according to temephos treatment, no difference ($P = 0.55$, by ANOVA) was found between treated and non-treated groups: 0.235 ± 0.045 for treated sites and 0.242 ± 0.051 for non-treated sites.

Genetic differentiation. Genetic differentiation assessed using allozymes showed that samples collected in Lohat Street were significantly differentiated ($F_{ST} = 0.046$, $P < 0.0001$) (Table 4). When estimating more precisely the pattern of differentiation, samples from the two types of breeding sites (i.e., abandoned recipient DC and water storage container WSC) were significantly differentiated ($P < 0.0001$). A higher F_{ST} value was estimated for the DC group ($F_{ST} = 0.071$) compared with those calculated for the WSC group ($F_{ST} = 0.029$). When estimating differentiation between pairs of samples, θ_{ij} ranged from -0.143 to 0.259 (0.045 ± 0.051) (Appendix 1): from -0.0060 to 0.2594 for the DC group (0.070 ± 0.067), from -0.0095 to 0.1386 for the WSC group (0.027 ± 0.030), and from -0.143 to 0.1765 for heterologous combinations (AR/WSC) (0.045 ± 0.045). The mean θ_{ij-WSC} was significantly lower than mean $\theta_{ij-heterologous}$ ($P = 0.023$, by Kruskal-Wallis test). In addition, the latter was significantly lower than the mean θ_{ij-DC} ($P = 0.036$, by Kruskal-Wallis test). Of 190 combinations, 61 showed significant differentiation (24.4%) among which 18 were DC combinations, 10 were WSC combinations, and 33 were AR/WSC combinations. Thus, genetic differentiation of AR/WSC combinations was intermediate between the WSC group and the DC group.

TABLE 2

F_{IS} and deviations from Hardy-Weinberg proportions observed at four polymorphic allozyme loci in 20 *Aedes aegypti* samples collected in the Mean Chey district of Phnom Penh, Cambodia, May 2001*

Hk1													Mdh				
Sample no.	N	N _{all}	F _{all}			F _{IS}	N	N _{all}	F _{all}			F _{IS}					
			1	2	3				1	2	3						
1	48	1	0	1	0	-	48	3	0.031	0.469	0.500	-0.082					
2	16	2	0.031	0.969	0	-	16	2	0	0.533	0.467	-0.038					
3	32	1	0	1	0	-	32	2	0	0.781	0.219	+0.103					
4	48	1	0	1	0	-	48	3	0.011	0.628	0.362	+0.103					
5	48	1	0	1	0	-	48	3	0.021	0.542	0.438	-0.093					
6	48	1	0	1	0	-	48	3	0.094	0.688	0.219	-0.127					
7	28	2	0	0.982	0.018	-	28	2	0	0.696	0.304	-0.253					
8	20	3	0.025	0.950	0.025	-0.001	20	2	0	0.375	0.625	-0.371					
9	48	2	0	0.958	0.042	-0.033	48	2	0	0.787	0.213	-0.262					
10	48	2	0	0.885	0.115	-0.120	48	3	0.010	0.604	0.385	-0.007					
11	48	1	0	1	0	-	48	2	0	0.625	0.375	-0.506					
12	48	1	0	1	0	-	48	3	0.012	0.430	0.558	-0.174					
13	48	1	0	1	0	-	48	3	0.010	0.396	0.594	-0.051					
14	48	1	0	1	0	-	48	3	0.042	0.521	0.438	+0.077					
15	24	1	0	1	0	-	24	3	0.021	0.521	0.458	+0.076					
16	48	1	0	1	0	-	48	3	0.010	0.635	0.354	-0.031					
17	48	1	0	1	0	-	48	3	0.042	0.479	0.479	+0.197					
18	48	1	0	1	0	-	48	2	0	0.448	0.552	+0.212					
19	48	1	0	1	0	-	48	2	0	0.458	0.542	+0.258					
20	48	2	0	0.990	0.010	-	48	2	0	0.628	0.372	-0.037					

Sample no.	Pgi										Pgm					All loci		
	N	N _{all}	F _{all}				F _{IS}	N	N _{all}	F _{all}			F _{IS}	H _{obs}	P deficit	P excess		
			1	2	3	4				1	2	3					4	
1	48	4	0	0.089	0.911	0	-0.004	48	3	0.010	0.875	0	0.115	-0.062	0.187	0.9397	0.945	
2	16	2	0	0.094	0.906	0	+0.687	16	1	0	1	0	0	-	0.126	0.0795	0.9612	
3	32	1	0	0	1	0	-	32	2	0	0.688	0	0.313	+0.589	0.100	0.0058	0.9956	
4	48	2	0	0	0.990	0.010	-	48	3	0.042	0.896	0	0.063	+0.216	0.113	0.0156	0.09874	
5	48	3	0.031	0	0.958	0.010	-0.011	48	3	0	0.938	0.010	0.052	-0.023	0.162	0.9339	0.1200	
6	48	3	0.042	0.010	0.948	0	-0.017	48	4	0.010	0.948	0.010	0.031	-0.008	0.142	0.9381	0.0716	
7	28	2	0	0.089	0.911	0	+0.366	28	3	0.107	0.875	0	0.018	-0.054	0.186	0.4469	0.5790	
8	20	1	0	0	1	0	-	20	1	0	1	0	0	-	0.150	0.9884	0.1215	
9	48	3	0	0.031	0.958	0.010	-0.011	48	2	0	0.896	0	0.104	-0.107	0.159	1	0.0391	
10	48	2	0	0	0.969	0.031	-0.022	48	3	0.083	0.802	0	0.115	-0.114	0.237	0.9710	0.0275	
11	48	3	0	0.073	0.792	0.135	-0.122	48	4	0.021	0.813	0.010	0.156	-0.058	0.295	1	<10⁻⁴	
12	48	1	0	0	1	0	-	48	3	0	0.958	0.010	0.031	-0.011	0.140	0.9871	0.0157	
13	48	3	0	0.031	0.948	0.021	-0.017	48	2	0.073	0.927	0	0	-0.069	0.158	0.8318	0.2407	
14	48	3	0	0.031	0.948	0.021	-0.017	48	2	0	0.958	0	0.042	-0.033	0.129	0.2280	0.7507	
15	24	1	0	0	1	0	-	24	2	0	0.854	0	0.146	-0.153	0.150	0.5131	0.5847	
16	48	3	0	0.052	0.906	0.042	+0.172	48	2	0	0.969	0	0.031	-0.022	0.142	0.1649	0.8347	
17	48	4	0.021	0.052	0.917	0.010	-0.012	48	3	0.010	0.885	0	0.104	-0.107	0.187	0.1752	0.8371	
18	48	3	0	0.011	0.947	0.043	-0.018	48	2	0	0.958	0	0.042	-0.033	0.117	0.2437	0.8413	
19	48	2	0	0	0.927	0.073	-0.069	48	3	0.094	0.875	0	0.031	-0.103	0.154	0.4258	0.5786	
20	48	1	0	0	1	0	-	48	3	0.010	0.979	0	0.010	-0.005	0.109	0.7272	0.5342	

TABLE 3
F_{IS} and deviations from Hardy-Weinberg proportions observed at five microsatellite loci in 10 *Aedes aegypti* samples collected in the Mean Chey district of Phnom Penh, Cambodia, May 2001 *

T3A7														
Sample no.	34/72													
	F _{all}													
	N	N _{all}	1	2	3	4	5	6	7	8	9	10	11	F _{IS}
1	25	3	0.10	0.40	0.50	0.00	+0.599	0.432	0.023	0.182	0.159	0	0	+0.003
7	28	2	0.036	0.00	0.964	0.00	-0.019	0.317	0.05	0.40	0.117	0	0	-0.016
12	29	3	0.052	0.069	0.879	0.00	-0.083	0.5	0.052	0.034	0.103	0.017	0	-0.134
13	30	3	0.067	0.017	0.917	0.00	-0.058	0.5	0.052	0.121	0.052	0	0	-0.179
14	30	3	0.283	0.067	0.650	0.00	-0.133	0.522	0.152	0.130	0.087	0	0.043	+0.376
16	27	2	0.278	0.00	0.722	0.00	-0.368	0.463	0.037	0.185	0.093	0.019	0.037	+0.197
17	29	3	0.021	0.021	0.958	0.00	+0.500	0.370	0.196	0.152	0.043	0	0.022	+0.165
18	29	3	0.190	0.017	0.793	0.00	-0.220	0.446	0.125	0	0.107	0.036	0	+0.169
19	27	3	0.093	0.00	0.889	0.019	-0.087	0.648	0.037	0	0	0	0	-0.315
20	25	3	0.192	0.058	0.750	0.00	-0.046	0.5	0.019	0.038	0.115	0.058	0	+0.130

Sample no.	38/38																	
	AED19																	
	F _{all}																	
N	N _{all}	1	2	3	4	5	6	7	8	9	10	11	12	F _{IS}	H _{obs}	P		
1	30	1	0	0	0	1	-	25	2	0	0.680	0.320	+0.664	29	1	1	0.198	<10 ⁻⁴
7	29	2	0	0	0.017	0.983	0.000	30	2	0	0.783	0.217	+0.714	30	2	0.983	0.197	0.001
12	29	1	0	0	0	1	-	29	2	0	0.931	0.069	+1.00	30	1	1	0.199	<10 ⁻⁴
13	30	2	0	0.033	0	0.967	0.018	30	2	0	0.717	0.283	+0.114	30	1	1	0.275	0.0865
14	30	1	0	0	0	1	-	29	2	0	0.776	0.224	-0.273	29	1	1	0.283	0.0002
16	25	2	0.02	0	0	0.98	0.000	28	2	0	0.804	0.196	0.000	28	1	1	0.304	0.0028
17	24	1	0	0	0	1	-	23	3	0.22	0.848	0.130	+0.197	24	1	1	0.178	<10 ⁻⁴
18	29	1	0	0	0	1	-	30	2	0	0.967	0.033	-0.018	30	1	1	0.212	<10 ⁻⁴
19	28	1	0	0	0	1	-	26	2	0	0.769	0.231	-0.282	27	1	1	0.267	-
20	28	1	0	0	0	1	-	29	2	0	0.759	0.241	+0.076	29	1	1	0.268	0.0118

* F_{IS} = inbreeding coefficient that measures the reduction of heterozygosity in a subpopulation due to nonrandom mating; N = sample size; N_{all} = number of alleles per locus; F_{all} = allele frequencies; H_{obs} = observed heterozygosity; P deficit = probability for rejecting Hardy-Weinberg equilibrium. Significant values after correction using Bonferroni's method are shown in **bold**.

TABLE 4

F_{ST} values for estimating *Aedes aegypti* differentiation based on allozyme and microsatellite polymorphisms according to two types of breeding sites and temephos distribution in Phnom Penh, Cambodia, May 2001

Comparisons	F_{ST}												
	Allozyme						Microsatellite						
	N	Pgm	Pgi	Mdh	Hk1	All loci	N	C2A8	34/72	T3A7	AED19	38/38	All loci
All samples	20	0.050†	0.034†	0.046†	0.061†	0.046†	10	0.006‡	0.116†	0.033†	0.032†	−0.001	0.053†
Breeding site type													
DC	10	0.085§	0.027†	0.075†	0.051§	0.071†	5	−0.001	0.038‡	0.062†	0.053‡	−0.001	0.054†
WSC	10	0.024†	0.036†	0.029†	—	0.029†	5	0.008	0.166†	0.007§	0.004	—	0.052†
Temephos treatment													
Yes	9	0.074†	0.047†	0.029†	0.086†	0.045†	5	—	0.060§	0.028†	0.041‡	—	0.039†
No	11	0.021†	0.014§	0.066†	0.022‡	0.051†	5	0.003	0.188†	0.029‡	0.030§	—	0.067†

* F_{IS} = inbreeding coefficient that measures the reduction of heterozygosity in a subpopulation due to nonrandom mating; N = number of samples; Pgm = phosphoglucosomerase; Pgi = phosphoglucosomerase; Mdh = malate dehydrogenase; Hk1 = hexokinase 1; DC = discarded container; WSC = water storage container. Significant P values for homogeneity by Fisher's exact test are shown in **bold**.

† $P < 0.0001$.

‡ $P < 0.05$.

§ $P < 0.001$.

When samples were pooled according to temephos treatment (Table 4), significant differentiation was detected in both groups ($P < 0.0001$). The F_{ST} value estimated from samples collected in treated houses ($F_{ST} = 0.045$) was similar to the F_{ST} value calculated from samples in non-treated houses ($F_{ST} = 0.051$). When genetic differentiation was estimated for pairs of samples, θ_{ij} values ranged from −0.014 to 0.195 (0.046 ± 0.052) for the group corresponding to treated samples, from −0.009 to 0.208 (0.049 ± 0.051) for the group of non-treated samples, and from −0.008 to 0.259 (0.042 ± 0.049) for heterologous comparisons (treated/non-treated) (Appendix 2). Of the 61 significant combinations, 30 were treated/non-treated combinations, 19 were treated combinations, and 12 were non-treated combinations. The mean $\theta_{ij\text{-treated}}$, $\theta_{ij\text{-non-treated}}$ and $\theta_{ij\text{-heterologous}}$ were similar ($P = 0.813$, by ANOVA).

When genetic divergence was estimated according to geographic distance, the relation $F_{ST} / (1 - F_{ST}) = a + b(\ln \text{distance})$ was not significant when one considered 1) all samples collected in Lohat Street ($P = 0.662$), 2) only samples from WSC ($P = 0.797$), and 3) only samples from DC ($P = 0.629$).

When all microsatellite loci were considered, genetic differentiation estimated for the 10 samples was significant ($F_{ST} = 0.053$, $P < 0.0001$) (Table 4). When differentiation was assessed considering pairs of samples, θ_{ij} ranged from −0.0034 to 0.1241 (0.051 ± 0.036). The level of genetic differentiation within each group (DC and WSC) was significant ($P < 0.0001$) and similar (i.e., $F_{ST} = 0.054$ for the DC group and $F_{ST} = 0.052$ for the WSC group). When pairs of samples were considered, $\theta_{ij\text{-DC}}$ ranged from 0.003 to 0.110 (0.052 ± 0.039), $\theta_{ij\text{-WSC}}$ from −0.005 to 0.122 (0.052 ± 0.036), and $\theta_{ij\text{-heterologous}}$ from −0.003 to 0.124 (0.050 ± 0.037). Of 45 possible combinations, 26 were significant ($P < 0.05$) with 15 referring to heterologous combinations. When comparing mean distributions, no significant difference ($P > 0.05$, by ANOVA) was detected. Using microsatellites, no particular pattern of differentiation was detected.

Even if differentiation was significant ($P < 0.0001$), samples originated from treated houses were less differentiated ($F_{ST} = 0.0392$) than those from non-treated houses ($F_{ST} = 0.0667$) (Table 4). When examining pairs of samples (Appendix 2), θ_{ij} ranged from −0.005 to 0.124. When mean distributions of θ_{ij}

values in each group were compared, no significant difference ($P = 0.250$, by ANOVA) was detected.

When estimating genetic divergence according to geographic distance, the relationship $F_{ST} / (1 - F_{ST}) = a + b(\ln \text{distance})$ was not significant when 1) all samples ($P = 0.258$), 2) samples from WSC ($P = 0.178$), and 3) samples from DC ($P = 0.312$) were considered.

DISCUSSION

We demonstrated that (1) *Ae. aegypti* populations were highly differentiated in the Lohat street, (2) the pattern of genetic differentiation depends on the type of breeding sites (pairs of DC samples were more differentiated than an DC sample was with an WSC sample or between two WSC samples), and (3) insecticide treatment mostly affects the population functioning of DC.

Curiously, the level of genetic differentiation ($F_{ST} = 0.046$) estimated for all 20 *Ae. aegypti* samples collected in Lohat Street using allozyme is more important than that observed for populations collected in the whole municipality of Phnom Penh ($F_{ST} = 0.027$) (Paupy C and others, unpublished data). In this latter study, only WSC were sampled in the city center. When only WSC samples were taken into account in the present study, the level of genetic differentiation decreases (Paupy C and others, unpublished data). Low levels of genetic differentiation have also been recorded for *Ae. aegypti* in San Juan, Puerto, suggesting that females tend to oviposit only a few eggs in individual sites and to disperse over long distances.²⁷ Thus, even if the collected samples were a maximum of 400 meters apart, a distance compatible with *Ae. aegypti* natural flight range in cities,²⁸ mosquito differentiation remains significant along Lohat Street. More polymorphic markers such as microsatellites can detect higher level of genetic differentiation (i.e., higher F_{ST} values; see Table 4 for more details) and could be proposed to evaluate genetic structure at a street scale.

The type of breeding sites tends to influence the level of genetic differentiation. Allozyme analysis revealed that *Ae. aegypti* populations from abandoned containers (i.e., peridomestic breeding sites) were more differentiated than are those from WSC (i.e., domestic breeding site). Water jars are rarely

emptied, ensuring permanent larval production throughout the year.^{10,15} Conversely, DC harbor larvae only in the rainy season. Populations could stem from dry eggs or from eggs laid *de novo* by females from neighboring breeding sites and, particularly, from domestic jars. The ecologic constraints (e.g., small amounts of water and food) that mosquitoes from DC have to face limit larval production.²⁹ Recurrent extinction events and founder effects enhance the inter-DC differentiation. Migration is more limited between populations from DC than between those from permanent water-filled containers. Migrants can be detected between permanent jars and temporary breeding sites as demonstrated by Huber and others¹² in Ho Chi Minh City, Vietnam.

Aedes aegypti populations in Phnom Penh have been exposed to temephos every 6–7 weeks since the setting up of the campaign Mosquitoblitz in 2001. Most water storage collections were treated with the larvicide. The efficiency of these treatments was assessed by the calculation of the Breteau index (number of jars with larvae for 100 houses); this index decreased from more than 50 to less than 5 one week after insecticide application. Unfortunately, the index recovered its pretreatment level seven weeks after the end of the treatment (Chantha N, unpublished data). Populations have therefore experienced intense selection by insecticides that probably resulted in periodic population bottlenecks. Thus, genetic polymorphism is expected to decrease rapidly during insecticide use.³⁰ Mean heterozygosities of our populations are low, confirming weak gene flow between them. During the insecticidal campaign, collections for household use were mainly treated, thus becoming unsuitable for females in which to oviposit. Thus, females were compelled to migrate in the search of untreated breeding sites (non-treated jars or DC) producing a mixture of two or more subpopulations. Thus, a reduction in the number of heterozygotes can be detected (the Walhund effect).

Our study confirms the different population functioning between peridomestic DC (temporal, small effective size) and domestic water storage (permanent, large effective size). *Aedes aegypti* females tend to oviposit in many sites, preferentially in WSC; thus, a single oviposition container may contain a mixture of several families that decreases genetic differentiation between neighboring sites. Domestic containers were more subjected to insecticide treatments that induce recurrent extinctions and emergence of insecticide-resistant populations. Thus, alleles conferring insecticide resistance could be rapidly fixed and invade untreated peridomestic sites. Moreover, resistant populations could spread at a larger scale assisted by human displacements.^{31,32} Thus, insecticide treatments of WSC should be associated with destruction of peridomestic breeding sites containing a part of *Ae. aegypti* populations.

Received September 18, 2003. Accepted for publication October 30, 2003.

Acknowledgments: We are grateful to the “Fondation de France, Jeunesse Internationale” for providing a six-month fellowship to Christophe Paupy during his training at the Pasteur Institute in Cambodia. We thank Nadia Ayad, Laurence Mousson, Regine Villeret, and Marie Vazeille for technical assistance. We also thank the staff of the Unité de Virologie, Institut Pasteur du Cambodge, the National Malaria Center, and the municipality of Phnom Penh for their invaluable help in mosquito collections. We are indebted to Professor Jean-Luc Durosoir (Pasteur Institute, Paris, France), Professor Yves Buis-

son (Pasteur Institute, Phnom Penh, Cambodia), and Eliane Chungue (Pasteur Institute, Noumea, New Caledonia) for their constant support.

Financial support: This research was supported by the program ACIP “Action Concertée des Instituts Pasteur” (no. 41204).

Authors’ addresses: Christophe Paupy, Laboratoire de Microbiologie et Environnement, Institut Pasteur de La Nouvelle Calédonie, BP 61, 98845 Noumea, New Caledonia. Ngan Chantha, National Malaria Center, 372 Boulevard Monivong, Phnom Penh, Cambodia. Karine Huber, Nicolas Lecoz, and Anna-Bella Failloux, Unité Insectes et Maladies Infectieuses, Institut Pasteur, 25-28 Rue du Dr. Roux, 75724 Paris Cedex 15, France. Telephone: 33-1-40-61-36-17, Fax: 33-1-40-61-30-89, E-mail: afaillou@pasteur.fr. Jean-Marc Reynes, Unité de Virologie, Institut Pasteur du Cambodge, 5 Boulevard Monivong, Phnom Penh, Cambodia. Francois Rodhain, Ecole Pasteurienne d’Infectiologie, Institut Pasteur, 25-28 Rue du Dr. Roux, 75724 Paris Cedex 15, France.

REFERENCES

- Gubler DJ, 2002. Epidemic dengue/dengue hemorrhagic fever as a public health, social and economic problem in the 21st century. *Trends Microbiol* 10: 100–103.
- Lam SK, 1998. Emerging infectious diseases—Southeast Asia. *Emerg Infect Diseases* 4: 145–147.
- Pant CP, Jatanasen S, Yasuno M, 1973. Prevalence of *Aedes aegypti* and *Aedes albopictus* and observations on the ecology of dengue haemorrhagic fever in several areas of Thailand. *Southeast Asian J Trop Med Public Health* 4: 113–121.
- Strickman D, Kittayapong P, 1993. Laboratory demonstration of oviposition by *Aedes aegypti* (Diptera: Culicidae) in covered water jars. *J Med Entomol* 30: 947–949.
- Tripis M, Hausermann W, 1986. Dispersal and other population parameters of *Aedes aegypti* in an African village and their possible significance in epidemiology of vector-borne diseases. *Am J Trop Med Hyg* 35: 1263–1279.
- Reiter P, Amador MA, Anderson RA, Clark GG, 1995. Dispersal of *Aedes aegypti* in an urban area after blood feeding as demonstrated by rubidium-marked eggs. *Am J Trop Med Hyg* 52: 177–179.
- Ordóñez-Gonzales JG, Mercado-Hernandez R, Flores-Suarez AE, Fernandez-Salas I, 2001. The use of sticky ovitraps to estimate dispersal of *Aedes aegypti* in northeastern Mexico. *J Am Mosq Control Assoc* 17: 93–97.
- Tsuda Y, Takagi M, Wang S, Wang Z, Tang L, 2001. Movement of *Aedes aegypti* (Diptera: Culicidae) released in a small isolated village on Hainan Island, China. *J Med Entomol* 38: 93–98.
- Kohn M, 1990. A survey on indoor resting mosquito species in Phnom Penh, Kampuchea. *Folia Parasitol* 37: 165–174.
- Paupy C, Chantha N, Reynes JM, Failloux AB, 2003. Variation over space and time of *Aedes aegypti* in Phnom Penh (Cambodia): genetic structure and oral susceptibility to a dengue virus. *Genet Res* (in press).
- Moore CG, Cline BL, Ruiz-Tiben E, Lee D, Romney-Joseph H, Rivera-Correa E, 1978. *Aedes aegypti* in Puerto-Rico: environmental determinants of larval abundance and relation to dengue virus transmission. *Am J Trop Med Hyg* 27: 1225–1231.
- Huber K, Luu Le L, Tran Huu H, Tran Khan T, Rodhain F, Failloux AB, 2002. Temporal genetic variation in *Aedes aegypti* populations in Ho Chi Minh City (Vietnam). *Heredity* 89: 7–14.
- Rathavuth H, Vaughn DW, Minn K, Nimmannitya S, Nisalak A, Raengsakulrach B, Rorabaugh ML, Yuvatha K, Sophal O, 1997. Hemorrhagic fever in Cambodia is caused by dengue viruses: evidence of transmission of all four serotypes. *Southeast Asian J Trop Med Public Health* 28: 120–125.
- Chantha N, Guyant P, Hoyer S, 1999. Control of DHF outbreak in Cambodia, 1998. *Dengue Bull* 22 (<http://w3.whosea.org/DengueBulletin22/ch12.html>).
- Rawlins SC, Wan JH, 1995. Resistance in some Caribbean populations of *Aedes aegypti* to several insecticides. *J Am Mosq Control Assoc* 11: 59–65.

16. Tran Khanh T, Vazeille-Falcoz M, Mousson L, Tran Huu H, Rodhain F, Nguyen Thi H, Failloux AB, 1999. *Aedes aegypti* in Ho Chi Minh City (Vietnam): susceptibility to dengue 2 virus and genetic differentiation. *Trans R Soc Trop Med Hyg* 93: 581–586.
17. Huber K, Luu Le L, Tran Huu H, Ravel S, Rodhain F, Failloux AB, 2002. Genetic differentiation of the dengue vector *Aedes aegypti* (Ho Chi Minh City, Viet Nam) using microsatellite markers. *Mol Ecol* 11: 1629–1635.
18. Mousson L, Vazeille M, Chawprom S, Prajakwong S, Rodhain F, Failloux AB, 2002. Genetic structure of *Aedes aegypti* populations in Chiang Mai (Thailand) and relation with dengue transmission. *Trop Med Int Health* 7: 865–872.
19. Paupy C, Vazeille-Falcoz M, Mousson L, Rodhain F, Failloux AB, 2000. *Aedes aegypti* in Tahiti and Moorea (French Polynesia): isoenzyme differentiation in the mosquito population according to human population density. *Am J Trop Med Hyg* 62: 217–224.
20. Raymond M, Rousset F, 1995. GENEPOP (version 1.2): population genetics software for exact tests and ecumenicism. *J Hered* 86: 248–249.
21. Haldane JBS, 1954. An exact test for randomness of mating. *J Genet* 52: 631–635.
22. Fisher RA, 1970. *Statistical Methods for Research Workers*. 14th edition. Edinburgh: Oliver and Boyd.
23. Rousset F, Raymond M, 1995. Testing heterozygote excess and deficiency. *Genetics* 140: 1413–1419.
24. Weir BS, Cockerham CC, 1984. Estimating F-statistics for the analysis of population structure. *Evolution* 38: 1358–1370.
25. Holm S, 1979. A simple sequentially rejective multiple test procedure. *Scand J Stat* 6: 65–70.
26. Slatkin M, 1993. Isolation by distance in equilibrium and non-equilibrium populations. *Evolution* 47: 264–279.
27. Apostol BL, Black WC, Reiter P, Miller BR, 1996. Population genetics with RAPD-PCR markers: the breeding structure of *Aedes aegypti* in Puerto Rico. *Heredity* 76: 325–334.
28. Honório N, Da Costa Silva W, José Leite P, Monteiro Gonçalves J, Lounibos LP, Lourenço-de-Oliveira R, 2003. Dispersal of *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae) in an urban endemic dengue area in the state of Rio de Janeiro, Brazil. *Mem Inst Oswaldo Cruz* 98: 191–198.
29. Strickman D, Kittayapong P, 2003. Dengue and its vector in Thailand: calculated transmission risk from total pupal counts of *Aedes aegypti* and association of wing-length measurements with aspects of the larval habitats. *Am J Trop Med Hyg* 68: 209–217.
30. Lerdthusnee K, Chareonviriyaphap T, 1999. Comparison of isozyme patterns of *Aedes aegypti* populations collected from pre- and post-*Bacillus thuringiensis israelensis* treatment sites in Thailand. *J Am Mosq Control Assoc* 15: 48–52.
31. Huber K, Luu Le L, Chantha N, Failloux AB, 2003. Human displacements shape *Aedes aegypti* gene flow in Southeast Asia. *Acta Trop*: (in press).
32. Raymond M, Callaghan A, Fort P, Pasteur N, 1991. Worldwide migration of amplified insecticide resistance genes in mosquitoes. *Nature* 350: 151–153.

APPENDIX 1

θ_{ij} estimates computed using allozyme (below the diagonal) and microsatellite variations (above the diagonal) for all pairs of samples according to the breeding site categories*

Breeding site	Discarded container (DC)										Water storage container (WSC)									
	2	3	7	8	9	10	12	18	19	20	4	5	6	1	11	13	14	15	16	17
DC	2	0.1584																		
	3	0.0246	0.0749							0.0851					0.1202		0.0718		0.0476	0.0240
	8	0.0255	0.2594	0.1319																
	9	0.0849	0.0479	0.0159	0.2086															
	10	0.0317	0.0580	0.0172	0.0756	0.0401														
	12	0.0151	0.2198	0.1013	-0.0006	0.1628	0.0624													
	18	-0.0028	0.1947	0.0801	-0.0009	0.1425	0.0519	-0.0060												
	19	0.0082	0.1591	0.0561	0.0159	0.1254	0.0332	0.0092	0.0282	0.0248										
	20	0.0103	0.1392	0.0214	0.0940	0.0467	0.0335	0.0540	0.0482	0.0255										
	4	0.0134	0.0805	0.0046	0.0886	0.0292	0.0124	0.0517	0.0422	0.0143										
	5	-0.0042	0.1273	0.0324	0.0402	0.0713	0.0251	0.0144	0.0075	-0.0022	0.0026									
	6	0.0540	0.0931	0.0141	0.1689	0.0145	0.0471	0.1240	0.1090	0.0054	0.0426	0.0021								
	1	0.0094	0.1241	0.0550	0.0241	0.1028	0.0260	0.0067	0.0026	0.0376	0.0021	0.0021	0.0786							
	11	0.0359	0.0622	0.0243	0.1060	0.0503	0.0267	0.6869	0.0604	0.0581	0.0317	0.0377	0.0543	0.0381						
	13	0.0154	0.2226	0.0955	-0.0008	0.1765	0.0644	0.0019	-0.0012	0.0740	0.0656	0.0285	0.1386	0.0130	0.0808					
	14	-0.0143	0.1403	0.0330	0.0321	0.0781	0.0300	0.0109	0.0032	0.0074	0.0070	-0.0078	0.0452	0.0026	0.0381	0.0513				
	15	0.0126	0.0849	0.0403	0.0463	0.0772	0.0108	0.0188	0.0111	0.0093	0.0039	-0.0014	0.0626	-0.0095	0.0296	0.0226	0.0010			
	16	-0.0013	0.1128	0.0062	0.0930	0.0329	0.0314	0.0608	0.0428	0.0014	0.0020	0.0074	0.0151	0.0348	0.0251	0.0713	0.0056	0.0265	0.0420	
	17	-0.0020	0.1172	0.0406	0.0273	0.0898	0.0246	0.0112	0.0025	0.0270	0.0184	0.0003	0.0651	-0.0072	0.0269	0.0140	-0.0017	-0.0065	0.0230	

* Significant differentiation (after application of Bonferroni's procedure) is underlined.

APPENDIX 2

θ_{ij} estimates computed using allozyme (below the diagonal) and microsatellite variations (above the diagonal) for all pairs of samples according to temephos treatment*

Breeding site	Treated (yes)										Non-treated (no)									
	2	3	10	11	14	17	18	19	20	1	4	5	6	7	8	9	12	13	15	16
Yes	2																			
	3	0.1584																		
	10	0.0317	0.0580																	
	11	0.0359	0.0622	0.0267																
	14	-0.0143	0.1403	0.0300	0.0381					0.0054	0.0528						0.0718	0.0513		-0.0054
	17	-0.0020	0.1172	0.0246	0.0269	-0.0017	0.0646	0.0453	0.0521	0.0387	0.1225						0.0107	0.0163		0.0420
	18	-0.0028	0.1947	0.0519	0.0604	0.0032	0.0025	0.0318	0.0554	0.0255	0.1241						0.0029	0.0584		0.0280
	19	0.0082	0.1591	0.0332	0.0428	0.0112	0.0036	-0.0005	0.0482	0.0143	0.1202						0.0282	0.0056		0.0462
	20	0.0103	0.1392	0.0335	0.0581	0.0074	0.0270	0.0450	0.0026	0.0458	0.0674						0.0248	0.0077		-0.0034
	1	0.0094	0.1241	0.0260	0.0381	0.0026	-0.0072	0.0014	0.0026	0.0376	0.1202						0.1176	0.0885		0.0622
No	4	0.0134	0.0805	0.0124	0.0317	0.0070	0.0184	0.0422	0.0335	-0.0022	0.0216									
	5	-0.0042	0.1273	0.0251	0.0377	-0.0078	0.0003	0.0075	0.09129	0.0054	0.0021	0.0026								
	6	0.0540	0.0931	0.0471	0.0543	0.0452	0.0651	0.1090	0.0986	0.0242	0.0786	0.0171	0.0426							
	7	0.0246	0.0749	0.0172	0.0243	0.0330	0.0406	0.0801	0.0561	0.0214	0.0550	0.0046	0.0324	0.0141			0.0851	0.0400		0.0476
	8	0.0255	0.2594	0.0756	0.1060	0.0321	0.0273	-0.0009	0.0159	0.0940	0.0241	0.0886	0.0402	0.1689	0.1319					
	9	0.0849	0.0479	0.0401	0.0503	0.0781	0.0898	0.1425	0.1254	0.0467	0.1028	0.0292	0.0713	0.0145	0.0159	0.2086				
	12	0.0151	0.2198	0.0624	0.0869	0.0109	0.0112	-0.0060	0.0092	0.0540	0.0067	0.0517	0.0144	0.1240	0.1013	-0.0006	0.1628	0.0297		
	13	0.0154	0.2226	0.0644	0.0808	0.0226	0.0140	-0.0012	0.0740	0.0240	0.0390	0.0656	0.0285	0.1386	0.0955	-0.0008	0.1765	0.0019		
	15	0.0126	0.0849	0.0108	0.0296	0.0010	-0.0065	0.0111	0.0093	0.0270	-0.0095	0.0039	-0.0014	0.0626	0.0403	0.0463	0.0772	0.0188	0.0307	0.0271
	16	-0.0013	0.1128	0.0314	0.0251	0.0056	0.0230	0.0428	0.0429	0.0014	0.0348	0.0020	0.0074	0.0151	0.0062	0.0930	0.0329	0.0608	0.0713	0.0265

* Yes = bags of temephos were distributed to the householders; No = bags of temephos were not distributed to the householders. Significant differentiation (after application of Bonferroni's procedure at 5%) is underlined.