Control of pathogens and microbiota by innate lymphoid cells

Sascha Cording, Jasna Medvedovic, Emelyne Lecuyer, Tegest Aychek, Gerard Eberl

► To cite this version:

Sascha Cording, Jasna Medvedovic, Emelyne Lecuyer, Tegest Aychek, Gerard Eberl. Control of pathogens and microbiota by innate lymphoid cells. *Microbes and Infection*, 2018, 10.1016/j.micinf.2018.05.003 . pasteur-01824045

HAL Id: pasteur-01824045

<https://hal-pasteur.archives-ouvertes.fr/pasteur-01824045>

Submitted on 26 Jun 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution - NonCommercial - ShareAlike| 4.0 International License

Accepted Manuscript

Control of pathogens and microbiota by innate lymphoid cells

Sascha Cording, Jasna Medvedovic, Emelyne Lecuyer, Tegest Aychek, Gérard Eberl

PII: S1286-4579(18)30109-6

DOI: [10.1016/j.micinf.2018.05.003](https://doi.org/10.1016/j.micinf.2018.05.003)

Reference: MICINF 4587

To appear in: *Microbes and Infection*

Received Date: 5 April 2018

Accepted Date: 22 May 2018

Please cite this article as: S. Cording, J. Medvedovic, E. Lecuyer, T. Aychek, G. Eberl, Control of pathogens and microbiota by innate lymphoid cells, *Microbes and Infection* (2018), doi: 10.1016/j.micinf.2018.05.003.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Control of pathogens and microbiota by innate lymphoid cells

Sascha Cording^{1,2,3}, Jasna Medvedovic^{1,2}, Emelyne Lecuyer^{1,2}, Tegest Aychek^{1,2}, Gérard Eberl^{1,2}

¹ Institut Pasteur, Microenvironment & Immunity Unit, 75724 Paris, France

² INSERM U1224, 75724 Paris, France

³ Present address: INSERM, U989, Institut IMAGINE, and Université Paris Descartes-Sorbonne Paris Cité, Paris, France

Corresponding author: Gérard Eberl

E-mail address: gerard.eberl@pasteur.fr

Summary

Innate lymphoid cells (ILCs) are the innate counterpart of T cells. Upon infection or injury, ILCs react promptly to direct the developing immune response to the one most adapted to the threat facing the organism. Therefore, ILCs play an important role early in resistance to infection, but also to maintain homeostasis with the symbiotic microbiota following perturbations induced by diet and pathogens. Such roles of ILCs have been best characterized in the intestine and lung, mucosal sites that are exposed to the environment and are therefore colonized with diverse but specific types of microbes. Understanding the dialogue between pathogens, microbiota and ILCs may lead to new strategies to re-inforce immunity for prevention, vaccination and therapy.

Keywords: Innate lymphoid cells, microbiota, pathogens, homeostasis, mucosal immunity

I. Introduction

ILCs derive from a lymphoid cell precursor common with T cells, with whom they share phenotypes and functions [1,2]. However, T cells are antigen-specific, carry immunological memory, and are selected by their cognate antigen in lymph nodes and Peyer's patches before expansion. Once activated, T cells recirculate through the blood to reach organs where their effector functions are required, eventually committing apoptosis or remaining in tissues as resident memory T (T_{RM}) cells [3]. Nevertheless, the process of T cell selection, activation and recirculation requires several days, leaving the early phase of the immune response without T cell-mediated orchestration of the response best adapted to the type of threat faced by the individual. However, early orchestration is directed by ILCs, which perform similar regulatory functions and provide prompt effector responses to infection and injury. ILCs also play important roles early in life, when the adaptive immune system is not yet in place [1].

An individual faced with intracellular threats, such as tumors, viruses and particular bacteria, responds with type 1 immune responses characterized by the production of type I interferons, the inducer cytokine IL-12, the effector cytokine $IFN\gamma$, the release of cytotoxic oxygen radicals and proteins, and the elimination of the transformed or infected cells (Figure 1). Extracellular microbes, such as bacteria and fungi, elicit type 3 responses that are characterized by the release of the inducer cytokine IL-23 and the production of the effector cytokines IL-17 and IL-22, which lead to the reinforcement of mucosal barriers and the recruitment of polymorphonuclear neutrophilic phagocytes that target, ingest and destroy the microbes. In contrast, when faced with large parasites, such as worms, the individual develops type 2 responses that lead to the release of fluids and the production of mucus at mucosal surfaces, and the deposition of collagen to resist parasite penetration. The type 2 inducer cytokines are IL-25, IL-33 and TSLP, which lead to the production of the effector cytokines IL-4, IL-5 and IL-13. Myeloid cells and stromal cells produce the appropriate inducer cytokines in response to a specific type of threat, activating ILCs and T cells to differentiate into ILC1s and Th1, ILC2s and Th2, or ILC3s and Th17, which produce the effector cytokines characteristic of type 1, 2 and 3 immunity [1,4].

Importantly, these immune responses are not only engaged in response to pathogens and tissue injury, but also in response to the symbiotic microbiota [5]. Microbes are present at all mucosal surfaces, as well as within tissues. An estimated 10^{14} bacteria reside in the intestine, together with viruses, fungi, protists and occasional worms, while smaller yet significant collections of microbes are present in the oral cavity, genitourinary system and skin [6,7]. Furthermore, the list of viruses found within our organism is expanding [8], and

bacteria have been identified in lymphoid tissues [9,10] and placenta [11]. As a consequence, the immune system is constantly activated by the presence of these diverse symbiotic microbes. We have therefore argued that the primary function of the immune system is to maintain homeostasis of the host with its microbiota [5], a microbiota that is necessary for digestion, production of metabolites and defense. In contrast, if microbes are associated with injury of the host's cells and tissues, the immune reaction will lead to the elimination of the pathogens.

In this review, we discuss the role of ILCs in the control of the symbiotic or pathogenic microbiota (Figure 2 and Table 1). From recent data, it emerges that ILCs play unique roles in microbiota control, observations that may lead to a new understanding of how chronic inflammatory pathologies emerge when such control is lost, and create new opportunities for prevention and therapy of infectious diseases. We will not discuss the role of NK cells, the oldest member of the ILC family, which has been extensively studied since 40 years and best reviewed elsewhere [12].

II. ILCs in the control of intestinal and hepatic infections

ILC3s are the oldest and best characterized ILC family member (not considering NK cells). In the late 90's, non-B non-T lymphoid cells, termed lymphoid tissue inducer (LTi) cells, have been described that colonize developing secondary lymphoid tissues, the lymph nodes and the Peyer's patches, and shown to be required for their development [13,14]. LTi cells form clusters in fetal lymphoid tissues and the intestinal lamina propria [15], and activate stromal cells to initiate organogenesis [16]. Ten years later, it was shown that LTi cells are part of a larger family of ILC3s that depend on the hormone receptor and transcription factor ROR γ t [17], and express the type 3 cytokines IL-17, IL-22 and lymphotoxin (LT) [18-21]. Non-LTi ILC3s do not cluster and are not involved in the development of lymphoid tissues. Rather, these cells resemble more freely moving lymphoid effector cells, and together with LTi cells, play a critical role early in defense against enteric pathogens.

Using mouse models that lack all lymphoid cells or only B and T cells, it was reported that ILC3s are required early in the control of intestinal infection by *Proteobacteria*, such as *Citrobacter rodentium*, the murine homologue of human enteropathogenic *Escherichia coli*. The production of IL-22 is paramount for this protective function, and induces the expression of anti-microbial peptides (AMPs) by epithelial cells, such as Reg3 γ [18,21-23]. Also involved in increasing epithelial defense is membrane-bound LT, which bind its receptor

LT β R on epithelial cells and induces expression of the neutrophils chemoattractants CXCL1 and CXCL2 [24]. Furthermore, ILC3s are involved in the response to infection by *Salmonella enterica* through the production of IL-17 and IL22, as well as IFN γ [25]. The production of IL-17 by ILC3s and the recruitment of neutrophils are essential in the response to infection by *E.coli K1* and *Klebsiella pneumoniae* and to prevent sepsis [26]. The protective role of ILC3s against pathogens extends to parasites and viruses. Infection by *Toxoplasma gondii* leads to more severe inflammatory pathology in the absence of ILC3s, presumably because of a failure to contain the parasites early [27]. Furthermore, IL-22 production by ILC3s potentiates the activity of IFN λ in epithelial cells and thus increases resistance to rotavirus [28]. However, the pro-inflammatory activity of ILC3s may also lead to pathology through the production of IFN γ in the context of *S. enterica* infection [25], as well as intestinal fibrosis through the expression of IL-17 and IL-22 in the context of *S. typhimurium* infection [29].

ILC2s were first reported in the context of intestinal infection with the helminth *Nippostrongylus brasiliensis*, which evokes a vigorous expansion of “non-B non-T cells” expressing IL-4, IL-5 and IL-13 [30,31]. ILC2s are also involved in defense against *Heligmosomoides polygyrus* [32], *Strongyloides venezuelensis* [33], *Trichinella spiralis* [34] and *T. muris* [35], and are diminished in the blood of children infected with the blood dweller *Schistosoma haematobium* [36]. Both inducer cytokines IL-25 and IL-33 are expressed upon helminth infection and activate ILC2s, noting that ILC2s responding to IL-33 have been suggested to be precursors of ILC2s responding to IL-25 [37]. While IL-33 is expressed by different types of stromal cells, as well as by mast cells [32], the source of IL-25 was only recently identified, in the intestine, as Tuft epithelial cells [38-40]. Interestingly, Tuft cells appear to detect helminth infection through chemosensory signaling via the G-protein coupled receptor Trpm5 [38]. The activity of ILC2s is also promoted, during helminth infection, by the neuropeptide neuromedin U that is expressed by cholinergic neurons in the intestine [41,42].

ILC1s are distinguished from NK cells by their lack of expression of, and requirement for, the transcription factor Eomesodermin [2]. ILC1s provide early protection to the liver from mouse cytomegalovirus infection, producing IFN γ before NK cells are engaged [43], but at the same time prevent the recruitment of NK cells and CD8⁺ T cells that optimally fight liver infection with adenovirus [44]. Mouse hepatitis virus inoculated orally is also cleared from the intestine by ILC1s, which are activated by IL-15 released from infected stromal cells [45]. Furthermore, ILC1s and their production of IFN γ are engaged in the defense against

bacteria, such as *Clostridium difficile* [46]. Nevertheless, ILC1s and NK cells have largely overlapping function, even though ILC1s are defined as non-cytotoxic. As both cells types react promptly to activators such as IL-12, their relative role is mostly dependent on their tissue distribution before infection.

III. The ILC crosstalk with intestinal microbiota

The large intestinal microbiota has been best characterized, so far, at the level of its bacteriome and of its cross-talk between symbiotic bacteria and ILC3s. In contrast, the interaction between the virome and ILCs remains largely unexplored, and ILC2s are not known to cross-talk with microbes, even though some bacteria have been reported to induce the expansion of ILC2s at the steady state [47]. Therefore, this chapter will discuss mostly our knowledge on the cross-talk between bacterial symbionts and ILC3s.

In 2008, we reported that peptidoglycan released by proliferating Gram⁻ bacteria in the intestine induce the activation of LT α i cells clustered in so-called cryptopatches [15], which are found near the base of crypts in the small intestine [48]. LT α i cells in turn activate underlying stromal cells to release chemokines and recruit CCR6⁺ B cells to form isolated lymphoid follicles (ILFs). ILFs generate microbiota-specific IgA-producing B cells, in a T-cell independent way, and play an important role in intestinal homeostasis [15,49]. The colonizing microbiota is therefore involved in the development of the intestinal immune system. At the same time, it provides a negative feedback on the number and activity of ILC3s, which include LT α i cells, by inducing epithelial cells to produce IL-25 [47]. Of note, in both these phenomena, epithelial cells translate the recognition of microbes into signals that regulate ILCs.

Myeloid cells are nevertheless the major relay between microbiota and ILC3s through the production of the type 3 inducer cytokines IL-23 and IL-1 β . In the context of *C. rodentium* infection, CX₃CR1⁺ macrophages produce higher amounts of these cytokines than do conventional CD103⁺ DCs, and thus, more efficiently induce the expression by ILC3s of IL-22 [50] and GM-CSF [51]. In addition, CD11b⁺ DCs, which include CX₃CR1⁺ cells, are obligate sources of IL-23 for host survival during *C. rodentium* infection [52], and TNF α produced by CD11b⁺ DCs during *Helicobacter typhlonius* infection synergizes with IL-23 for the expression of IL-17 by ILC3s [53]. CD11b⁺ DCs are also activated by microbiota-derived ATP to produce IL-23 during the steady state [54]. Other members of the symbiotic microbiota induce the expression of IL-23 by myeloid cells and as a consequence, IL-22 by ILC3s, such as SFB [55] and *Lactobacilli* species [56]. ILC3s feedback positively on the

production of IL-23 by myeloid cells through their expression of membrane-bound LT and activation of LT β R on DCs [57]. Unexpectedly, intestinal glial cells sense microbiota through a Myd88-dependent pathway and produce neurotrophic factors that activate the Ret receptor on ILC3s and promote their expression of IL-22 [58].

At the steady state, ILC3s are the major source of IL-22 [47]. IL-22, as well as IL-17, induce the production of AMPs by epithelial cells [59], and therefore, are critical for the containment not only of pathogens, but also of the symbiotic microbiota. IL-22 also regulates the fucosylation of epithelial cells, a dietary carbohydrate for many symbionts, and thereby increases resistance to infection by *Salmonella typhimurium* [60] and *C. rodentium* [61]. Furthermore, membrane-bound (LT $\alpha_1\beta_2$) and soluble (LT α_3) LT expressed by ILC3s play an important role in the containment of the microbiota [62] through the activation of epithelial cells [24], as well as of DCs and T cells that lead to the production of anti-microbial IgA [63]. The containment of microbiota through the enhancement of epithelial defenses and adaptive immunity must nevertheless be controlled to avoid exaggerate responses to the symbiotic microbiota and consequent pathologic inflammation. To this end, ILCs express MHC class II molecules, as well as the machinery required to process proteins for presentation of peptides onto class II molecules, allowing ILCs to dampen the CD4⁺ T cell response to microbiota [64] and to SFB in particular [65]. Reciprocally, the presence of CD4⁺ T cells in the intestinal lamina propria decreases the expression of IL-22 by ILC3s [47,66], presumably through competition for survival and inducing cytokines such as IL-23.

Bacterial symbionts of the *Alcaligenes* genus have been identified that are present in Peyer's patches. ILC3s and IL-22 are critical for the containment of *Alcaligenes*, as the depletion of ILCs using an anti-CD90 antibody leads to the presence of the bacteria in spleen and liver [9]. The depletion experiments were performed in Rag-deficient mice, as well as in mice reconstituted with mature B and T cells, indicating that adaptive immunity is not necessary for the containment of *Alcaligenes*. It remains nevertheless possible that, in normal mice, adaptive immunity develops over time that contributes to the containment of these symbionts through the generation of antibodies. As a matter of fact, *Alcaligenes* promotes the generation of IgA, which contribute to the construction of its niche [10].

IV. Microbe-ILC interactions in the lungs

The impact of ILCs on microbes (and parasites) at mucosal sites other than the intestine has been best studied in the lungs. ILC2s are the dominant ILC subset in the lungs at the steady state, as a consequence of the production of IL-33 and TSLP by the alveolar epithelium [67].

The production of IL-5 and IL-13 by ILC2s promotes the recruitment of eosinophils, the differentiation of monocyte into M2 (or alternatively activated) macrophages, and the differentiation of T cells into Th2 cells [68-70]. ILC2s contribute to the resistance to lung parasites, such as *Strongyloides venezuelensis* [71], *Nippostrongylus brasiliensis* [72,73], *Litomosoides sigmodontis* [74], as well as to tissue repair [75]. However, type 2 immunity induced by ILC2s also leads to increased susceptibility to fungal [76,77] and bacterial infection [78], by mechanisms of immunological cross-regulation [4].

Type 2 immunity promoted by ILC2s plays an important role in the repair phase of immune responses, which occurs, for example, after influenza A virus infection through the production of the EGFR ligand amphiregulin [79]. This repair response is nevertheless inhibited by IFN γ that is induced by the virus infection [80]. Furthermore, the ILC2-initiated repair responses, induced by virus-induced tissue damage, can lead to exacerbated type 2 responses and, as a consequence, asthma. Such an association, and the mechanisms driving this association, have been reported in the context of infection with rhinovirus [81-83] and respiratory syncytial virus [84,85]. Similar pathological consequences are also induced by the fungi *Alternaria alternata* [86] and *Cryptococcus neoformans* [87].

ILC3s also play an important role in the lungs in response to bacterial and fungal infections. IL-22 produced by ILC3s is instrumental in lung immunity to *Streptococcus pneumoniae* [88] and *Pseudomonas aeruginosa* [89]. Interestingly, neonatal colonization of mice with intestinal bacteria induces the influx of IL-22-producing ILC3s into the lung, which confer life-saving resistance to early infection by *Streptococcus pneumoniae* [90].

V. Concluding remarks

We have discussed recent data showing the critical role played by ILC early in the immune responses to pathogens, as these cells can respond promptly to inducer cytokines provided by myeloid and epithelial cells upon infection and injury. Of note, T_{RM} cells, as well as particular subsets of invariant T $\alpha\beta$ or T $\gamma\delta$ cells, can play similar roles as their activation state alleviates the requirement for antigen-specific TCR activation. The prompt activation of ILCs is also important in response to changes in the symbiotic microbiota, in order to maintain homeostasis.

The current knowledge on the role of ILCs in defense and homeostasis has been inferred from mouse models and association studies in human. It is now possible to design strategies to harness the central role of ILCs in homeostasis, defense and immunoregulation for the prevention and therapy of infectious diseases, and of pathological consequences of

microbiota dysbiosis. It is also possible that novel vaccination strategies benefit from an early manipulation of ILCs, in order to force the response to specific antigens into the type most adapted to fight the pathogen. The manipulation of ILCs may involve the administration of inducer cytokines, or other molecules, yet to be discovered, that specifically activate subsets of ILCs. Another possibility is the isolation of ILC precursors from the blood of patients, and their expansion *in vitro* into the desired subset before re-administration and induction of type 1, 2 or 3 immune responses. We are still in the early days of ILC biology, but also at the exciting transition phase when clinical applications become possible.

240 **Table 1**

ILC type	Role	Microbe type	Microbe species	Organ	References
ILC1	+	Bacteria	<i>Clostridium difficile</i>	Intestine	[46]
	+	Virus	<i>Adenovirus</i>	Liver	[44]
	+		<i>Mouse hepatitis virus</i>	Liver	[45]
	+		<i>Cytomegalovirus</i>	Liver	[43]
ILC2	–	Bacteria	<i>Streptococcus pneumoniae</i>	Lung	[78]
	–	Virus	<i>Rhinovirus</i>	Lung	[81-83]
	–		<i>Respiratory syncytial virus</i>	Lung	[84,85]
	+/-		<i>Influenza</i>	Lung	[79,80]
	+/-	Fungi	<i>Cryptococcus neoformans</i>	Lung	[76,77,87]
	–		<i>Alternaria alternata</i>	Lung	[86]
	+	Parasites	<i>Trichinella spiralis</i>	Intestine	[34]
	+		<i>Trichuris muris</i>	Intestine	[35]
	+		<i>Heligmosomoides polygyrus</i>	Intestine	[32]
	+		<i>Nippostrongylus brasiliensis</i>	Intestine/lung	[30,31,72,73]
	+		<i>Strongyloides venezuelensis</i>	Intestine/lung	[33,71]
	+		<i>Litomosoides sigmodontis</i>	Lung	[74]
	+		<i>Schistosoma haematobium</i>	Blood	[36]
ILC3	+	Bacteria	<i>Commensal microbiota</i>	Intestine	[15,24,47,49,54,62-64]
	+		<i>Citrobacter rodentium</i>	Intestine	[18,21-23,50,52,61]
	+		<i>Escherichia coli K1</i>	Intestine	[26]
	+		<i>SFB</i>	Intestine	[55,65]
	+		<i>Lactobacilli</i>	Intestine	[56]
	+		<i>Alcaligenes</i>	Intestine	[9,10]
	+/-		<i>Salmonella enterica</i>	Intestine	[25]
	+/-		<i>Salmonella typhimurium</i>	Intestine	[29,60]
	+/-		<i>Helicobacter typhlonius</i>	Intestine	[53]
	+		<i>Streptococcus pneumoniae</i>	Lung	[88,90]
	+		<i>Pseudomonas aeruginosa</i>	Lung	[89]
	+		<i>Klebsiella pneumoniae</i>	Blood	[26]
	+	Virus	<i>Rotavirus</i>	Intestine	[28]
	+	Fungi	<i>Candida albicans</i>	Lung	[89]
	+	Parasites	<i>Toxoplasma gondii</i>	Intestine	[27]

241

242

243 **Conflict of interests**

244 We declare no conflicts of interests

References

- [1] Eberl G, Colonna M, Di Santo JP, McKenzie AN. Innate lymphoid cells: a new paradigm in immunology. *Science* 2015;348:aaa6566.
- [2] Serafini N, Voshenrich CA, Di Santo JP. Transcriptional regulation of innate lymphoid cell fate. *Nat Rev Immunol* 2015;15:415-28.
- [3] Mueller SN, Mackay LK. Tissue-resident memory T cells: local specialists in immune defence. *Nat Rev Immunol* 2016;16:79-89.
- [4] Eberl G. Immunity by equilibrium. *Nat Rev Immunol* 2016;16:524-32.
- [5] Eberl G. A new vision of immunity: homeostasis of the superorganism. *Mucosal Immunol* 2010;3:450-60.
- [6] Proctor DM, Relman DA. The landscape ecology and microbiota of the human nose, mouth, and throat. *Cell Host Microbe* 2017;21:421-32.
- [7] Consortium HMP. Structure, function and diversity of the healthy human microbiome. *Nature* 2012;486:207-14.
- [8] Virgin HW, Wherry EJ, Ahmed R. Redefining chronic viral infection. *Cell* 2009;138:30-50.
- [9] Sonnenberg GF, Monticelli LA, Alenghat T, Fung TC, Hutnick NA, Kunisawa J et al. Innate lymphoid cells promote anatomical containment of lymphoid-resident commensal bacteria. *Science* 2012;336:1321-5.
- [10] Obata T, Goto Y, Kunisawa J, Sato S, Sakamoto M, Setoyama H et al. Indigenous opportunistic bacteria inhabit mammalian gut-associated lymphoid tissues and share a mucosal antibody-mediated symbiosis. *Proc Natl Acad Sci U S A* 2010;107:7419-24.
- [11] Aagaard K, Ma J, Antony KM, Ganu R, Petrosino J, Versalovic J. The placenta harbors a unique microbiome. *Sci Transl Med* 2014;6:237ra65.
- [12] Trinchieri G. Natural killer cells wear different hats: effector cells of innate resistance and regulatory cells of adaptive immunity and of hematopoiesis. *Semin Immunol* 1995;7:83-8.
- [13] Mebius RE, Rennert P, Weissman IL. Developing lymph nodes collect CD4⁺CD3⁻LTb⁺ cells that can differentiate to APC, NK cells, and follicular cells but not T or B cells. *Immunity* 1997;7:493-504.
- [14] Adachi S, Yoshida H, Kataoka H, Nishikawa S. Three distinctive steps in Peyer's patch formation of murine embryo. *Int Immunol* 1997;9:507-14.
- [15] Bouskra D, Brezillon C, Berard M, Werts C, Varona R, Boneca IG et al. Lymphoid tissue genesis induced by commensals through NOD1 regulates intestinal homeostasis. *Nature* 2008;456:507-10.
- [16] Honda K, Nakano H, Yoshida H, Nishikawa S, Rennert P, Ikuta K et al. Molecular basis for hematopoietic/mesenchymal interaction during initiation of Peyer's patch organogenesis. *J Exp Med* 2001;193:621-30.
- [17] Eberl G, Marmon S, Sunshine MJ, Rennert PD, Choi Y, Littman DR. An essential function for the nuclear receptor ROR γ t in the generation of fetal lymphoid tissue inducer cells. *Nat Immunol* 2004;5:64-73.
- [18] Satoh-Takayama N, Voshenrich CA, Lesjean-Pottier S, Sawa S, Lochner M, Rattis F et al. Microbial flora drives interleukin 22 production in intestinal NKp46⁺ cells that provide innate mucosal immune defense. *Immunity* 2008;29:958-70.
- [19] Cella M, Fuchs A, Vermi W, Facchetti F, Otero K, Lennerz JK et al. A human natural killer cell subset provides an innate source of IL-22 for mucosal immunity. *Nature* 2009;457:722-5.
- [20] Luci C, Reynders A, Ivanov, II, Cognet C, Chiche L, Chasson L et al. Influence of the transcription factor ROR γ on the development of NKp46⁺ cell populations in gut and skin. *Nat Immunol* 2009;10:75-82.

- [21] Sanos SL, Bui VL, Mortha A, Oberle K, Heners C, Johner C et al. RORgammat and commensal microflora are required for the differentiation of mucosal interleukin 22-producing NKp46+ cells. *Nat Immunol* 2009;10:83-91.
- [22] Sonnenberg GF, Monticelli LA, Elloso MM, Fouser LA, Artis D. CD4(+) Lymphoid tissue-inducer cells promote innate immunity in the gut. *Immunity* 2011;34:122-34.
- [23] Guo X, Qiu J, Tu T, Yang X, Deng L, Anders RA et al. Induction of innate lymphoid cell-derived interleukin-22 by the transcription factor STAT3 mediates protection against intestinal infection. *Immunity* 2014;40:25-39.
- [24] Wang Y, Koroleva EP, Kruglov AA, Kuprash DV, Nedospasov SA, Fu YX et al. Lymphotoxin beta receptor signaling in intestinal epithelial cells orchestrates innate immune responses against mucosal bacterial infection. *Immunity* 2010;32:403-13.
- [25] Klose CS, Kiss EA, Schwierzeck V, Ebert K, Hoyer T, d'Hargues Y et al. A T-bet gradient controls the fate and function of CCR6-RORgammat+ innate lymphoid cells. *Nature* 2013;494:261-5.
- [26] Deshmukh HS, Liu Y, Menkiti OR, Mei J, Dai N, O'Leary CE et al. The microbiota regulates neutrophil homeostasis and host resistance to *Escherichia coli* K1 sepsis in neonatal mice. *Nat Med* 2014;20:524-30.
- [27] Wagage S, Harms Pritchard G, Dawson L, Buza EL, Sonnenberg GF, Hunter CA. The group 3 innate lymphoid cell defect in aryl hydrocarbon receptor deficient mice is associated with T cell hyperactivation during intestinal infection. *PLoS One* 2015;10:e0128335.
- [28] Hernandez PP, Mahlakoiv T, Yang I, Schwierzeck V, Nguyen N, Guendel F et al. Interferon-lambda and interleukin 22 act synergistically for the induction of interferon-stimulated genes and control of rotavirus infection. *Nat Immunol* 2015;16:698-707.
- [29] Lo BC, Gold MJ, Hughes MR, Antignano F, Valdez Y, Zaph C et al. The orphan nuclear receptor ROR alpha and group 3 innate lymphoid cells drive fibrosis in a mouse model of Crohn's disease. *Sci Immunol* 2016;1:pii: eaaf8864.
- [30] Moro K, Yamada T, Tanabe M, Takeuchi T, Ikawa T, Kawamoto H et al. Innate production of T(H)2 cytokines by adipose tissue-associated c-Kit(+)Sca-1(+) lymphoid cells. *Nature* 2009;463:540-4.
- [31] Fallon PG, Ballantyne SJ, Mangan NE, Barlow JL, Dasvarma A, Hewett DR et al. Identification of an interleukin (IL)-25-dependent cell population that provides IL-4, IL-5, and IL-13 at the onset of helminth expulsion. *J Exp Med* 2006;203:1105-16.
- [32] Shimokawa C, Kanaya T, Hachisuka M, Ishiwata K, Hisaeda H, Kurashima Y et al. Mast cells are crucial for induction of group 2 innate lymphoid cells and clearance of helminth infections. *Immunity* 2017;46:863-74 e4.
- [33] Mukai K, Karasuyama H, Kabashima K, Kubo M, Galli SJ. Differences in the Importance of Mast Cells, Basophils, IgE, and IgG versus That of CD4(+) T Cells and ILC2 Cells in Primary and Secondary Immunity to *Strongyloides venezuelensis*. *Infect Immun* 2017;85:
- [34] Angkasekwinai P, Sodthawon W, Jeerawattanawart S, Hansakon A, Pattanapanyasat K, Wang YH. ILC2s activated by IL-25 promote antigen-specific Th2 and Th9 functions that contribute to the control of *Trichinella spiralis* infection. *PLoS One* 2017;12:e0184684.
- [35] Spencer SP, Wilhelm C, Yang Q, Hall JA, Bouladoux N, Boyd A et al. Adaptation of innate lymphoid cells to a micronutrient deficiency promotes type 2 barrier immunity. *Science* 2014;343:432-7.
- [36] Nausch N, Appleby LJ, Sparks AM, Midzi N, Mduluza T, Mutapi F. Group 2 innate lymphoid cell proportions are diminished in young helminth infected children and restored by curative anti-helminthic treatment. *PLoS Negl Trop Dis* 2015;9:e0003627.

- [37] Huang Y, Guo L, Qiu J, Chen X, Hu-Li J, Siebenlist U et al. IL-25-responsive, lineage-negative KLRG1(hi) cells are multipotential 'inflammatory' type 2 innate lymphoid cells. *Nat Immunol* 2015;16:161-9.
- [38] Howitt MR, Lavoie S, Michaud M, Blum AM, Tran SV, Weinstock JV et al. Tuft cells, taste-chemosensory cells, orchestrate parasite type 2 immunity in the gut. *Science* 2016;351:1329-33.
- [39] Gerbe F, Sidot E, Smyth DJ, Ohmoto M, Matsumoto I, Dardalhon V et al. Intestinal epithelial tuft cells initiate type 2 mucosal immunity to helminth parasites. *Nature* 2016;529:226-30.
- [40] von Moltke J, Ji M, Liang HE, Locksley RM. Tuft-cell-derived IL-25 regulates an intestinal ILC2-epithelial response circuit. *Nature* 2016;529:221-5.
- [41] Klose CSN, Mahlakoiv T, Moeller JB, Rankin LC, Flamar AL, Kabata H et al. The neuropeptide neuromedin U stimulates innate lymphoid cells and type 2 inflammation. *Nature* 2017;549:282-86.
- [42] Cardoso V, Chesne J, Ribeiro H, Garcia-Cassani B, Carvalho T, Bouchery T et al. Neuronal regulation of type 2 innate lymphoid cells via neuromedin U. *Nature* 2017;549:277-81.
- [43] Weizman OE, Adams NM, Schuster IS, Krishna C, Pritykin Y, Lau C et al. ILC1 confer early host protection at initial sites of viral infection. *Cell* 2017;171:795-808 e12.
- [44] Krueger PD, Narayanan S, Surette FA, Brown MG, Sung SJ, Hahn YS. Murine liver-resident group 1 innate lymphoid cells regulate optimal priming of anti-viral CD8⁺ T cells. *J Leukoc Biol* 2017;101:329-38.
- [45] Gil-Cruz C, Perez-Shibayama C, Onder L, Chai Q, Cupovic J, Cheng HW et al. Fibroblastic reticular cells regulate intestinal inflammation via IL-15-mediated control of group 1 ILCs. *Nat Immunol* 2016;17:1388-96.
- [46] Abt MC, Lewis BB, Caballero S, Xiong H, Carter RA, Susac B et al. Innate immune defenses mediated by two ILC subsets are critical for protection against acute *Clostridium difficile* infection. *Cell Host Microbe* 2015;18:27-37.
- [47] Sawa S, Lochner M, Satoh-Takayama N, Dulauroy S, Berard M, Kleinschek M et al. ROR γ t⁺ innate lymphoid cells regulate intestinal homeostasis by integrating negative signals from the symbiotic microbiota. *Nat Immunol* 2011;12:320-26.
- [48] Kanamori Y, Ishimaru K, Nanno M, Maki K, Ikuta K, Nariuchi H et al. Identification of novel lymphoid tissues in murine intestinal mucosa where clusters of c-kit⁺ IL-7R⁺ Thy1⁺ lympho-hemopoietic progenitors develop. *J Exp Med* 1996;184:1449-59.
- [49] Tsuji M, Suzuki K, Kitamura H, Maruya M, Kinoshita K, Ivanov, II et al. Requirement for lymphoid tissue-inducer cells in isolated follicle formation and T cell-independent immunoglobulin a generation in the gut. *Immunity* 2008;29:261-71.
- [50] Longman RS, Diehl GE, Victorio DA, Huh JR, Galan C, Miraldi ER et al. CX(3)CR1(+) mononuclear phagocytes support colitis-associated innate lymphoid cell production of IL-22. *J Exp Med* 2014;211:1571-83.
- [51] Mortha A, Chudnovskiy A, Hashimoto D, Bogunovic M, Spencer SP, Belkaid Y et al. Microbiota-dependent crosstalk between macrophages and ILC3 promotes intestinal homeostasis. *Science* 2014;343:1249288.
- [52] Satpathy AT, Briseno CG, Lee JS, Ng D, Manieri NA, Kc W et al. Notch2-dependent classical dendritic cells orchestrate intestinal immunity to attaching-and-effacing bacterial pathogens. *Nat Immunol* 2013;14:937-48.
- [53] Powell N, Walker AW, Stolarczyk E, Canavan JB, Gokmen MR, Marks E et al. The transcription factor T-bet regulates intestinal inflammation mediated by interleukin-7 receptor⁺ innate lymphoid cells. *Immunity* 2012;37:674-84.

- [54] Atarashi K, Nishimura J, Shima T, Umesaki Y, Yamamoto M, Onoue M et al. ATP drives lamina propria T(H)17 cell differentiation. *Nature* 2008;455:808-12.
- [55] Sano T, Huang W, Hall JA, Yang Y, Chen A, Gavzy SJ et al. An IL-23R/IL-22 circuit regulates epithelial serum amyloid A to promote local effector Th17 responses. *Cell* 2015;163:381-93.
- [56] Nakamoto N, Amiya T, Aoki R, Taniki N, Koda Y, Miyamoto K et al. Commensal *lactobacillus* controls immune tolerance during acute liver injury in mice. *Cell Rep* 2017;21:1215-26.
- [57] Tumanov AV, Koroleva EP, Guo X, Wang Y, Kruglov A, Nedospasov S et al. Lymphotoxin controls the IL-22 protection pathway in gut innate lymphoid cells during mucosal pathogen challenge. *Cell Host Microbe* 2011;10:44-53.
- [58] Ibiza S, Garcia-Cassani B, Ribeiro H, Carvalho T, Almeida L, Marques R et al. Glial-cell-derived neuroregulators control type 3 innate lymphoid cells and gut defence. *Nature* 2016;535:440-43.
- [59] Liang SC, Tan XY, Luxenberg DP, Karim R, Dunussi-Joannopoulos K, Collins M et al. Interleukin (IL)-22 and IL-17 are coexpressed by Th17 cells and cooperatively enhance expression of antimicrobial peptides. *J Exp Med* 2006;203:2271-9.
- [60] Goto Y, Obata T, Kunisawa J, Sato S, Ivanov, II, Lamichhane A et al. Innate lymphoid cells regulate intestinal epithelial cell glycosylation. *Science* 2014;345:1254009.
- [61] Pickard JM, Maurice CF, Kinnebrew MA, Abt MC, Schenten D, Golovkina TV et al. Rapid fucosylation of intestinal epithelium sustains host-commensal symbiosis in sickness. *Nature* 2014;514:638-41.
- [62] Zhang Y, Kim TJ, Wroblewska JA, Tesic V, Upadhyay V, Weichselbaum RR et al. Type 3 innate lymphoid cell-derived lymphotoxin prevents microbiota-dependent inflammation. *Cell Mol Immunol* 2017;
- [63] Kruglov AA, Grivennikov SI, Kuprash DV, Winsauer C, Prepens S, Seleznik GM et al. Nonredundant function of soluble LTalpha3 produced by innate lymphoid cells in intestinal homeostasis. *Science* 2013;342:1243-6.
- [64] Hepworth MR, Monticelli LA, Fung TC, Ziegler CG, Grunberg S, Sinha R et al. Innate lymphoid cells regulate CD4⁺ T-cell responses to intestinal commensal bacteria. *Nature* 2013;498:113-7.
- [65] Goto Y, Panea C, Nakato G, Cebula A, Lee C, Diez MG et al. Segmented filamentous bacteria antigens presented by intestinal dendritic cells drive mucosal Th17 cell differentiation. *Immunity* 2014;40:594-607.
- [66] Korn LL, Thomas HL, Hubbeling HG, Spencer SP, Sinha R, Simkins HM et al. Conventional CD4⁺ T cells regulate IL-22-producing intestinal innate lymphoid cells. *Mucosal Immunol* 2014;7:1045-57.
- [67] McKenzie AN, Spits H, Eberl G. Innate lymphoid cells in inflammation and immunity. *Immunity* 2014;41:366-74.
- [68] Molofsky AB, Nussbaum JC, Liang HE, Van Dyken SJ, Cheng LE, Mohapatra A et al. Innate lymphoid type 2 cells sustain visceral adipose tissue eosinophils and alternatively activated macrophages. *J Exp Med* 2013;210:535-49.
- [69] Nussbaum JC, Van Dyken SJ, von Moltke J, Cheng LE, Mohapatra A, Molofsky AB et al. Type 2 innate lymphoid cells control eosinophil homeostasis. *Nature* 2013;502:245-8.
- [70] Halim TY, Steer CA, Matha L, Gold MJ, Martinez-Gonzalez I, McNagny KM et al. Group 2 innate lymphoid cells are critical for the initiation of adaptive T helper 2 cell-mediated allergic lung inflammation. *Immunity* 2014;40:425-35.
- [71] Yasuda K, Matsumoto M, Nakanishi K. Importance of both innate immunity and acquired immunity for rapid expulsion of *S. venezuelensis*. *Front Immunol* 2014;5:118.

- [72] Bando JK, Nussbaum JC, Liang HE, Locksley RM. Type 2 innate lymphoid cells constitutively express arginase-I in the naive and inflamed lung. *J Leukoc Biol* 2013;94:877-84.
- [73] Bouchery T, Kyle R, Camberis M, Shepherd A, Filbey K, Smith A et al. ILC2s and T cells cooperate to ensure maintenance of M2 macrophages for lung immunity against hookworms. *Nature communications* 2015;6:6970.
- [74] Boyd A, Killoran K, Mitre E, Nutman TB. Pleural cavity type 2 innate lymphoid cells precede Th2 expansion in murine *Litomosoides sigmodontis* infection. *Exp Parasitol* 2015;159:118-26.
- [75] Turner JE, Morrison PJ, Wilhelm C, Wilson M, Ahlfors H, Renauld JC et al. IL-9-mediated survival of type 2 innate lymphoid cells promotes damage control in helminth-induced lung inflammation. *J Exp Med* 2013;210:2951-65.
- [76] Flaczyk A, Duerr CU, Shourian M, Lafferty EI, Fritz JH, Qureshi ST. IL-33 signaling regulates innate and adaptive immunity to *Cryptococcus neoformans*. *J Immunol* 2013;191:2503-13.
- [77] Piehler D, Grahnert A, Eschke M, Richter T, Kohler G, Stenzel W et al. T1/ST2 promotes T helper 2 cell activation and polyfunctionality in bronchopulmonary mycosis. *Mucosal Immunol* 2013;6:405-14.
- [78] Saluzzo S, Gorki AD, Rana BMJ, Martins R, Scanlon S, Starkl P et al. First-breath-induced type 2 pathways shape the lung immune environment. *Cell Rep* 2017;18:1893-905.
- [79] Monticelli LA, Sonnenberg GF, Abt MC, Alenghat T, Ziegler CG, Doering TA et al. Innate lymphoid cells promote lung-tissue homeostasis after infection with influenza virus. *Nat Immunol* 2011;12:1045-54.
- [80] Califano D, Furuya Y, Roberts S, Avram D, McKenzie ANJ, Metzger DW. IFN-gamma increases susceptibility to influenza A infection through suppression of group II innate lymphoid cells. *Mucosal Immunol* 2018;11:209-19.
- [81] Hong JY, Bentley JK, Chung Y, Lei J, Steenrod JM, Chen Q et al. Neonatal rhinovirus induces mucous metaplasia and airways hyperresponsiveness through IL-25 and type 2 innate lymphoid cells. *J Allergy Clin Immunol* 2014;134:429-39.
- [82] Jackson DJ, Makrinioti H, Rana BM, Shamji BW, Trujillo-Torralbo MB, Footitt J et al. IL-33-dependent type 2 inflammation during rhinovirus-induced asthma exacerbations in vivo. *Am J Respir Crit Care Med* 2014;190:1373-82.
- [83] Han M, Rajput C, Hong JY, Lei J, Hinde JL, Wu Q et al. The innate cytokines IL-25, IL-33, and TSLP cooperate in the induction of type 2 innate lymphoid cell expansion and mucous metaplasia in rhinovirus-infected immature mice. *J Immunol* 2017;199:1308-18.
- [84] Saravia J, You D, Shrestha B, Jaligama S, Siefker D, Lee GI et al. Respiratory Syncytial Virus Disease Is Mediated by Age-Variable IL-33. *PLoS Pathog* 2015;11:e1005217.
- [85] Stier MT, Bloodworth MH, Toki S, Newcomb DC, Goleniewska K, Boyd KL et al. Respiratory syncytial virus infection activates IL-13-producing group 2 innate lymphoid cells through thymic stromal lymphopoietin. *J Allergy Clin Immunol* 2016;138:814-24 e11.
- [86] Stier MT, Zhang J, Goleniewska K, Cephus JY, Rusznak M, Wu L et al. IL-33 promotes the egress of group 2 innate lymphoid cells from the bone marrow. *J Exp Med* 2018;215:263-81.
- [87] Piehler D, Eschke M, Schulze B, Protschka M, Muller U, Grahnert A et al. The IL-33 receptor (ST2) regulates early IL-13 production in fungus-induced allergic airway inflammation. *Mucosal Immunol* 2016;9:937-49.

- [88] Van Maele L, Carnoy C, Cayet D, Ivanov S, Porte R, Deruy E et al. Activation of type 3 innate lymphoid cells and interleukin 22 secretion in the lungs during *Streptococcus pneumoniae* infection. J Infect Dis 2014;210:493-503.
- [89] Mear JB, Gosset P, Kipnis E, Faure E, Dessein R, Jawhara S et al. Candida albicans airway exposure primes the lung innate immune response against *Pseudomonas aeruginosa* infection through innate lymphoid cell recruitment and interleukin-22-associated mucosal response. Infect Immun 2014;82:306-15.
- [90] Gray J, Oehrle K, Worthen G, Alenghat T, Whitsett J, Deshmukh H. Intestinal commensal bacteria mediate lung mucosal immunity and promote resistance of newborn mice to infection. Sci Transl Med 2017;9: pii: eaaf9412.

Legends to figures

Figure 1. The activation of innate lymphoid cells. The three types of ILCs are promptly activated by distinct types of threats. Such threats are detected by myeloid and non-hematopoietic cells, which express inducer cytokines (in boxes). In reaction to inducer cytokines, ILCs express effector cytokines that both activate defense mechanisms and regulate immunity.

Figure 2. The control of microbiota and pathogens by ILCs. The response of ILCs to symbiotic microbes and pathogens have been extensively described in the intestine and lung, to include bacteria, viruses, fungi and parasites, as well as viruses in the liver. Given their prompt activation, ILCs play a critical role early in the effector response to perturbations in the microbiota and to pathogens, and in the regulation of adaptive immunity. In some cases, however, the reactivity of ILCs contributes to pathology.

Figure 1

Intracellular threats
(Tumors, viruses and intracellular bacteria)

Large parasites
(Helminths)

Extracellular microbes
(Bacteria, fungi)

Figure 2