



HAL
open science

Immune Responses to Dengue and Zika Viruses-Guidance for T Cell Vaccine Development

Claude Roth, Félix Delgado, Etienne Simon-Loriere, Anavaj Sakuntabhai

► **To cite this version:**

Claude Roth, Félix Delgado, Etienne Simon-Loriere, Anavaj Sakuntabhai. Immune Responses to Dengue and Zika Viruses-Guidance for T Cell Vaccine Development. *International Journal of Environmental Research and Public Health*, 2018, 15 (2), pp.385. 10.3390/ijerph15020385 . pasteur-02003866

HAL Id: pasteur-02003866

<https://hal-pasteur.archives-ouvertes.fr/pasteur-02003866>

Submitted on 25 Feb 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution| 4.0 International License



Review

Immune Responses to Dengue and Zika Viruses—Guidance for T Cell Vaccine Development

Claude Roth ^{1,2,*}, Félix G. Delgado ^{1,2,3} , Etienne Simon-Lorière ^{1,2}
and Anavaj Sakuntabhai ^{1,2}

¹ Functional Genetics of Infectious Diseases Unit, Institut Pasteur, 75015 Paris, France; fdelgadot@unbosque.edu.co (F.G.D.); etienne.simon-lorriere@pasteur.fr (E.S.-L.); anavaj.sakuntabhai@pasteur.fr (A.S.)

² CNRS UMR 2000–Génomique Évolutive, Modélisation et Santé, Institut Pasteur, 75015 Paris, France

³ Virology Group, Universidad El Bosque, Bogotá D.C. 110121, Colombia

* Correspondence: claude.roth@pasteur.fr; Tel.: +33-1-45-68-89-62

Received: 31 January 2018; Accepted: 19 February 2018; Published: 23 February 2018

Abstract: Despite numerous efforts to identify the molecular and cellular effectors of the adaptive immunity that induce a long-lasting immunity against dengue or Zika virus infection, the specific mechanisms underlying such protective immunity remain largely unknown. One of the major challenges lies in the high level of dengue virus (DENV) seroprevalence in areas where Zika virus (ZIKV) is circulating. In the context of such a pre-existing DENV immunity that can exacerbate ZIKV infection and disease, and given the lack of appropriate treatment for ZIKV infection, there is an urgent need to develop an efficient vaccine against DENV and ZIKV. Notably, whereas several ZIKV vaccine candidates are currently in clinical trials, all these vaccine candidates have been designed to induce neutralizing antibodies as the primary mechanism of immune protection. Given the difficulty to elicit simultaneously high levels of neutralizing antibodies against the different DENV serotypes, and the potential impact of pre-existing subneutralizing antibodies induced upon DENV infection or vaccination on ZIKV infection and disease, additional or alternative strategies to enhance vaccine efficacy, through T cell immunity, are now being considered. In this review, we summarize recent discoveries about cross-reactive B and T cell responses against DENV and ZIKV and propose guidelines for the development of safe and efficient T cell vaccines targeting both viruses.

Keywords: Zika virus; dengue virus; T cell epitopes; vaccination

1. History

Zika virus (ZIKV) is a Flavivirus transmitted by *Aedes* species mosquitoes. It is a single positive-stranded RNA virus closely related to yellow fever virus (YFV), dengue virus (DENV) and West Nile virus (WNV) [1]. First isolated in Uganda in 1947 [2], it remained confined to several regions in Africa and Asia from that time until the early 2000s. In 2007, however, it caused an explosive outbreak for the first time outside of Africa and Asia, on Yap Island, Federated States of Micronesia [3,4], followed by subsequent outbreaks with higher numbers of cases in 2013–2014 in French Polynesia and other South Pacific Islands and more recently in the Americas [5–9]. Although initially believed to only cause mild disease, the 2013–2014 and 2015 outbreaks in French Polynesia and Brazil clearly revealed that ZIKV causes neurological complications, such as Guillain-Barré syndrome in adults and microcephaly in infants born to ZIKV-infected women [10–13]. Phylogenetic studies indicated the presence of two lineages of ZIKV, the African and Asian lineages, the latter being responsible for the recent major outbreaks in French Polynesia and South America [14,15]. Notably, it was suggested that the enhanced infectivity of the Asian lineage of ZIKV was due to a spontaneous mutation in the gene coding for Non-Structural Protein 1 (NS1) leading to its higher secretion in the serum and infectivity to

mosquitoes [16], which could explain its recent re-emergence in the Americas [14,15] despite its relative absence in South East Asia. More strikingly, several amino acid substitutions in the proteome or more specifically in the precursor membrane (prM) protein with possible functional implications for ZIKV biology and pathogenesis have been identified from ZIKV outbreak strains in South America [17,18].

In addition to the high infectivity of the Asian lineage in the Americas, one of the most important concerns today is related to the high level of DENV seroprevalence in areas where ZIKV is circulating [19]. This is particularly important given the structural similarities between ZIKV and DENV [20–22], and the existence of cross-reactive immune responses associated with disease pathogenesis [23–27]. Nevertheless, while it is now well established that a secondary infection with a heterologous DENV serotype represents a risk factor for the development of severe dengue disease, because of serotype cross-reactive or sub-neutralizing antibodies which can mediate antibody dependent enhancement (ADE) [28], it remains to be determined whether a previous DENV infection can also increase the risk of developing a more severe ZIKV disease in humans, as suggested by studies in mice [26]. Likewise, while ZIKV-immune plasma can enhance DENV infection in immune-deficient mice [24], the role of ZIKV immunity in protection or enhancement of dengue disease in humans is still unknown.

In this review, we address the most recent findings regarding the adaptive immune response against ZIKV, focusing on the effect of DENV pre-existing immunity on ZIKV infection, the underlying idea being to identify immunological parameters predictive of increased susceptibility or protection against ZIKV infection and disease. In this respect, we will review the current state of knowledge on the impact of anti-DENV antibodies on ZIKV infection and disease, and then summarize the recent data on the potential role of T cells in DENV and ZIKV immunity, with the aim to promote a long lasting immune protection against these two viruses.

2. Antibody Cross-Reactivity between Zika and Dengue Viruses

The high level of cross-reactivity among flaviviruses, especially DENV and ZIKV which share 54–59% sequence identity in the E protein [20,29,30], and their co-circulation in the same endemic regions have complicated serological approaches to discriminate between these two viral infections. In most cases, reverse transcription-polymerase chain reaction (RT-PCR)-based assays within a week post-infection, in combination with serological binding assays to recombinant proteins and functional neutralization assays *in vitro*, either by Plaque Reduction Neutralization Test (PRNT) or Flow-Cytometry-Based Neutralization Assay have been developed to distinguish between ZIKV and DENV infections [23,24,31–34].

Upon primary DENV infection, different types of antibodies have been found in polyclonal human sera: serotype-specific antibodies with strong neutralizing activity against the homologous DENV serotype, and cross-reactive antibodies with weak neutralizing activity and strong enhancing potential against heterologous DENV serotypes [35,36]. In secondary DENV infections, both type-specific and cross-reactive neutralizing antibodies, as well as cross-reactive enhancing antibodies to the other DENV serotypes, are elicited [37]. Following multiple DENV infections, the polyclonal response has been shown to contain mainly cross-reactive antibodies that recognize the different DENV serotypes, these antibodies being potentially associated with a more durable cross-protective immunity [37,38].

Upon ZIKV infection, it has been shown that the quality and type of antibody response depends on previous infection with DENV, with a more restricted specificity against ZIKV in DENV naïve individuals [24,25]. Using a panel of monoclonal antibodies (mAbs) derived from ZIKV-infected individuals with or without previous DENV immunity, it was shown notably that the antibody response against ZIKV includes envelope domain III (EDIII)-specific antibodies with strong neutralizing activity against ZIKV, as well as envelope domains I and II (EDI/II)-specific antibodies with high degree of cross-reactivity against the E protein of all four DENV serotypes and low neutralizing potential against ZIKV [24]. Thus, just like how EDI/II-specific and cross-reactive antibodies induced after DENV infection can enhance DENV and ZIKV infection [23,39–43], EDI/II

cross-reactive antibodies raised by ZIKV infection can also potently enhance DENV and ZIKV infection [24,27]. Interestingly, not all cross-reactive antibodies are enhancing, as shown in a recent study where memory B cell clones derived from different ZIKV-infected individuals could produce antibodies that neutralize both ZIKV and DENV serotype 1 (DENV1) [44]. In this case, cross-reactivity was shown to result from a previous DENV1 infection and the clonal expansion of memory B cells specific of an EDIII epitope of ZIKV and DENV1 and neutralize both types of viruses. This observation provides the first evidence for a ZIKV-neutralizing antibody response that derives from pre-existing immunity to DENV. However, the low frequency of these cross-reactive memory B cells within the polyclonal population of circulating memory B cells raises the question of the impact of such responses in the immune protection against ZIKV infection. In this sense, it would be helpful to determine whether multiple DENV infections could boost such memory B cells with a broad spectrum of neutralization, in the same manner as the DENV1-induced cross-neutralizing response [44].

At the polyclonal level, it was confirmed that while a primary ZIKV infection could induce predominantly ZIKV-specific antibodies that poorly cross-react with the four DENV serotypes, a ZIKV infection in individuals with previous DENV infections results in the production of both ZIKV-specific and DENV-cross-reactive antibodies [45]. Interestingly, depletion of these DENV-cross-reactive antibodies did not affect the level of anti-ZIKV neutralizing antibodies, suggesting that ZIKV-specific neutralizing antibodies are produced after ZIKV infection regardless of previous DENV immunity [45].

Thus, while it is clear that DENV infections can induce the production of cross-reactive antibodies that recognize the four DENV serotypes, and to a lesser extent ZIKV, in most cases, they do not induce durable, high-level ZIKV cross-neutralizing antibodies [45].

Similarly, upon primary or secondary DENV infections, it is now well established that the level of DENV-specific neutralizing antibodies decay rapidly in the absence of re-exposure, or more slowly in endemic settings [46–49], and individuals with high neutralizing antibody titers have lower probability of symptomatic infections, in comparison with individuals with low neutralizing antibody titers [49]. In this sense, a positive correlation was clearly established between the low level of pre-existing anti-DENV antibodies and the severity of secondary dengue disease in humans, with a higher risk of severe dengue with anti-DENV antibody titers of 1:21 to 1:80 and a stronger protection against symptomatic dengue disease observed with higher antibody levels [19,50]. More recently and finally, while type-specific neutralizing responses were initially thought to elicit life-long immunity against homologous reinfections [51,52], there is now clear evidence that homologous reinfection can occur in the presence of serotype-specific neutralizing antibodies, leading to symptomatic disease [53,54]. While in some cases these homotypic reinfections have been observed more than 10 years after the initial infection [53], in other cases, such reinfections were identified with a shorter time interval of 1 to 2 years between the successive infections [54]. In both situations, the specific level of pre-existing anti-DENV antibodies provides an explanation for the homologous serotype reinfection and symptomatic disease observed in individuals from different endemic regions [53,54]. Although it is too early to determine whether the same correlation applies to ZIKV infection, it seems likely that the decay in the level of ZIKV-specific neutralizing antibodies in the absence of re-exposure to this virus should affect the outcome of future ZIKV infection and disease.

3. T Cell Responses against DENV and Prospects for a Vaccine

Although the exact role of T cells during dengue virus infection and disease is still a matter of debate [55], there is currently a growing body of evidence supporting a protective role for T cells in dengue virus infection, both in human and mouse studies [56,57].

Briefly, similarly to the ADE phenomenon hypothesis associated to cross-reactive antibodies, it was proposed that DENV-specific T cells could play a detrimental role during secondary dengue infection. In this “original antigenic sin” scenario, an expansion of cross-reactive T cells with higher avidity to the previous infecting serotype would mask the specific T cell response against secondary infection, and would result in less efficient elimination of DENV-infected cells [58–60]. These cross-reactive

T cells, stimulated upon a secondary infection with a different serotype, also displayed quantitative and qualitative differences in their response to the cross-reactive epitope [61], with higher ratios of Tumor Necrosis Factor alpha (TNF- α) to Interferon gamma (IFN- γ)-producing CD4 T cells [62], suboptimal degranulation but high cytokine production [63], or more specifically impaired IFN- γ production [64].

However, in spite of these studies, the direct demonstration of a pathogenic role of DENV-specific T cells in patients experiencing natural secondary dengue infection is still missing, and recent reports do not support a causative role for cross-reactive T cells in the pathogenesis of dengue hemorrhagic fever during secondary infections [65,66]. Indeed, increased frequencies of DENV-specific CD4⁺ and CD8⁺ T cells were detected in school children who subsequently experienced subclinical infection, in comparison with symptomatic secondary DENV infections [67]. In addition, protection or susceptibility to severe dengue disease has been associated with the expression of certain Human Leukocyte Antigen (HLA) molecules [68–73] and a beneficial function of CD8⁺ T cells against DENV infection was demonstrated after depletion of the CD8 T-cell compartment in interferon- α receptor knock-out mice (*ifnar*^{-/-}) [74]. More strikingly, a strong correlation was established between protection against severe dengue and a polyfunctional memory CD8⁺ T cell response with a high magnitude in healthy dengue-immune individuals [75]. Finally and more recently, we observed a higher activation of Natural Killer (NK) cells and T cells in asymptomatic dengue viral infection; different T cell populations proliferate more and have an activated phenotype, with increased pathogen recognition, signal transduction and higher cytotoxic activity in asymptomatic DENV infected individuals compared to individuals with symptoms during DENV infection [76].

While CD4⁺ T cell responses are mainly directed toward the structural proteins capsid (C) and envelope (E) and the non-structural protein NS1, similarly to DENV-specific B cells that target prM, E and NS1, most CD8 T cell epitopes reside in the non-structural proteins NS3, NS4B and NS5, and to a lesser extent in the structural protein E [60,75,77–80].

Besides the identification of DENV-specific T cell epitopes recognized by activated T cells, these studies contributed to identifying correlates of protection, as the result of an efficient T cell activation by immunodominant peptides restricted by certain HLA alleles, and a robust and polyfunctional CD4 or CD8 response [75,81–85]. This is true, for example, for the HLA-B*0702 and -B*3501 class I molecules, and the HLA-A*0101 and -A*2402 associated with a high and a low CD8 T-response frequency and magnitude, respectively [75]. The same goes for the HLA-DRB1*0401 or -DRB1*0802 class II alleles, which are associated with an increased resistance or susceptibility to severe dengue disease, respectively, and the phenotype of responding cytotoxic CD4 T cells [81].

4. Identification ZIKV-Specific and DENV/ZIKV Cross-Reactive T Cells

In contrast to a large number of studies on T cell responses to DENV infection, to date, relatively little is known regarding the analysis of T cell responses to ZIKV infection.

First predictions of ZIKV T cell antigens were conducted by modelling potential epitopes that could bind to different HLA class I or class II alleles, from the ZIKV proteome [86–88], and by identifying short peptides targeted by DENV-specific CD8⁺ T cells with conserved sequences between DENV and ZIKV [89], with the underlying assumption that these epitopes should stimulate cross-reactive T cells after sequential DENV and ZIKV infection.

While such sequence conservation between several immunogenic CD8⁺ T cell peptides from DENV and the corresponding ZIKV sequence was highlighted in the E, NS1, NS3 and NS5 proteins [89], a study revealed that memory T cells against NS1 or E proteins were poorly cross-reactive, even in donors previously infected by DENV [24].

To identify first the dominant epitopes of ZIKV recognized by CD8⁺ T cells in the context of human HLA class I molecules, and to clarify the protective role of CD8⁺ T cells in ZIKV infection, different mouse models have been used. Type I interferon receptor-deficient mice expressing human HLA class I molecules (HLA-B*0702 or HLA-A*0101), which are susceptible to DENV and ZIKV

infection, enabled identification of ZIKV peptides that are targeted by CD8⁺ T cells, and to show that dengue or Zika virus infections can induce the development of cross-reactive CD8⁺ T cells that are protective against ZIKV infection [90]. In HLA class II transgenic mice, CD4 immunodominant epitopes have also been mapped from ZIKV Envelope, and from non-structural proteins NS1, NS3 and NS5, among which several peptides in the envelope revealed cross-reactivity with other flaviviruses [91] (Figure 1).




Figure 1. Schematic representation of class I- and class II-restricted cross-reactive Zika virus (ZIKV) epitopes identified in humans and transgenic mice. The ZIKV genome is not to scale. E: Envelope; HLA: Human Leukocyte Antigen; M: Protein M; NS: Non-Structural Proteins; pr: Precursor; Tg: Transgenic; UTR: Untranslated Region.

In another mouse model, which lacks the type I interferon receptor in a subset of myeloid cells [92], adoptive transfer of ZIKV-immune CD8⁺ T cells was shown to inhibit primary ZIKV infection and replication [93]. Likewise, in *ifnar*^{-/-} and in wild-type mice, adoptive transfer and depletion experiments have demonstrated that DENV-immune CD8⁺ T cells, but not DENV-immune sera can mediate cross-protective responses against ZIKV infection [94]. Although a correlation was previously established in mice and rhesus monkeys between the level of neutralizing antibodies induced upon ZIKV vaccination and the immune protection against ZIKV challenge [95,96], these studies highlight the essential role of CD8⁺ T cells in providing a protective immunity, in the presence or not of pre-existing DENV immunity, a finding that could help define future strategies of vaccine against DENV and ZIKV [97]. Finally, wild-type immunocompetent mice, *ifnar*^{-/-} mice and Interferon Alpha Receptor (IFNAR)-depleted *rag1*^{-/-} mice, have helped to characterize the phenotype of responding T cells after ZIKV infection and to demonstrate the protective effect of CD8⁺ T cells against ZIKV infection in various organs, including the brain and testes [98–100]. However, while these studies clearly demonstrate a major role for CD8⁺ T cells in controlling ZIKV infection, they all relied on experiments performed in mice with an altered immune system, a model that supports DENV or ZIKV infection but does not reflect the human situation. In addition, these mouse models do not take into account the human innate immune response against these flaviviruses, more specifically the NK cells that kill infected cells through the recognition of viral peptides in the context of HLA-C*0102 [101]. In summary, even though HLA transgenic mice expressing a single HLA allele do not mimic the human situation, the identification of immunodominant epitopes in these transgenic mice constitutes an important step towards preclinical evaluation of T-cell based vaccines.

For this purpose, to formally identify ZIKV-derived peptides in humans, and to determine whether a previous DENV infection can activate cross-reactive T cells, functional studies have been performed using human Peripheral Blood Mononuclear Cells (PBMC) collected from ZIKV endemic areas. From blood donors from different countries in Central and South America, the majority of the CD8⁺ T cell responses in DENV seronegative individuals were shown to be directed against structural proteins, whereas in DENV seropositive donors, a large proportion of CD8⁺ T cell responses were directed against the non-structural proteins with cross-reactivity for peptides in NS2A, NS3, and NS5 [102] (Figure 1). Strikingly, cross-reactive CD4⁺ and CD8⁺ T cells recognized peptides with identical or highly conserved sequences between DENV and ZIKV, with a higher magnitude of response, showing that DENV-specific memory CD8⁺ T cells can enhance the T cell responses to ZIKV.

In light of the low protection observed in individuals DENV-seronegative at the time of vaccination with Dengvaxia, which lacks DENV non-structural proteins and fails to induce a competent T cell

response [103], and taking into account the role of CD8⁺ T cells in preventing ADE [104,105] it is now becoming obvious that efficient vaccine candidates against DENV and ZIKV should be formulated to include CD4 and CD8 T cell epitopes, either alone, or in combination with B cell epitopes.

Based on the recent identification of these T cell epitopes derived from ZIKV, work is currently ongoing to define a minimal antigen, which includes the most immunodominant peptides recognized by cross-reactive T cells, and which can induce a long lasting immune protection against DENV and ZIKV infection and disease.

5. Conclusions

Our understanding of the immune response to ZIKV has dramatically increased in the last years through in vitro studies with polyclonal and monoclonal antibodies from ZIKV-infected individuals, and through the identification of the epitopes inducing a strong memory and cross-reactive T cell response. While several animal models that could mimic the human situation have been developed to clarify the beneficial or detrimental role of the different immune mediators in disease protection, further phenotypic analyses of ZIKV-specific T cells, in asymptomatic or symptomatic donors will help define correlates of protection in natural immunity and vaccination against ZIKV infection and disease.

Acknowledgments: The authors would like to acknowledge the Ecole Doctoral BioSPC (Université Paris Diderot) and the Colombian Department of Science, Technology and Innovation (COLCIENCIAS) for the doctoral scholarship program (Nb. 679) of F.G.D. The authors acknowledge also support from the “Integrative Biology of Emerging Infectious Diseases” Labex (Laboratoire d’excellence) grant ANR Num. ANR-10-LABX-62-IBEID (French Government’s “Investissements d’Avenir” program), the European Commission Seventh Framework Program (PF7/2007-2013) for the DENFREE project under Grant Agreement n° 282 378 and Prix Duquesne.

Author Contributions: C.R., F.G.D., E.S.-L. and A.S. provided valuable scientific discussion and expertise in the field of flavivirus virology. C.R. wrote the manuscript.

Conflicts of Interest: The authors declare no conflicts of interest. The founding sponsors had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, and in the decision to publish the results.

References

1. Kuno, G.; Chang, G.J.; Tsuchiya, K.R.; Karabatsos, N.; Cropp, C.B. Phylogeny of the genus Flavivirus. *J. Virol.* **1998**, *72*, 73–83. [[PubMed](#)]
2. Dick, G.W.; Kitchen, S.F.; Haddow, A.J. Zika virus. I. Isolations and serological specificity. *Trans. R. Soc. Trop. Med. Hyg.* **1952**, *46*, 509–520. [[CrossRef](#)]
3. Duffy, M.R.; Chen, T.-H.; Hancock, T.; Powers, A.M.; Kool, J.L.; Lanciotti, R.S.; Pretrick, M.; Marfel, M.; Holzbauer, S.; Dubray, C.; et al. Zika virus outbreak on Yap Island, Federated States of Micronesia. *N. Engl. J. Med.* **2009**, *360*, 2536–2543. [[CrossRef](#)] [[PubMed](#)]
4. Lanciotti, R.S.; Kosoy, O.L.; Laven, J.J.; Velez, J.O.; Lambert, A.J.; Johnson, A.J.; Stanfield, S.M.; Duffy, M.R. Genetic and serologic properties of Zika virus associated with an epidemic, Yap State, Micronesia, 2007. *Emerg. Infect. Dis.* **2008**, *14*, 1232–1239. [[CrossRef](#)] [[PubMed](#)]
5. Cao-Lormeau, V.M.; Roche, C.; Teissier, A.; Robin, E.; Berry, A.-L.; Mallet, H.-P.; Sall, A.A.; Musso, D. Zika virus, French polynesia, South pacific, 2013. *Emerg. Infect. Dis.* **2014**, *20*, 1085–1086. [[CrossRef](#)] [[PubMed](#)]
6. Campos, G.S.; Bandeira, A.C.; Sardi, S.I. Zika virus outbreak, Bahia, Brazil. *Emerg. Infect. Dis.* **2015**, *21*, 1885–1886. [[CrossRef](#)] [[PubMed](#)]
7. Dupont-Rouzeyrol, M.; O’Connor, O.; Calvez, E.; Daurès, M.; John, M.; Grangeon, J.-P.; Gourinat, A.-C. Co-infection with Zika and dengue viruses in 2 patients, New Caledonia, 2014. *Emerg. Infect. Dis.* **2015**, *21*, 381–382. [[CrossRef](#)] [[PubMed](#)]
8. Zanoluca, C.; de Melo, V.C.A.; Mosimann, A.L.P.; dos Santos, G.I.V.; dos Santos, C.N.D.; Luz, K. First report of autochthonous transmission of Zika virus in Brazil. *Mem. Inst. Oswaldo Cruz* **2015**, *110*, 569–572. [[CrossRef](#)] [[PubMed](#)]

9. Pacheco, O.; Beltrán, M.; Nelson, C.A.; Valencia, D.; Tolosa, N.; Farr, S.L.; Padilla, A.V.; Tong, V.T.; Cuevas, E.L.; Espinosa-Bode, A.; et al. Zika virus disease in Colombia—Preliminary report. *N. Engl. J. Med.* **2016**. [[CrossRef](#)] [[PubMed](#)]
10. Oehler, E.; Watrin, L.; Larre, P.; Le Parc-Goffart, I.; Lastère, S.; Valour, F.; Baudouin, L.; Mallet, H.P.; Musso, D.; Ghawche, F. Zika virus infection complicated by Guillain-Barre syndrome—Case report, French Polynesia, December 2013. *Eurosurveillance* **2014**, *19*, 20720. [[CrossRef](#)] [[PubMed](#)]
11. Cao-Lormeau, V.M.; Blake, A.; Mons, S.; Lastère, S.; Roche, C.; Vanhomwegen, J.; Dub, T.; Baudouin, L.; Teissier, A.; Larre, P.; et al. Guillain-Barre Syndrome outbreak associated with Zika virus infection in French Polynesia: A case-control study. *Lancet* **2016**, *387*, 1531–1539. [[CrossRef](#)]
12. Cauchemez, S.; Besnard, M.; Bompard, P.; Dub, T.; Guillemette-Artur, P.; Eyrolle-Guignot, D.; Salje, H.; van Kerkhove, M.D.; Abadie, V.; Garel, C.; et al. Association between Zika virus and microcephaly in French Polynesia, 2013–15: A retrospective study. *Lancet* **2016**, *387*, 2125–2132. [[CrossRef](#)]
13. Soares De Araujo, J.S.; Teixeira Regis, C.; Silva Gomes, R.G.; Riberio Tavares, T.; Rocha dos Santos, C.; Melo Assunção, P.; Nóbrega, R.V.; Alves Pinto, D.F.; Dantas Bezerra, B.V.; da Silva Mattos, S. Microcephaly in north-east Brazil: A retrospective study on neonates born between 2012 and 2015. *Bull. World Health Organ.* **2016**, *94*, 835–840. [[CrossRef](#)] [[PubMed](#)]
14. Enfissi, A.; Codrington, J.; Roosblad, J.; Kazanji, M.; Rousset, D. Zika virus genome from the Americas. *Lancet* **2016**, *387*, 227–228. [[CrossRef](#)]
15. Musso, D.; Gubler, D.J. Zika Virus. *Clin. Microbiol. Rev.* **2016**, *29*, 487–524. [[CrossRef](#)] [[PubMed](#)]
16. Liu, Y.; Liu, J.; Du, S.; Shan, C.; Nie, K.; Zhang, R.; Li, X.-F.; Zhang, R.; Wang, T.; Qin, C.-F.; et al. Evolutionary enhancement of Zika virus infectivity in *Aedes aegypti* mosquitoes. *Nature* **2017**, *545*, 482–486. [[CrossRef](#)] [[PubMed](#)]
17. Metsky, H.C.; Matranga, C.B.; Wohl, S.; Schaffner, S.F.; Freije, C.A.; Winnicki, S.M.; West, K.; Qu, J.; Baniecki, M.L.; Gladden-Young, A.; et al. Zika virus evolution and spread in the Americas. *Nature* **2017**, *546*, 411–415. [[CrossRef](#)] [[PubMed](#)]
18. Yuan, L.; Huang, X.-Y.; Liu, Z.-Y.; Zhang, F.; Zhu, X.-L.; Yu, J.-Y.; Ji, X.; Xu, Y.-P.; Li, G.; Li, C.; et al. A single mutation in the prM protein of Zika virus contributes to fetal microcephaly. *Science* **2017**, *358*, 933–936. [[CrossRef](#)] [[PubMed](#)]
19. Katzelnick, L.C.; Coloma, J.; Harris, E. Dengue: Knowledge gaps, unmet needs, and research priorities. *Lancet Infect. Dis.* **2017**, *17*, e88–e100. [[CrossRef](#)]
20. Barba-Spaeth, G.; Dejnirattisai, W.; Rouvniski, A.; Vaney, M.-C.; Medits, I.; Sharma, A.; Simon-Lorière, E.; Sakuntabhai, A.; Cao-Lormeau, V.-M.; Haouz, A.; et al. Structural basis of potent Zika-dengue virus antibody cross-neutralization. *Nature* **2016**, *536*, 48–53. [[CrossRef](#)] [[PubMed](#)]
21. Sirohi, D.; Chen, Z.; Sun, L.; Klose, T.; Pierson, T.C.; Rossmann, M.G.; Kuhn, R.J. The 3.8 Å resolution cryo-EM structure of Zika virus. *Science* **2016**, *352*, 467–470. [[CrossRef](#)] [[PubMed](#)]
22. Xu, X.; Song, H.; Qi, J.; Liu, Y.; Wang, H.; Su, C.; Shi, Y.; Gao, G.F. Contribution of intertwined loop to membrane association revealed by Zika virus full-length NS1 structure. *EMBO J.* **2016**, *35*, 2170–2178. [[CrossRef](#)] [[PubMed](#)]
23. Priyamvada, L.; Quicke, K.M.; Hudson, W.H.; Onlamoon, N.; Sewatanon, J.; Edupuganti, S.; Pattanapanyasat, K.; Chokeybulkit, K.; Mulligan, M.J.; Wilson, P.C.; et al. Human antibody responses after dengue virus infection are highly cross-reactive to Zika virus. *Proc. Natl. Acad. Sci. USA* **2016**, *113*, 7852–7857. [[CrossRef](#)] [[PubMed](#)]
24. Stettler, K.; Beltramello, M.; Espinosa, D.A.; Graham, V.; Cassotta, A.; Bianchi, S.; Vanzetta, F.; Minola, A.; Jaconi, S.; Mele, F.; et al. Specificity, cross-reactivity, and function of antibodies elicited by Zika virus infection. *Science* **2016**, *353*, 823–826. [[CrossRef](#)] [[PubMed](#)]
25. Andrade, D.V.; Harris, E. Recent advances in understanding the adaptive immune response to Zika virus and the effect of previous flavivirus exposure. *Virus Res.* **2017**. [[CrossRef](#)] [[PubMed](#)]
26. Bardina, S.V.; Bunduc, P.; Tripathi, S.; Duehr, J.; Frere, J.J.; Brown, J.A.; Nachbagauer, R.; Foster, G.A.; Krysztof, D.; Tortorella, D.; et al. Enhancement of Zika virus pathogenesis by preexisting ant flavivirus immunity. *Science* **2017**, *356*, 175–180. [[CrossRef](#)] [[PubMed](#)]
27. Rogers, T.F.; Goodwin, E.C.; Briney, B.; Sok, D.; Beutler, N.; Strubel, A.; Nedellec, R.; Le, K.; Brown, M.E.; Burton, D.R.; et al. Zika virus activates de novo and cross-reactive memory B cell responses in dengue-experienced donors. *Sci. Immunol.* **2017**, *2*. [[CrossRef](#)] [[PubMed](#)]

28. Halstead, S.B. Dengue antibody-dependent enhancement: Knowns and unknowns. *Microbiol. Spectr.* **2014**, *2*. [[CrossRef](#)] [[PubMed](#)]
29. Priyamvada, L.; Hudson, W.; Ahmed, R.; Wrammert, J. Humoral cross-reactivity between Zika and dengue viruses: Implications for protection and pathology. *Emerg. Microbes Infect.* **2017**, *6*, e33. [[CrossRef](#)] [[PubMed](#)]
30. Xu, X.; Vaughan, K.; Weiskopf, D.; Grifoni, A.; Diamond, M.S.; Sette, A.; Peters, B. Identifying Candidate Targets of Immune Responses in Zika Virus Based on Homology to Epitopes in Other Flavivirus Species. *PLoS Curr.* **2016**, *8*. [[CrossRef](#)] [[PubMed](#)]
31. Thomas, S.J.; Nisalak, A.; Anderson, K.B.; Libraty, D.H.; Kalayanarooj, S.; Vaughn, D.W.; Putnak, R.; Gibbons, R.V.; Jarman, R.; Endy, T.P. Dengue plaque reduction neutralization test (PRNT) in primary and secondary dengue virus infections: How alterations in assay conditions impact performance. *Am. J. Trop. Med. Hyg.* **2009**, *81*, 825–833. [[CrossRef](#)] [[PubMed](#)]
32. De Alwis, R.; de Silva, A.M. Measuring antibody neutralization of dengue virus (DENV) using a flow cytometry-based technique. *Methods Mol. Biol.* **2014**, *1138*, 27–39. [[PubMed](#)]
33. Calvo, E.P.; Sanchez-Quete, F.; Duran, S.; Sandoval, I.; Castellanos, J.E. Easy and inexpensive molecular detection of dengue, chikungunya and zika viruses in febrile patients. *Acta Trop.* **2016**, *163*, 32–37. [[CrossRef](#)] [[PubMed](#)]
34. Aubry, M.; Teissier, A.; Huart, M.; Merceron, S.; Vanhomwegen, J.; Roche, C.; Vial, A.-L.; Teururai, S.; Sicard, S.; Paulous, S.; et al. Zika virus seroprevalence, French Polynesia, 2014–2015. *Emerg. Inf. Dis.* **2017**, *23*, 669–672. [[CrossRef](#)] [[PubMed](#)]
35. De Alwis, R.; Smith, S.A.; Olivarez, N.P.; Messer, W.B.; Huynh, J.P.; Wahala, W.M.P.B.; White, L.J.; Diamond, M.S.; Baric, R.S.; Crowe, J.E.; et al. Identification of human neutralizing antibodies that bind to complex epitopes on dengue virions. *Proc. Natl. Acad. Sci. USA* **2012**, *109*, 7439–7444. [[CrossRef](#)] [[PubMed](#)]
36. Halstead, S.B. Neutralization and antibody-dependent enhancement of dengue viruses. *Adv. Virus Res.* **2003**, *60*, 421–467. [[PubMed](#)]
37. Patel, B.; Longo, P.; Miley, M.J.; Montoya, M.; Harris, E.; de Silva, A.M. Dissecting the human serum antibody response to secondary dengue virus infections. *PLoS Neglect. Trop. Dis.* **2017**, *11*, e0005554. [[CrossRef](#)] [[PubMed](#)]
38. Olkowski, S.; Forshey, B.M.; Morrison, A.C.; Rocha, C.; Vilcarromero, S.; Halsey, E.S.; Kochel, T.J.; Scott, T.W.; Stoddard, S.T. Reduced risk of disease during postsecondary dengue virus infections. *J. Infect. Dis.* **2013**, *208*, 1026–1033. [[CrossRef](#)] [[PubMed](#)]
39. Beltramello, M.; Williams, K.L.; Simmons, C.P.; Macagno, A.; Simonelli, L.; Ha Quyen, N.T.; Sukupolvi-Petty, S.; Navarro-Sanchez, E.; Young, P.R.; de Silva, A.M.; et al. The human immune response to dengue virus is dominated by highly cross-reactive antibodies endowed with neutralizing and enhancing activity. *Cell Host Microbe* **2010**, *8*, 271–283. [[CrossRef](#)] [[PubMed](#)]
40. Dejnirattisai, W.; Jumnainsong, A.; Onsirisakul, N.; Fitton, P.; Vasanawathana, S.; Limpitikul, W.; Puttikhunt, C.; Edwards, C.; Duangchinda, T.; Supasa, S.; et al. Cross-reacting antibodies enhance dengue virus infection in humans. *Science* **2010**, *328*, 745–748. [[CrossRef](#)] [[PubMed](#)]
41. Smith, S.A.; Zhou, Y.; Olivarez, N.P.; Broadwater, A.H.; de Silva, A.M.; Crowe, J.E., Jr. Persistence of circulating memory B cell clones with potential for dengue virus disease enhancement for decades following infection. *J. Virol.* **2012**, *86*, 2665–2675. [[CrossRef](#)] [[PubMed](#)]
42. Dejnirattisai, W.; Supasa, P.; Wongwiwat, W.; Rouvinski, A.; Barba-Spaeth, G.; Duangchinda, T.; Sakuntabhai, A.; Cao-Lormeau, V.-M.; Malasit, P.; Rey, F.A.; et al. Dengue virus sero-cross-reactivity drives antibody-dependent enhancement of infection with zika virus. *Nat. Immunol.* **2016**, *17*, 1102–1108. [[CrossRef](#)] [[PubMed](#)]
43. Priyamvada, L.; Cho, A.; Onlamoon, N.; Zheng, N.-Y.; Huang, M.; Kovalenkov, Y.; Chokephaibulkit, K.; Angkasekwinai, N.; Pattanapanyasat, K.; Ahmed, R.; et al. B cell responses during secondary dengue virus infection are dominated by highly cross-reactive, memory-derived plasmablasts. *J. Virol.* **2016**, *90*, 5574–5585. [[CrossRef](#)] [[PubMed](#)]
44. Robbiani, D.F.; Bozzacco, L.; Keeffe, J.R.; Khouri, R.; Olsen, P.C.; Gazumyan, A.; Schaefer-Babajew, D.; Avila-Ros, S.; Nogueira, L.; Patel, R.; et al. Recurrent potent human neutralizing antibodies to Zika virus in Brazil and Mexico. *Cell* **2017**, *169*, 597–609. [[CrossRef](#)] [[PubMed](#)]

45. Collins, M.H.; McGowan, E.; Jadi, R.; Young, E.; Lopez, C.A.; Baric, R.S.; Lazear, H.M.; de Silva, A.M. Lack of durable cross-neutralizing antibodies against Zika virus from dengue virus infection. *Emerg. Infect. Dis.* **2017**, *23*, 773–781. [[CrossRef](#)] [[PubMed](#)]
46. Puschnik, A.; Lau, L.; Cromwell, E.A.; Balmaseda, A.; Zompi, S.; Harris, E. Correlation between dengue-specific neutralizing antibodies and serum avidity in primary and secondary dengue virus 3 natural infections in humans. *PLoS Neglect. Trop. Dis.* **2013**, *7*, e2274. [[CrossRef](#)] [[PubMed](#)]
47. Corbett, K.S.; Katzelnick, L.; Tissera, H.; Amerasinghe, A.; Dharshan de Silva, A.; de Silva, A.M. Preexisting neutralizing antibody responses distinguish clinically inapparent and apparent dengue virus infections in a Sri Lankan pediatric cohort. *J. Infect. Dis.* **2015**, *211*, 590–599. [[CrossRef](#)] [[PubMed](#)]
48. Clapham, H.E.; Rodriguez-Barraquer, I.; Azman, A.S.; Althouse, B.M.; Salje, H.; Gibbons, R.V.; Rothman, A.L.; Jarman, R.G.; Nisalak, A.; Thaisomboonsuk, B.; et al. Dengue virus (DENV) neutralizing antibody kinetics in children after symptomatic primary and postprimary DENV infection. *J. Infect. Dis.* **2016**, *213*, 1428–1435. [[CrossRef](#)] [[PubMed](#)]
49. Katzelnick, L.C.; Montoya, M.; Gresh, L.; Balmaseda, A.; Harris, E. Neutralizing antibody titers against dengue virus correlate with protection from symptomatic infection in a longitudinal cohort. *Proc. Natl. Acad. Sci. USA* **2016**, *113*, 728–733. [[CrossRef](#)] [[PubMed](#)]
50. Katzelnick, L.C.; Gresh, L.; Halloran, M.E.; Mercado, J.C.; Kuan, G.; Gordon, A.; Balmaseda, A.; Harris, E. Antibody-dependent enhancement of severe dengue disease in humans. *Science* **2017**, *358*, 929–932. [[CrossRef](#)] [[PubMed](#)]
51. Sabin, A.B. Research on dengue during World War II. *Am. J. Trop. Med. Hyg.* **1952**, *1*, 30–50. [[CrossRef](#)] [[PubMed](#)]
52. Halstead, S.B.; Casals, J.; Shotwell, H.; Palumbo, N. Studies on the immunization of monkeys against dengue. I. Protection derived from single and sequential virus infections. *Am. J. Trop. Med. Hyg.* **1973**, *22*, 365–374. [[CrossRef](#)] [[PubMed](#)]
53. Forshey, B.M.; Reiner, R.C.; Olkowski, S.; Morrison, A.C.; Espinoza, A.; Long, K.C.; Vilcarrromero, S.; Casanova, W.; Wearing, H.J.; Halsey, E.S.; et al. Incomplete protection against dengue virus type 2 re-infection in Peru. *PLoS Neglect. Trop. Dis.* **2016**, *10*, e0004398. [[CrossRef](#)] [[PubMed](#)]
54. Waggoner, J.J.; Balmaseda, A.; Gresh, L.; Sahoo, M.K.; Montoya, M.; Wang, C.; Abeynayake, J.; Kuan, G.; Pinsky, B.A.; Harris, E. Homotypic dengue virus reinfections in Nicaraguan children. *J. Infect. Dis.* **2016**, *214*, 986–993. [[CrossRef](#)] [[PubMed](#)]
55. Culshaw, A.; Mongkolsapaya, J.; Screaton, G.R. The immunopathology of dengue and Zika virus infections. *Curr. Opin. Immunol.* **2017**, *48*, 1–6. [[CrossRef](#)] [[PubMed](#)]
56. Rivino, L. T cell immunity to dengue virus and implications for vaccine design. *Expert Rev. Vaccines* **2016**, *15*, 443–453. [[CrossRef](#)] [[PubMed](#)]
57. Katzelnick, L.C.; Harris, E. Participants in the Summit on Dengue Immune Correlates of P Immune correlates of protection for dengue: State of the art and research agenda. *Vaccine* **2017**, *35*, 4659–4669. [[CrossRef](#)] [[PubMed](#)]
58. Mongkolsapaya, J.; Dejnirattisai, W.; Xu, X.-N.; Vasanawathana, S.; Tangthawornchaikul, N.; Chairunsri, A.; Sawasdivorn, S.; Duangchinda, T.; Dong, T.; Rowland-Jones, S.; et al. Original antigenic sin and apoptosis in the pathogenesis of dengue hemorrhagic fever. *Nat. Med.* **2003**, *9*, 921–927. [[CrossRef](#)] [[PubMed](#)]
59. Yachi, P.P.; Ampudia, J.; Zal, T.; Gascoigne, N.R. Altered peptide ligands induce delayed CD8-T cell receptor interaction—A role for CD8 in distinguishing antigen quality. *Immunity* **2006**, *25*, 203–211. [[CrossRef](#)] [[PubMed](#)]
60. Duangchinda, T.; Dejnirattisai, W.; Vasanawathana, S.; Limpitikul, W.; Tangthawornchaikul, N.; Malasit, P.; Mongkolsapaya, J.; Screaton, G. Immunodominant T-cell responses to dengue virus NS3 are associated with DHF. *Proc. Natl. Acad. Sci. USA* **2010**, *107*, 16922–16927. [[CrossRef](#)] [[PubMed](#)]
61. Bashyam, H.S.; Green, S.; Rothman, A.L. Dengue virus-reactive CD8⁺ T cells display quantitative and qualitative differences in their response to variant epitopes of heterologous viral serotypes. *J. Immunol.* **2006**, *176*, 2817–2824. [[CrossRef](#)] [[PubMed](#)]
62. Mangada, M.M.; Rothman, A.L. Altered cytokine responses of dengue-specific CD4⁺ T cells to heterologous serotypes. *J. Immunol.* **2005**, *175*, 2676–2683. [[CrossRef](#)] [[PubMed](#)]

63. Mongkolsapaya, J.; Duangchinda, T.; Dejnirattisai, W.; Vasanawathana, S.; Avirutnan, P.; Jairungsri, A.; Khemnu, N.; Tangthawornchaikul, N.; Chotiyarnwong, P.; Sae-Jang, K.; et al. T cell responses in dengue hemorrhagic fever: Are cross-reactive T cells suboptimal? *J. Immunol.* **2006**, *176*, 3821–3829. [[CrossRef](#)] [[PubMed](#)]
64. Zivny, J.; DeFronzo, M.; Jarry, W.; Jameson, J.; Cruz, J.; Ennis, F.A.; Rothman, A.L. Partial agonist effect influences the CTL response to a heterologous dengue virus serotype. *J. Immunol.* **1999**, *163*, 2754–2760. [[PubMed](#)]
65. Weiskopf, D.; Angelo, M.A.; Bangs, D.J.; Sidney, J.; Paul, S.; Peters, B.; de Silva, A.D.; Lindow, J.C.; Diehl, S.A.; Whitehead, S.; et al. The human CD8⁺ T cell responses induced by a live attenuated tetravalent dengue vaccine are directed against highly conserved epitopes. *J. Virol.* **2015**, *89*, 120–128. [[CrossRef](#)] [[PubMed](#)]
66. Weiskopf, D.; Cerpas, C.; Angelo, M.A.; Bangs, D.J.; Sidney, J.; Paul, S.; Peters, B.; Sanches, F.P.; Silvera, C.G.T.; Costa, P.R.; et al. Human CD8⁺ T-cell responses against the 4 dengue virus serotypes are associated with distinct patterns of protein targets. *J. Infect. Dis.* **2015**, *212*, 1743–1751. [[CrossRef](#)] [[PubMed](#)]
67. Hatch, S.; Endy, T.P.; Thomas, S.; Mathew, A.; Potts, J.; Pazoles, P.; Libraty, D.H.; Gibbons, R.; Rothman, A.L. Intracellular cytokine production by dengue virus-specific T cells correlates with subclinical secondary infection. *J. Infect. Dis.* **2011**, *203*, 1282–1291. [[CrossRef](#)] [[PubMed](#)]
68. Loke, H.; Bethell, D.B.; Phuong, C.X.T.; Dung, M.; Scheinder, J.; White, N.J.; Day, N.P.; Farrar, J.; Hill, A.V.S. Strong HLA class I-restricted T cell responses in dengue hemorrhagic fever: A double-edged sword? *J. Infect. Dis.* **2001**, *184*, 1369–1373. [[CrossRef](#)] [[PubMed](#)]
69. Stephens, H.A.; Klaythong, R.; Sirikong, M.; Vaughn, D.W.; Green, S.; Kalayanaoij, S.; Endy, T.P.; Libraty, D.H.; Nisalak, A.; Innis, B.L.; et al. HLA-A and -B allele associations with secondary dengue virus infections correlate with disease severity and the infecting viral serotype in ethnic Thais. *Tissue Antigens* **2002**, *60*, 309–318. [[CrossRef](#)] [[PubMed](#)]
70. Sierra, B.; Alegre, R.; Pérez, A.B.; García, G.; Sturn-Ramirez, K.; Obasanjo, O.; Aguirre, E.; Alvarez, M.; Rodriguez-Roche, R.; Valdés, L.; et al. HLA-A, -B, -C, and -DRB1 allele frequencies in Cuban individuals with antecedents of dengue 2 disease: Advantages of the Cuban population for HLA studies of dengue virus infection. *Hum. Immunol.* **2007**, *68*, 531–540. [[CrossRef](#)] [[PubMed](#)]
71. Nguyen, T.P.; Kikuchi, M.; Huong, V.T.Q.; Ha, D.Q.; Thuy, T.T.; Tham, V.D.; Tuan, H.M.; Tuong, V.V.; Nga, C.T.P.; Dat, T.V.; et al. Protective and enhancing HLA alleles, HLA-DRB1*0901 and HLA-A*24, for severe forms of dengue virus infection, dengue hemorrhagic fever and dengue shock syndrome. *PLoS Neglect. Trop. Dis.* **2008**, *2*, e304.
72. Coffey, L.L.; Mertens, E.; Brehin, A.-C.; Fernandez-Garcia, M.D.; Amara, A.; Després, P.; Sakuntabhai, A. Human genetic determinants of dengue virus susceptibility. *Microbes Infect.* **2009**, *11*, 143–156. [[CrossRef](#)] [[PubMed](#)]
73. Malavige, G.N.; Rostron, T.; Rohanachandra, L.T.; Fernando, N.; Dharshan de Silva, A.; Liyanage, M.; Ogg, G. HLA class I and class II associations in dengue viral infections in a Sri Lankan population. *PLoS ONE* **2011**, *6*, e20581. [[CrossRef](#)] [[PubMed](#)]
74. Yauch, L.E.; Zellweger, R.M.; Kotturi, M.F.; Qutubuddin, A.; Sidney, J.; Peters, B.; Prestwood, T.R.; Sette, A.; Shresta, S. A protective role for dengue virus-specific CD8⁺ T cells. *J. Immunol.* **2009**, *182*, 4865–4873. [[CrossRef](#)] [[PubMed](#)]
75. Weiskopf, D.; Angelo, M.A.; de Azeredo, E.L.; Sidney, J.; Greenbaum, J.A.; Fernando, A.N.; Broadwater, A.; Kolla, R.V.; de Silva, A.D.; de Silva, A.M.; et al. Comprehensive analysis of dengue virus-specific responses supports an HLA-linked protective role for CD8⁺ T cells. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, E2046–E2053. [[CrossRef](#)] [[PubMed](#)]
76. Simon-Lorière, E.; Duong, V.; Tawfik, A.; Ung, S.; Ly, S.; Casadémont, I.; Prot, M.; Courtejoie, N.; Bleakley, K.; Buchy, P.; et al. Increased adaptive immune responses and proper feedback regulation protect against clinical dengue. *Sci. Transl. Med.* **2017**, *9*. [[CrossRef](#)] [[PubMed](#)]
77. Livingston, P.G.; Kurane, I.; Dai, L.C.; Okamoto, Y.; Lai, C.J.; Men, R.; Karaki, S.; Takiguchi, M.; Ennis, F.A. Dengue virus-specific, HLA-B35-restricted, human CD8⁺ cytotoxic T lymphocyte (CTL) clones. Recognition of NS3 amino acids 500 to 508 by CTL clones of two different serotype specificities. *J. Immunol.* **1995**, *154*, 1287–1295. [[PubMed](#)]

78. Kurane, I.; Zeng, L.; Brinton, M.A.; Ennis, F.A. Definition of an epitope on NS3 recognized by human CD4⁺ cytotoxic T lymphocyte clones cross-reactive for dengue virus types 2, 3, and 4. *Virology* **1998**, *240*, 169–174. [[CrossRef](#)] [[PubMed](#)]
79. Rivino, L.; Kumaran, E.A.P.; Jovanovic, V.; Nadua, K.; Teo, E.W.; Pang, S.W.; Teo, G.H.; Hao Gan, V.C.; Lye, D.C.; et al. Differential targeting of viral components by CD4⁺ versus CD8⁺ T lymphocytes in dengue virus infection. *J. Virol.* **2013**, *87*, 2693–2706. [[CrossRef](#)] [[PubMed](#)]
80. Rivino, L.; Tan, A.T.; Chia, A.; Kumaran, E.A.; Grotenbreg, G.M.; MacAry, P.A.; Bertoletti, A. Defining CD8⁺ T cell determinants during human viral infection in populations of Asian ethnicity. *J. Immunol.* **2013**, *191*, 4010–4019. [[CrossRef](#)] [[PubMed](#)]
81. Weiskopf, D.; Bangs, D.J.; Sidney, J.; Kolla, R.V.; de Silva, A.D.; de Silva, A.M.; Crotty, S.; Peters, B.; Sette, A. Dengue virus infection elicits highly polarized CX3CR1⁺ cytotoxic CD4⁺ T cells associated with protective immunity. *Proc. Natl. Acad. Sci. USA* **2015**, *112*, E4256–E4263. [[CrossRef](#)] [[PubMed](#)]
82. Zeng, L.; Kurane, I.; Okamoto, Y.; Ennis, F.A.; Brinton, M.A. Identification of amino acids involved in recognition by dengue virus NS3-specific, HLA-DR15-restricted cytotoxic CD4⁺ T-cell clones. *J. Virol.* **1996**, *70*, 3108–3117. [[PubMed](#)]
83. Okamoto, Y.; Kurane, I.; Leporati, A.M.; Ennis, F.A. Definition of the region on NS3 which contains multiple epitopes recognized by dengue virus serotype-cross-reactive and flavivirus-cross-reactive, HLA-DPw2-restricted CD4⁺ T cell clones. *J. Gen. Virol.* **1998**, *79*, 697–704. [[CrossRef](#)] [[PubMed](#)]
84. Simmons, C.P.; Dong, T.; Chau, N.V.; Phuong Dung, N.T.; Bich Chau, T.N.; Thu Thao, L.T.; Dung, N.T.; Hien, T.T.; Rowland-Jones, S.; Farrar, J. Early T-cell responses to dengue virus epitopes in Vietnamese adults with secondary dengue virus infections. *J. Virol.* **2005**, *79*, 5665–5675. [[CrossRef](#)] [[PubMed](#)]
85. Nascimento, E.J.; Mailliard, R.B.; Khan, A.M.; Sidney, J.; Sette, A.; Guzman, N.; Paulaitis, M.; de Melo, A.B.; Cordeiro, M.T.; Gil, L.V.G.; et al. Identification of conserved and HLA promiscuous DENV3 T-cell epitopes. *PLoS Neglect. Trop. Dis.* **2013**, *7*, e2497. [[CrossRef](#)] [[PubMed](#)]
86. Alam, A.; Ali, S.; Ahamad, S.; Malik, M.Z.; Ishrat, R. From ZikV genome to vaccine: In silico approach for the epitope-based peptide vaccine against Zika virus envelope glycoprotein. *Immunology* **2016**, *149*, 386–399. [[CrossRef](#)] [[PubMed](#)]
87. Dar, H.; Zaheer, T.; Talha Rehman, M.; Ali, A.; Javed, A.; Ayub Khan, G.; Mujtaba Babar, M.; Waheed, Y. Prediction of promiscuous T-cell epitopes in the Zika virus polyprotein: An in silico approach. *Asian Pac. J. Trop. Med.* **2016**, *9*, 844–850. [[CrossRef](#)] [[PubMed](#)]
88. Usman Mirza, M.; Rafique, S.; Ali, A.; Munir, M.; Ikram, N.; Manan, A.; Salo-Ahen, O.M.H.; Idrees, M. Towards peptide vaccines against Zika virus: Immunoinformatics combined with molecular dynamics simulations to predict antigenic epitopes of Zika viral proteins. *Sci. Rep.* **2016**, *6*, 37313. [[CrossRef](#)] [[PubMed](#)]
89. Rivino, L.; Lim, M.Q. CD4⁺ and CD8⁺ T-cell immunity to Dengue—Lessons for the study of Zika virus. *Immunology* **2016**, *150*, 146–154. [[CrossRef](#)] [[PubMed](#)]
90. Wen, J.; Weihao Tang, W.; Sheets, N.; Ellison, J.; Sette, A.; Kim, K.; Shresta, S. Identification of Zika virus epitopes reveals immunodominant and protective roles for dengue virus cross-reactive CD8⁺ T cells. *Nat. Microbiol.* **2017**, *2*, 17036. [[CrossRef](#)] [[PubMed](#)]
91. Reynolds, C.J.; Sulyeman, O.M.; Ortega-Prieto, A.M.; Skelton, J.K.; Bonnesoeur, P.; Blohm, A.; Carregaro, V.; Silva, J.S.; James, E.A.; Maillère, B.; et al. T cell immunity to Zika virus targets immunodominant epitopes that show cross-reactivity with other Flaviviruses. *Sci. Rep.* **2018**, *8*, 672. [[CrossRef](#)] [[PubMed](#)]
92. Zust, R.; Toh, Y.-X.; Valdés, I.; Cerny, D.; Heinrich, J.; Hermida, L.; Marcos, E.; Guillén, G.; Kalinke, U.; Shi, P.-Y.; et al. Type I Interferon signals in macrophages and dendritic cells control dengue virus infection: Implications for a new mouse model to test dengue vaccines. *J. Virol.* **2014**, *88*, 7276–7285. [[CrossRef](#)] [[PubMed](#)]
93. Elong Ngono, A.; Vizcarra, E.A.; Tang, W.W.; Sheets, N.; Joo, Y.; Kim, K.; Gorman, M.J.; Diamond, M.S.; Shresta, S. Mapping and role of the CD8⁺ T cell response during primary Zika virus infection in mice. *Cell Host Microbe* **2017**, *21*, 35–46. [[CrossRef](#)] [[PubMed](#)]
94. Wen, J.; Elong Ngono, A.; Regla-Nava, J.A.; Kim, K.; Gorman, M.J.; Diamond, M.S.; Shresta, S. Dengue virus-reactive CD8⁺ T cells mediate cross-protection against subsequent Zika virus challenge. *Nat. Commun.* **2017**, *8*, 1459. [[CrossRef](#)] [[PubMed](#)]

95. Larocca, R.A.; Abbink, P.; Peron, J.P.S.; Zanotto, P.M.A.; Iampietro, M.J.; Badamchi-Zadeh, A.; Boyd, M.; Ng'ang'a, D.; Kirilova, M.; Nityanandam, R.; et al. Vaccine protection against Zika virus from Brazil. *Nature* **2016**, *536*, 474–478. [[CrossRef](#)] [[PubMed](#)]
96. Abbink, P.; Larocca, R.A.; de la Barrera, R.A.; Bricault, C.A.; Moseley, E.T.; Boyd, M.; Kirilova, M.; Li, Z.; Ng'ang'a, D.; Nanayakkara, O.; et al. Protective efficacy of multiple vaccine platforms against Zika virus challenge in rhesus monkeys. *Science* **2016**, *353*, 1129–1132. [[CrossRef](#)] [[PubMed](#)]
97. Lima, N.S.; Rolland, M.; Modjarrad, K.; Trautmann, L. T Cell immunity and Zika virus vaccine development. *Trends Immunol.* **2017**, *38*, 594–605. [[CrossRef](#)] [[PubMed](#)]
98. Huang, H.; Li, S.; Zhang, Y.; Han, X.; Jia, B.; Liu, H.; Liu, D.; Tan, S.; Wang, Q.; Bi, Y.; et al. CD8⁺ T cell immune response in immunocompetent mice during Zika virus infection. *J. Virol.* **2017**, *91*. [[CrossRef](#)] [[PubMed](#)]
99. Pardy, R.D.; Rajah, M.M.; Condotta, S.A.; Taylor, N.G.; Sagan, S.M.; Richer, M.J. Analysis of the T cell response to Zika virus and identification of a novel CD8⁺ T cell epitope in immunocompetent mice. *PLoS Pathog.* **2017**, *13*, e1006184. [[CrossRef](#)] [[PubMed](#)]
100. Winkler, C.W.; Myers, L.M.; Woods, T.A.; Messer, R.J.; Carmody, A.B.; McNally, K.L.; Scott, D.P.; Hasenkrug, K.J.; Best, S.M.; Peterson, K.E. Adaptive immune responses to Zika virus are important for controlling virus infection and preventing infection in brain and testes. *J. Immunol.* **2017**, *198*, 3526–3535. [[CrossRef](#)] [[PubMed](#)]
101. Naiyer, M.M.; Cassidy, S.A.; Magri, A.; Cowton, V.; Chen, K.; Mansour, S.; Kranidioti, C.; Mbiribindi, B.; Rettman, P.; Harris, S.; et al. KIR2DS2 recognizes conserved peptides derived from viral helicases in the context of HLA-C. *Sci. Immunol.* **2017**, *2*. [[CrossRef](#)] [[PubMed](#)]
102. Grifoni, A.; Pham, J.; Sidney, J.; O'Rourke, P.; Paul, S.; Peters, B.; Martini, S.R.; de Silva, A.D.; Ricciardi, M.J.; Magnani, D.M.; et al. Prior Dengue virus exposure shapes T cell immunity to Zika virus in humans. *J. Virol.* **2017**, *91*. [[CrossRef](#)] [[PubMed](#)]
103. Halstead, S.B. Which Dengue Vaccine Approach Is the Most Promising, and Should We Be Concerned about Enhanced Disease after Vaccination? There Is Only One True Winner. *C.S.H. Perspect. Biol.* **2017**. [[CrossRef](#)] [[PubMed](#)]
104. Zellweger, R.M.; Eddy, W.E.; Tang, W.W.; Miller, R.; Shresta, S. CD8⁺ T cells prevent antigen-induced antibody-dependent enhancement of dengue disease in mice. *J. Immunol.* **2014**, *193*, 4117–4124. [[CrossRef](#)] [[PubMed](#)]
105. Zellweger, R.M.; Tang, W.W.; Eddy, W.E.; King, K.; Sanchez, M.C.; Shresta, S. CD8⁺ T cells can mediate short-term protection against heterotypic dengue virus reinfection in mice. *J. Virol.* **2015**, *89*, 6494–6505. [[CrossRef](#)] [[PubMed](#)]

