



HAL
open science

Improving dengue virus capture rates in humans and vectors in Kamphaeng Phet Province, Thailand, using an enhanced spatiotemporal surveillance strategy

Stephen J. Thomas, Jared Aldstadt, Richard G. Jarman, Darunee Buddhari, In-Kyu Yoon, Jason H. Richardson, Alongkot Ponlawat, Sapon Iamsirithaworn, Thomas W. Scott, Alan L. Rothman, et al.

► To cite this version:

Stephen J. Thomas, Jared Aldstadt, Richard G. Jarman, Darunee Buddhari, In-Kyu Yoon, et al.. Improving dengue virus capture rates in humans and vectors in Kamphaeng Phet Province, Thailand, using an enhanced spatiotemporal surveillance strategy. *American Journal of Tropical Medicine and Hygiene*, 2015, 93 (1), pp.24-32. 10.4269/ajtmh.14-0242 . pasteur-02011017

HAL Id: pasteur-02011017

<https://pasteur.hal.science/pasteur-02011017>

Submitted on 8 Feb 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

1 LRH: THOMAS AND OTHERS

2 RRH: ENHANCED SURVEILLANCE FOR DENGUE

3

4 Improving Dengue Virus Capture Rates in Humans and Vectors in Kamphaeng
5 Phet Province, Thailand, Using an Enhanced Spatiotemporal Surveillance Strategy

6 Stephen J. Thomas ^{1*}, Jared Aldstadt ³, Richard G. Jarman ^{1*}, Darunee Buddhari ², In-Kyu Yoon
7 ², Jason H. Richardson ⁴, Alongkot Ponlawat ⁴, Sopon Iamsirithaworn ⁵, Thomas W. Scott ^{6,10}

8 Alan L. Rothman,⁷ Robert V. Gibbons,² Louis Lambrechts,⁸ and Timothy P. Endy⁹

9 ¹Viral Diseases Branch, Walter Reed Army Institute of Research, Silver Spring, Maryland,
10 U.S.A.

11 ²Department of Virology, United States Army Medical Component-Armed Forces Research
12 Institute of Medical Sciences, Bangkok, Thailand

13 ³Department of Geography, University at Buffalo, Buffalo, New York, U.S.A.

14 ⁴Department of Entomology, Armed Forces Research Institute of Medical Sciences, 315/6
15 Rajvithi Rd., Bangkok, Thailand

16 ⁵Bureau of Epidemiology, Department of Disease Control Sciences, Ministry of Public Health,
17 Nonthaburi, Thailand

18 ⁶Department of Entomology, University of California, Davis, Davis, California, U.S.A.

19 ⁷Institute for Immunology and Informatics, University of Rhode Island, Providence, Rhode Island

20 ⁸Insect-Virus Interactions Group, Department of Genomes and Genetics, Institut Pasteur, Centre
21 National de la Recherche Scientifique, Paris, France

22 ⁹Department of Infectious Diseases, State University of New York, Syracuse, New York, U.S.A.

23 ¹⁰Fogarty International Center, National Institutes of Health, Bethesda, Maryland 20892, U.S.A.

24 *Work was completed while at the Department of Virology, United States Army Medical
25 Component-Armed Forces Research Institute of Medical Sciences, Bangkok, Thailand
26 Corresponding Author: Stephen J. Thomas, MD
27 Mailing Address: Viral Diseases Branch, Walter Reed Army Institute of Research, 503 Robert
28 Grant Ave., Silver Spring, MD, 20910
29 Tel: 301-319-9210
30 Fax: 301-319-9661
31 E-mail: Stephen.j.thomas3.mil@mail.mil

32

33

34 **Abstract:**

35 Dengue is of public health importance in tropical and sub-tropical regions. DENV transmission
36 dynamics was studied in Kamphaeng Phet Province, Thailand using an enhanced spatiotemporal
37 surveillance of 93 hospitalized subjects with confirmed dengue (initiates) and associated cluster
38 individuals (associates) with entomologic sampling. A total of 438 associates were enrolled from
39 208 houses with household members with fever history located within a 200 meter radius of an
40 initiate case. Of 409 associates, 86 (21%) had laboratory-confirmed DENV infection. A total of
41 63 (1.8%) of the 3,565 mosquitoes collected were dengue PCR+. There was a significant
42 relationship between spatial proximity to the initiate case and likelihood of detecting DENV
43 from associate cases and *Aedes* mosquitoes. The viral detection rate from human hosts and
44 mosquito vectors in this study was higher than previously observed by the study team in the
45 same geographic area using different methodologies. We propose that the sampling strategy used
46 in this study could support surveillance of DENV transmission and vector interactions.

47

48 Abstract Word Count: 150

49 Manuscript Word Count: 4,279

50 **Introduction**

51 Dengue illness is a disease of increasing public health importance [1]. Available data and
52 modeling estimate that there are 390 million dengue virus (DENV) infections annually with 96
53 million manifesting clinically [2]. International travel, population growth, increasing
54 urbanization, and a changing global ecology foster an increasingly favorable environment for the
55 expanding dengue endemic areas and the peridomestic *Aedes aegypti* which transmit the viruses
56 [3]. There are currently no licensed drugs or vaccines to treat or prevent dengue. When applied
57 properly, vector control and personal protective measures have successfully disrupted epidemic
58 and endemic DENV transmission [4,5]. Unfortunately, successful vector control programs have
59 been the exception and difficult to sustain [6]. The strategic use of safe and efficacious dengue
60 vaccines in combination with appropriately targeted and sustained vector control measures is
61 increasingly being considered as the optimal approach to produce a sustained reduction in
62 dengue's global burden [7].

63 Once a vaccine is available, numerous questions will remain about how to most
64 effectively target and co-implement vaccination and vector control programs. The prospect of
65 implementing large-scale, control programs raises a number of questions:

- 66 1) How will vaccination and vector control affect the complex, dynamic, and evolving
67 interactions between vector, virus, and host occurring at the macro (i.e., country or
68 region) and micro (i.e., province, district, village or neighborhood) spatial scales?
- 69 2) How will vaccination and vector control affect the complex, dynamic, and evolving
70 interactions between vector, virus, and host at the population (i.e., *Aedes* species, DENV
71 serotypes and genotypes, and people of various ethnic backgrounds) level?

72 3) How will “herd immunity” be affected and how will this influence DENV evolution at
73 the micro and macro population levels and the associated observed clinical phenotypes?

74 4) What effect, if any, will existing herd immunity (due to vaccination or natural infection)
75 to non-dengue flaviviruses (e.g. Japanese encephalitis and yellow fever viruses) have on
76 DENV transmission and the observed clinical phenotypes following infection?

77 The overarching study objective was to explore DENV transmission dynamics and virus-
78 vector-host interactions prior to, during, and following the introduction of dengue vaccines into
79 central Thailand. The authors pursued this objective by building upon observations from previous
80 prospective studies such as the focality of DENV transmission, presence or history of fever
81 increasing the likelihood of isolating virus from a household, and the significance of year-round
82 DENV transmission [8-10]. Study methods were modified in an effort to maximize DENV
83 isolation rates from human hosts and mosquito vectors and further explore earlier observations,
84 which included:

85 1) Enrolling initiate and associate cases throughout the year (i.e., high dengue season and
86 low dengue season) to explore trends in seasonal and spatial DENV transmission;

87 2) Only enrolling DENV PCR+ initiate cases to increase the likelihood of capturing active
88 transmission;

89 3) Only enrolling associates with fever or a history of fever within the last 7 days or sharing
90 a household with someone meeting these criteria to increase the likelihood of identifying
91 associates with recent infection;

92 4) Reassessing associates for the occurrence of fever between the acute and convalescent
93 blood collection to capture additional viremic cases;

- 94 5) Expanding the age of enrollment to include children above the age of 6 months and adults
95 to improve understanding of transmission inside the home; and
96 6) Extending the enrollment of associates and mosquito collection from a 100 to 200m
97 (meter) radius around the initiate.

98 In this report the authors describe initiation of the baseline phase (i.e. prior to vaccine
99 introduction) and include detailed accounts of study methodology and the entomologic, clinical,
100 epidemiologic, virologic, serologic, and molecular characterization of human cases captured by
101 active and passive surveillance methods between November 2009 and December 2010.

102

103

104 **Methods:**

105 Ethics Statement

106 The study protocol was approved by the Institutional Review Boards (IRB) of the Thai
107 Ministry of Public Health (MOPH), Walter Reed Army Institute of Research (WRAIR), and the
108 State University of New York (SUNY), Upstate Medical University. The IRB's of the University
109 of California, Davis (UCD), University of Rhode Island (URI) and University at Buffalo
110 established relying agreements with WRAIR IRB.

111 All study subjects engaged in documented informed consent or assent process, as
112 applicable, prior to participating in any study activities. In the event the subject was unable to
113 participate in the informed consent/assent process, a recognized health care proxy represented
114 them in the process and documented consent. From this point forward, when the authors discuss
115 consent, assent is also implied as applicable.

116 Role of the Funding Source

117 Funding sources for this project included National Institutes of Health grants R01
118 GM083224-01 and P01 AI034533. Additional funding was provided by the US Military
119 Infectious Diseases Research Program. The funding sources had no involvement in study design,
120 data collection, analysis or interpretation, report writing, or publication submission. The
121 corresponding author had full access to all study data and final responsibility for the decision for
122 publication.

123 Study Location

124 The study was conducted in Kamphaeng Phet (KPP) province in north-central Thailand.
125 There were 725,846 registered residents of KPP in 2009 (Thailand, Department of Provincial
126 Administration, 2010). The KPP Provincial Hospital (KPPPH) is located in the province's

127 central district. The Armed Forces Research Institute of Medical Sciences (AFRIMS),
128 Department of Virology field site (KPP AFRIMS Virology Research Unit [KAVRU]) is located
129 on the KPPPH grounds. The Department of Entomology, AFRIMS field site is located a short
130 distance from KPPPH [11].

131 Demographics of house residents were collected and house spatial coordinates were
132 identified using a Geographic Positioning System hand held unit (Trimble(r) GeoXH(tm),
133 GeoExplorer(r) 2008 series, Trimble Navigation Limited, CO, USA) and geo-coded into a
134 Geographic Information System (GIS) database (CorporationArcMap™, version 9.1, ESRI, CA,
135 USA).

136 Study Definitions

137 The authors recognize the potential confusion using terms such as, “index case,”
138 “contacts,” and “cluster investigations.” As the index case may not be the true first infection in
139 space and time and the contact may not be a true infection resulting from DENV transmission
140 from the initiating case we have attempted to more accurately define the relationships
141 investigated in this study without making claims of causation. The hospitalized dengue cases
142 serving as the initiator of community transmission investigations are referred to as the “initiating
143 case,” or “initiate”. The enrolled individuals residing within the initiate’s home or within 200
144 meters of the initiate are referred to as “associates” and if found to have DENV infection, as
145 “associate cases”. The combination of the initiate, associate and associate cases is referred to as
146 a “spatiotemporal group.”

147 Symptomatic DENV infection in either initiating or associated cases are defined as any
148 febrile illness (reported or measured fever) paired with a confirmatory molecular or serologic
149 assay run on the acute or acute and convalescent blood sample pair, respectively. Subclinical

150 DENV infections are defined as associated cases (all initiating cases were symptomatic and
151 hospitalized) with a positive molecular or serologic result, but without reported or measured fever
152 during the period between the acute and convalescent blood sample collection (0 to 14 days).

153 Subjects are serologically classified as having an “acute” DENV infection if their acute
154 and/or convalescent blood sample pair (or day 0 and/or day 14 blood samples in associates) was
155 dengue IgM positive; or, if IgM is negative, IgG is positive with rising titer. Subjects are
156 classified as “recent” DENV infection if IgM was negative and IgG was positive with declining
157 titer. These are further categorized as “primary” infection if the IgM to IgG ratio is $\geq 1 \cdot 8$ and
158 “secondary” infection if the ratio is $< 1 \cdot 8$ [12].

159 Initiate Case Identification and Evaluation

160 KAVRU provides the KPPPH dengue diagnostic research assays for patients presenting
161 with fever or history of fever and dengue-like symptoms as outlined in the World Health
162 Organization (WHO) guidelines for the diagnosis and treatment of dengue [13]. Hospital staff
163 members identified suspected dengue cases, completed and documented the informed consent
164 process allowing the testing of a blood sample, and then sent the sample to KAVRU for RT-PCR
165 testing. Virus RNA was extracted from human serum or mosquito suspension using Qiagen Viral
166 RNA Extraction kits. Serotype-specific DNA fragments from each unknown sample were
167 amplified by TaqDNA polymerase through RT-PCR performed at KAVRU following
168 modifications to the Lanciotti protocol [14]. Suspected cases > 6 months of age, that provided a
169 sample collected within the prior 24 hours, and had detectable DENV RNA by RT-PCR were
170 provided the opportunity for study enrollment as initiate cases. Following acquisition of informed
171 consent, the acute blood sample was accessed and served as the baseline sample. Demographic
172 and clinical laboratory information was collected. Initiate case house spatial coordinates were

173 recorded and geo-coded into a GIS database. Mosquitoes inside and outside of the initiate home
174 were collected by aspiration and processed as described in preparation for DENV RNA detection
175 RT-PCR [10]. In approximately 14 days a second blood sample was collected from the initiating
176 case for serologic testing by in-house IgM/IgG EIA.

177 Associate Case Identification and Evaluation

178 Residents >6 months of age sharing the initiate household as their primary residence were
179 provided the opportunity for study enrollment. Residents previously enrolled as an associate
180 within the past 6 months were excluded in an attempt to improve geographic diversity and the
181 likelihood of isolating DENV (i.e. reduce chances of recent infection and lingering homo- or
182 heterotypic immunity). Following the informed consent process, clinical information and a blood
183 sample were collected. In approximately 14 days a second blood sample was collected for
184 serologic testing. General well-being of the associates was tracked by active surveillance (i.e.,
185 combination of self-reporting and outreach by staff via phone or daily home visit by village
186 health worker over the 14 day period). If, between the day of enrollment and the day of
187 convalescent blood sampling, the associate develops fever, a second acute blood sample was
188 collected and the 14 day “clock” started again. Therefore, it was possible an associate might have
189 two acute samples, one triggered as an associate of the initiating case (acute sample #1) and one
190 triggered by the development of illness (acute sample #2), as well as one convalescent sample.

191 Individuals not residing within the same house as the initiate case, but living in a home
192 within a 200m radius of the initiate case were also considered for enrollment as associates. If
193 anyone within the home reported an active fever or history of fever (temperature $\geq 38^{\circ}\text{C}$) within
194 the past 7 days, all residents of the home >6 months of age were eligible for enrollment.

195 Following the informed consent process, residents had clinical and demographic information and
196 an acute blood sample collected. Associates were followed and blood collected as detailed above.

197 House Mapping, Associate Case Home Identification, and Entomologic Sampling

198 GPS mapped initiate households were used to construct a digital map, enabling the team
199 to precisely identify houses located within a 200m radius of the initiate case [10,15]. Study
200 nurses visited households starting closest to the initiate house and moving in sequential fashion
201 to the periphery of the area and back to the initiate house, then back out again along a different
202 line until the circle was complete or 25 eligible associates were enrolled, whichever occurred
203 first (Figure 1).

204 On day 1 of each initiate/associate case investigation, adult *Ae aegypti* were collected
205 using standard backpack aspirators from inside and within the immediate vicinity of each
206 potential associate's house. The end of each aspirator tube was fitted with a 1-pint cardboard
207 cage. After completing the collection for each home, the cage was labeled and stored on dry ice
208 for transportation to the laboratory where the chilled mosquitoes were examined, speciated, and
209 processed for DENV detection. A thorough adult aspiration collection usually requires ~10-15
210 minutes per house. It was estimated that approximately 25% of the adult *Ae. aegypti* were
211 captured in a single pass through the house (TW Scott, unpublished data). All specimens were
212 identified to species by entomology field supervisors to ensure speciation accuracy and quality
213 control. As previously mentioned, mosquitoes inside and outside of the initiate home were
214 collected by aspiration and processed as described with serotype-specific DENV RNA detection
215 RT-PCR performed on each individual mosquito [10].

216 Statistical Analyses

217 Data were analyzed using SPSS (SPSS for Windows version 19) and R (The R Project for
218 Statistical Computing 2.12). Demographic, clinical, and laboratory parameters were analyzed at
219 the initiate, associate, and house levels. Student's *t*-test and analysis of variance (ANOVA) were
220 used to test for differences in continuous variables. Fisher's exact test and Pearson's chi-squared
221 were used to examine associations between categorical variables.

222 **Results**

223 Enrollment of Initiates and Associates

224 Figure 2 depicts enrollment of initiates and associates and results of serologic and
225 molecular testing for DENV infection. Approximately 49% of patients hospitalized with
226 suspected dengue were PCR+ and of those 62% were enrolled in the study. Of the 4,345
227 households within a 200m radius of the 93 initiate homes, 124 (2.8%) had someone with active
228 fever or history of fever; 115 of these households had volunteers who consented to enroll. Initiate
229 and associate households contained 793 individuals meeting enrollment criteria, and 438 (55%)
230 consented to enroll as associates. Complete serologic data was available on 93% of the enrolled
231 associates and of these 21% had a positive dengue serology. Of the 86 associates with positive
232 serology, 42% also were PCR+. All PCR+ associates had fever at the time of enrollment. Eleven
233 associates who enrolled without fever developed fever following their first acute blood sampling;
234 of these 8 were serology positive and 7 PCR+.

235 Of note logistic limitations guided enrollment on days with heavy dengue patient census
236 accounting for the difference between the 149 PCR+ cases meeting inclusion criteria and the 93
237 ultimately enrolled. In these instances the study team randomly selected which initiate cases
238 would lead to an associate case investigation using a random numbers table.

239 Characteristics of the Initiating Cases

240 Each of the 93 initiate cases was a hospitalized acute, PCR+ DENV infection (Table 1).
241 Infections with DENV-2 were most numerous followed by DENV-3 and then DENV-1; there
242 were no DENV-4 infections. Most cases were acute secondary DENV infections (85%).
243 Approximately half of all cases were DF (48%) and half DHF (52%); there were no deaths among
244 the initiates.

245 Associated Cases Available for Enrollment

246 The number of associate houses within 200m of an initiating case ranged from 1-232 with
247 a mean of 47.7 (SD 42 houses). The number of households enrolled ranged from 1-9 with a mean
248 of 2.2 (SD 1.5 households). For each spatiotemporal group there was a range of associates
249 enrolled from 0-18 people with a mean of 4.7 (SD 4.0 people).

250 Characteristics of the Associated Cases

251 The majority (79%) of 409 enrolled associates had no serologic evidence of infection,
252 20% had evidence of an acute DENV infection, and there was 1 case with evidence of JE
253 infection (IgM positive, no encephalitis; Table 1). There was complete concordance between the
254 DENV serotype of the initiating case and the DENV serotypes detected in associate cases in that
255 spatiotemporal unit. Nested PCR results among associates revealed that DENV-3 was the most
256 common infection followed by DENV-2 and then DENV-1.

257 There was a significant difference in the reporting of nausea between primary and
258 secondary DENV infections and significant variation in reporting of headache, rhinorrhea, cough,
259 and retro-orbital pain among the age groups (Tables 2 and 3). There was a statistically significant
260 difference in the probability of an associate experiencing a DENV infection based on the DENV
261 type infecting the initiate. DENV-3 infection in the initiate carried the highest probability of
262 associate infection (Table 4).

263 Spatial Distribution of Associate Case Households

264 Closer proximity of a household to the initiate was correlated with an increased
265 associate household enrollment (Table 5). Households further from the initiate household
266 had a lower rate of DENV infection among associate household residents (Table 6). This
267 pattern of transmission was not gradual with distance. In an associate from a household with
268 fever for example, the rate of infection was similarly high (around 0.3) within 120 meters
269 and very low in households with fever beyond 120 meters. This would indicate that
270 spatiotemporal transmission in this investigation was primarily limited to 120 meters and
271 extending the radius from 100 to 200 meters did not substantially increase the efficiency of
272 detecting dengue cases or viremia.

273 DENV Infection in *Ae. aegypti* Collected in the Homes of Human Dengue Cases

274 A total of 3,565 *Ae. aegypti* were collected from 4,438 households (i.e. all households).
275 A total of 233 (6.5%) mosquitoes were from initiate households, of which 23 (9.9%) were
276 DENV PCR+. There was a greater likelihood one or more collected mosquitoes was DENV
277 PCR+ within the initiate house or a house closer to the initiate's house (Table 7).
278 There were 162 initiate households with at least one associate resident who had complete
279 serology and 212 associate households with at least one associate with complete serology. Of the
280 initiate households, female *Ae. aegypti* were collected in 54.9% and 58.5% of the associate
281 households had female *Ae. aegypti* collected. Of the female mosquitoes collected in initiate
282 households 29% of households had a DENV PCR+ mosquito while 6.5% of associate
283 households had a DENV PCR+ mosquito. There was a higher likelihood of finding a DENV
284 PCR+ mosquito in an initiate household (Table 8). Of the 31 initiate households with a DENV
285 serology and an associate resident, 54.8% of the houses had female *Ae. aegypti* captured and

286 64.7% of these households had a DENV PCR+ mosquito. Of the 42 associate households with a
287 DENV serology + associate, 57.1% of households had a female *Ae. aegypti* mosquito and one
288 (4.1%) of these households had a DENV PCR+ mosquito. Despite a similar likelihood of
289 finding female *Ae. aegypti* in households with a DENV+ serology resident, there was a higher
290 likelihood of finding DENV PCR+ mosquitoes in the initiate household (Table 8). In total, there
291 were 46 households with a DENV PCR+ mosquito and an associate residing with complete
292 serology; 80.4% were initiate households (Table 8.). There was a high degree of concordance
293 between the isolated DENV serotypes from mosquitoes within a cluster and the infecting
294 serotype of the index case (Table 9). In year 2010, only one cluster (10-077) was found
295 discordant from the mosquito (DENV-2) as compared to the index case (DENV-1). In 2012,
296 one cluster (12-025) was found DENV-1 in index case while PCR results were DENV-2 and
297 DENV-4 in 2 mosquito samples. The rest of the clusters were concordant.

298

299 **Discussion**

300 Our current study further characterizes the complex transmission dynamics and virus-
301 vector-host interactions in a well characterized spatial area around an infected viremic inpatient
302 in Central Thailand. We demonstrated a significant relationship between spatial proximity to the
303 initiate case and likelihood of detecting DENV from associate cases and *Ae aegypti* with higher
304 than anticipated virus detection from both human hosts and mosquito vectors. We propose that
305 the sampling strategy described is valuable for ongoing surveillance of DENV transmission
306 during and after field studies and the introduction of dengue vaccines.

307 The design of this study was built upon observations from previous prospective cohort
308 studies of DENV transmission conducted with and without initiate case and associate

309 (contact/cluster) investigations [8-10,16,17]. Modifications were made to the study design in an
310 attempt to increase the detection of virus, symptomatic and subclinical associates of initiate cases,
311 and infected mosquito vectors. The result was a demographically diverse group of DENV
312 infected people representing a broad virologic, serologic, and clinical spectrum. Substantial virus
313 detection rates were observed in both human associate cases and mosquitoes.

314 The age range of initiate cases was surprisingly wide (2.6-56 years), and the mean age
315 was higher than expected at 18.7 years [18]. This observation is consistent with unpublished data
316 from the authors across additional dengue seasons and data from other sources. For example, in
317 2010, the Thailand MOPH reported that the highest case rate for DHF was in the 10-14 year age
318 group, and that the 5-9 and 15-24 year age groups had the second and third highest rates,
319 respectively (Thailand, Ministry of Public Health). These findings are in contrast to decades of
320 data where the mean age of hospitalized dengue was in the age range of 5-9 years [19-24]. One
321 explanation for the shift is that smaller birth cohorts had reduced number of susceptibles, thereby
322 impacting the force of infection and the time in a person's life when they acquire their first and
323 second DENV infection, the latter being more often associated with clinically significant disease
324 in Thailand [18].

325 We also enrolled associates across a wide age range (7 months – 94.2 years) with a mean
326 age almost double the initiate cases (31.4 years vs. 18.7 years). Serologic evidence of DENV
327 infection and symptomatic DENV infection was observed in a number of subjects aged 40 years
328 and older. Most of these were secondary (i.e., post-primary) DENV infections, but we also
329 detected primary DENV infections, defined serologically, in the 50-59 and 85+ year age groups.
330 Dengue occurs throughout the year with all four serotypes circulating with high rates of infection
331 and transmission (hyperendemic) in KPP. Pediatric cohort studies have demonstrated high

332 DENV infection attack rates (combined symptomatic and subclinical) in the range of 2.2% to
333 7.9% per year (average incidence 5.8%) over at least the preceding 10 years [9]. Based on these
334 observations, it was assumed that lifelong KPP residents have experienced multiple DENV
335 infections by their late 20's. A modified plaque reduction neutralization assay (single dilution
336 neutralization assay, 1:30 dilution, 70% viral plaque reduction) completed on a cohort of children
337 enrolled in a KPP prospective study from 1998 to 2002 revealed a gradually increasing
338 prevalence of neutralizing antibodies to at least one DENV type from 45% among 4 year olds to
339 91% among 13 year olds (Yoon, IK, unpublished data) [10,16,25]. Our finding that all PCR+
340 associate cases had fever at the time of enrollment may also underscore the increasing infection
341 burden in adults as they are more likely to experience symptomatic DENV infections compared to
342 primary infection in children. Our observation of symptomatic dengue in older individuals merits
343 reconsideration of traditional views on the force of infection in hyperendemic areas and the
344 durability of homotypic and heterotypic immunity.

345 There was complete concordance between the infecting DENV serotypes in the initiate
346 case and the corresponding associate cases. This finding supports the concept that DENV
347 transmission is focal and serotype-conserved in space and time.[17,26] The authors'
348 acknowledge that the assumption that each initiating case represents the true index (i.e., first)
349 infection in each spatiotemporal group may be incorrect. Because most DENV infections are
350 asymptomatic or not clinically severe enough to drive health care-seeking behavior it is possible
351 the initiating case simply represents the first clinically overt infection in that defined geographic
352 area.

353 The ratio of symptomatic to subclinical infection among primary infections was 1:0.2; the
354 ratio in secondary infections was 1:0.4 and similar (1:0.4) among the 86 serologically positive

355 associates. These ratios are consistent with previous work by our group and others, which
356 reported symptomatic to subclinical ratios ranging from 1:0.9 to 1:18 [9,10,27-30]. The
357 differences in results are most likely associated with the study's surveillance focus in and around
358 symptomatic, PCR+ initiating cases. Previous studies conducted routine biologic sampling to
359 capture subclinical cases and cluster studies exploring houses with or without active or a recent
360 history of fever [9].

361 The probability of an associate becoming infected was associated with a number of factors
362 [31-33]. As mentioned above, a significant association existed between the DENV serotype
363 infecting the initiating case and the proportion of associates subsequently infected with the same
364 DENV serotype (p-value <0.001). Associates in samples initiated by a DENV-3 infection had a
365 30% chance of being serologically positive. The likelihood of infection was 19% for associates
366 in DENV-2 samples and 9% in DENV-1 samples. Taken by itself, this may point to viral
367 properties allowing for more efficient transmission, such as higher titer viremia in the human host
368 or mosquito vector, a longer duration of viremia increasing the potential window for transmission
369 to vectors and / or shorter incubation period in the mosquito. Another significant association was
370 between the distance a potential associate lived from an initiating case and the likelihood of
371 becoming infected (negative correlation, p-value <0.001). It is reasonable to assume the factors
372 culminating in an initiate infection (i.e., convergence of susceptible host and infected vector in
373 space and time) would also drive efficiency of transmission and associate infections until a
374 geographic barrier was introduced (i.e. next susceptible beyond the dispersal distance of the
375 vector) or other factors limited transmission (i.e., protective herd immunity of associates). We
376 did not define the relative roles of humans and mosquitoes in moving virus from one house to

377 another within clusters, although in general, humans tend to move more often and greater
378 distances than *Ae. Aegypti* [31-33].

379 Approximately 29% of all households had female *Ae. aegypti* and 1.8% of the females
380 collected were infected with DENV. Although initiating case houses and associate houses with a
381 PCR+ contact had similar rates of female *Ae. aegypti* infestation (55% and 59%, respectively),
382 the percentage of PCR+ mosquitoes was higher in the initiate case households (9.9%) compared
383 to non-initiate households with and without a DENV positive contact, 1.2% and 1.1%
384 respectively. There was a significant association (p-value < 0.001) between the distance of a
385 home from the initiating case and the proportion of PCR+ mosquitoes found within the house.
386 These observations are concordant with previous reports and intuitive; once a clinical case of
387 dengue is identified there is a high likelihood of finding infected vector(s) co-residing in the same
388 geographic location as the ill human.

389 Initiate cases represented the full spectrum of DENV infections clinically, serologically,
390 and virologically. We consider the finding of more than 1 in 5 associates of an initiating case
391 having serologic and, in some cases, clinical evidence of an acute DENV infection significant
392 from an epidemiologic and transmission dynamics standpoint. These results represent the initial
393 phase of a multi-year prospective study. The study design employed improved the efficiency of
394 capturing DENV in associates and vector populations compared to previous efforts by focusing
395 surveillance in areas and households where DENV transmission was most likely to be occurring.
396 Our observations emphasize the focal spread of DENV and its spatial restriction associated with
397 the mosquito vector and suggests that initiate cases were important in this study (ill and PCR+) in
398 transmitting DENV to mosquitoes and thus to others living near the initiate. Previous studies
399 support the observation that patient DENV viremia was a marker of human infectiousness and

400 blood meals containing high concentrations of DENV were positively associated with the
401 prevalence of infectious mosquitoes.[34] Our results and study design may be useful when
402 designing experiments to study fine scale patterns of DENV transmission and to increase
403 detection of case contacts with inapparent infections.

404 **Acknowledgements**

405 This research would not have been possible without outstanding support of the staff of the
406 Kamphaeng Phet Provincial Hospital and the excellent research capabilities of those at the
407 Kamphaeng Phet AFRIMS Virology Research Unit. The authors specifically acknowledge the
408 efforts of Chaleaw Saengchan and Rungkarn Hangsuwan for clinical research coordinator
409 services, and Udom Kijchalao and Arissara Pongsiri for their work in the field investigations. We
410 thank Dr. Angkana Uppapong and Dr. Kamchai Rungsimunpaiboon for expert medical
411 consultation and leadership and Dr. Nuttaporn Wongsutthipakorn for the support of the
412 provincial health office. The authors are especially grateful to the citizens of Kamphaeng Phet
413 for their willingness to participate in this study and their support of over three decades of
414 biomedical research in their community.

415 **Conflicts of Interest**

416 None of the authors on this manuscript have a conflict of interest with the manuscript and data
417 presented.

418 **Disclaimer**

419 The opinions or assertions contained herein are the private views of the authors and are not to be
420 construed as reflecting the official views of the United States Army, the United States
421 Department of Defense, or the National Institutes of Health.

422 **Key words:** dengue; transmission; viremia, vector, flavivirus, arbovirus, epidemiology
423 **Abbreviations:** Analysis of variance, ANOVA; Armed Forces Research Institute of Medical
424 Sciences, AFRIMS; Dengue virus DENV; Dengue hemorrhagic fever, DHF; Dengue shock
425 syndrome, DSS; Deoxyribonucleic acid, DNA; Enzyme-linked immunosorbent assay, ELISA;
426 Institutional Review Boards, IRB; Kamphaeng Phet, KPP; Kamphaeng Phet AFRIMS Virology
427 Research Unit, KAVRU; Kamphaeng Phet Provincial Hospital, KPPPH; Polymerase chain
428 reaction, PCR; Reverse transcriptase PCR, RT-PCR; Ribonucleic acid, RNA; State University of
429 New York Upstate Medical Center, SUNY; Thai Ministry of Public Health, MOPH; University
430 of Massachusetts Medical School, UMMS; University of California, Davis, UCD; Walter Reed
431 Army Institute of Research, WRAIR; World Health Organization, WHO;
432

433 **Author Contributions:**

434 SJT- study design, study execution, data analysis, manuscript writing

435 JA- data analysis, manuscript writing

436 DT- study design, study execution, manuscript writing

437 IKY - study design, study execution, laboratory assays, data analysis, manuscript writing.

438 JR- study execution, entomologic data analysis, manuscript writing

439 AP- study execution, entomologic data analysis, manuscript writing

440 SI- - study design, study execution, data analysis, manuscript writing

441 TWS- - study design, study execution, data analysis, manuscript writing

442 ALR- study design, data interpretation, and editing manuscript

443 RVG – study design, study planning, interpreting results and editing manuscript

444 LL – study design, data interpretation, and editing manuscript


445 TPE- - study design, execution, data analysis, manuscript writing

446

447

448 **Tables and Figures:**

449 Figure 1. Method of identifying and enrolling associates around an initiating case. The clinical
450 research team would begin at the initiate's home, identify a collection of homes within 200
451 meters of the initiate's home, identify homes with an occupant having active fever or history of
452 fever within the past 7 days, and enroll associates only from those homes while bypassing homes
453 without active fever or history of fever. The entomology research team, meanwhile, collected
454 mosquitoes inside and outside the home from all homes within a 200-meter radius of the initiating
455 case home regardless of fever history. Once investigations and enrollment of associates were
456 completed in one grouping, the teams would return to the initiating case's home and then identify,
457 using the aerial map, the next grouping of homes moving in a clockwise manner.




458 Circling home has no active fever or history of fever and is therefore not included in the investigation of associates.

459 Table 1. Initiate and Associate Group Demographic Data and Infection Characterization.

		Initiate (N = 93)	Associate		
			Available Serology (N = 409)	Available Serology + Fever (N = 156)	Dengue + Serology (N = 86)
Age (years)					
	minimum	2.6	0.58	0.58	0.83
	maximum	56.0	94.2	77.0	82.0
	mean	18.7	31.4	20.1	23.1
Sex (%)					
	Female	48	54	57	55
	Male	52	46	43	45
Infecting DENV type (% of total typed cases)		Total case count = 93	Acute sample not available, no clinical symptoms.		Total case count = 86

	DENV-1	16			1
	DENV-2	60			16
	DENV-3	24			24
	DENV-4	0			0
	Not detected				58
Serology Results (% of total results)					
	Acute Primary	2	3	6	16
	Acute Secondary	85	17	25	80
	Recent Secondary	2	1	1	3
	No Serologic Diagnosis	8	79	67	
	JE Infection		0.2		

460 Figure 2. Flow Diagram of Initiate and Associate Enrollment and Testing Outcome



461

462

463 Table 2. Symptom Complex in Primary vs. Secondary Associate Infections

Symptom	Primary (N=14)	Secondary (N=72)	p-value
Fever	11 (0.79)	50 (0.69)	0.749
Headache	5 (0.36)	37 (0.51)	0.384
Rhinorrhea	3 (0.21)	16 (0.22)	1.000
Anorexia	5 (0.36)	27 (0.38)	1.000
Cough	4 (0.29)	27 (0.38)	0.762
Nausea	0 (0.00)	25 (0.35)	0.008
Drowsiness	0 (0.00)	12 (0.17)	0.202
Muscle or Joint Pain	3 (0.21)	36 (0.50)	0.077
Abdominal Pain	2 (0.14)	15 (0.21)	0.727
Retro-orbital Pain	3 (0.21)	22 (0.31)	0.749
Rash	4 (0.29)	10 (0.14)	0.231
Diarrhea	2 (0.14)	14 (0.19)	1.000
Bleeding	0 (0.00)	5 (0.07)	0.586
Hospitalized	1 (0.07)	6 (0.05)	1.000

464 Table 3. Clinical Spectrum of EIA Positive Associates by Age Groups

Symptom	0-9 Years (N=22)	10-19 Years (N=31)	20-29 Years (N=10)	>30 Years (N=23)	p-value
Fever	18 (0.82)	24 (0.77)	8 (0.80)	11 (0.48)	0.052
Headache	6 (0.27)	20 (0.65)	7 (0.70)	9 (0.39)	0.021
Rhinorrhea	8 (0.36)	7 (0.23)	3 (0.30)	1 (0.04)	0.043
Anorexia	9 (0.41)	11 (0.35)	6 (0.60)	6 (0.26)	0.325
Cough	13 (0.59)	11 (0.35)	4 (0.40)	3 (0.13)	0.012
Nausea	4 (0.18)	12 (0.39)	5 (0.50)	4 (0.17)	0.101
Drowsiness	2 (0.09)	7 (0.23)	1 (0.10)	2 (0.09)	0.519
Muscle or Joint Pain	7 (0.32)	16 (0.52)	7 (0.70)	9 (0.39)	0.189
Abdominal Pain	3 (0.14)	8 (0.26)	2 (0.20)	4 (0.17)	0.741
Retro-orbital Pain	1 (0.05)	15 (0.48)	4 (0.40)	5 (0.22)	0.002
Rash	5 (0.23)	6 (0.19)	2 (0.20)	1 (0.04)	0.282
Diarrhea	2 (0.09)	8 (0.26)	3 (0.30)	3 (0.13)	0.285
Bleeding	2 (0.09)	2 (0.06)	1 (0.10)	0 (0.00)	0.440
Hospitalized	1 (0.05)	3 (0.10)	2 (0.20)	1 (0.04)	0.432

465 Table 4. Probability of Dengue Virus Infection in Associates According to the Initiating Case
 466 Infecting DENV Serotype.

467

Serologic Diagnosis	Infecting DENV of Initiate		
	DENV-1	DENV-2	DENV-3
EIA Negative	43	191	88
EIA Positive	4 (0.09)	44 (0.19)	38 (0.30)

468 Footnote: Numbers in parentheses are the proportions EIA positive. The infecting DENV type is
 469 taken from the initiating case. The differences are significant (Fisher's exact test, p-value =
 470 0.003).

471 Table 5. Spatial Distribution of Households and Enrolled Households

472

	Distance from Initiate Household (meters)				
	>0-40	>40-80	>80-120	>120-160	>160-200
Total	539	814	971	1051	1063
Households					
Enrolled	103	53	32	13	7
Households	(0.191)	(0.065)	(0.033)	(0.012)	(0.007)

473 Footnote: Numbers in parentheses indicate proportions of houses enrolled for each distance

474 category. Differences are significant, p-value < 0.001 by Fisher's exact test –.

475 Table 6. Spatial Distribution of Associates and Associates with DENV Infection by Serology

	Initiate Household	Distance from Initiate Household				
		>0-40	>40-80	>80-120	>120-160	>160-200
Total Associates with Available Serology	187	41	67	51	38	25
EIA Positive Associates	42 (0.225)	14 (0.341)	12 (0.179)	16 (0.314)	0 (0.000)	2 (0.080)

476 Footnote: Numbers in parentheses indicate proportions of associates that were EIA positive
 477 within the initiate house and for each distance category. Differences are significant, p-value <
 478 0.001 by Fisher's exact test.

479

480 Table 7. Spatial Distribution of Female *Ae. aegypti* within Houses and PCR+ Mosquitoes within
 481 Houses
 482

	Initiate Household	Distance to Initiate Household				
		>0-40	>40-80	>80-120	>120-160	>160-200
Total Mosquitoes	233	328	545	733	745	981
PCR+ Mosquitoes	23 (0.099)	10 (0.030)	7 (0.013)	13 (0.018)	7 (0.009)	3 (0.003)

483 Footnote: Numbers in parentheses indicate proportions of mosquitoes that were PCR+ for each
 484 distance category. Differences are significant, p-value < 0.001 by Fisher's exact test.
 485

486 Table 8. Associate Infection Rates by Infection Status of Female *Ae. aegypti* Captured in
 487 Households.

488

	Initiate Houses			Associate Houses		
	PCR Positive Female <i>Ae. aegypti</i>	Females Captured <i>Ae. aegypti</i>	No Females Captured <i>Ae. aegypti</i>	PCR Positive Females <i>Ae. aegypti</i>	Females Captured <i>Ae. aegypti</i>	No Females Captured <i>Ae. aegypti</i>
Total Associates with Available Serology	26	88	71	8	125	90
Serology Positive Associates	11 (0.423)	17 (0.193)	14 (0.197)	1 (0.125)	25 (0.200)	18 (0.200)

489 Footnote: Numbers in parentheses indicate proportions of associates that were serology positive
 490 within the initiate house and for each category.

491

492 Table Nine. Isolated DENV serotypes from clusters.

493

Year	2009-2010	2009-2012
Number of Ae. Samples	3545	9322
DENV-1	5	13
DENV-2	36	87
DENV-3	22	28
DENV-4	0	3

494

495 Year 2010, one cluster (10-077) was found discordant mosquito result (DENV-2) from the index
496 case (DENV-1)

497

498 Year 2012, one cluster (12-025) was found DENV-1 in index case while PCR result were
499 DENV-2 and DENV-4 in 2 mosquito samples.

500

501

502

503

504

505

506

507

508

509

510

511

512

513

514

515

516

517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541

References

1. Morens DM, Fauci AS (2008) Dengue and hemorrhagic fever: a potential threat to public health in the United States. *JAMA* 299: 214-216.
2. Bhatt S, Gething PW, Brady OJ, Messina JP, Farlow AW, et al. (2013) The global distribution and burden of dengue. *Nature* 496: 504-507.
3. Diaz FJ, Farfan-Ale JA, Olson KE, Lorono-Pino MA, Gubler DJ, et al. (2002) Genetic variation within the premembrane coding region of dengue viruses from the Yucatan peninsula of Mexico. *Am J Trop Med Hyg* 67: 93-101.
4. Ooi EE, Goh KT, Gubler DJ (2006) Dengue prevention and 35 years of vector control in Singapore. *Emerg Infect Dis* 12: 887-893.
5. Goh KT (1997) Dengue--a re-emerging infectious disease in Singapore. *Ann Acad Med Singapore* 26: 664-670.
6. Morrison AC, Zielinski-Gutierrez E, Scott TW, Rosenberg R (2008) Defining challenges and proposing solutions for control of the virus vector *Aedes aegypti*. *PLoS medicine* 5: e68.
7. Thomas SJ (2011) The necessity and quandaries of dengue vaccine development. *J Infect Dis* 203: 299-303.
8. Endy TP, Nisalak A, Chunsuttiwat S, Libraty DH, Green S, et al. (2002) Spatial and temporal circulation of dengue virus serotypes: a prospective study of primary school children in Kamphaeng Phet, Thailand. *Am J Epidemiol* 156: 52-59.

- 542 9. Endy TP, Chunsuttiwat S, Nisalak A, Libraty DH, Green S, et al. (2002) Epidemiology of inapparent and
543 symptomatic acute dengue virus infection: a prospective study of primary school children in Kamphaeng
544 Phet, Thailand. *Am J Epidemiol* 156: 40-51.
- 545 10. Mammen MP, Pimgate C, Koenraadt CJ, Rothman AL, Aldstadt J, et al. (2008) Spatial and temporal clustering of
546 dengue virus transmission in Thai villages. *PLoS Med* 5: e205.
- 547 11. Gibbons RV, Nisalak A, Yoon IK, Tannitisupawong D, Rungsimunpaiboon K, et al. (2013) A model international
548 partnership for community-based research on vaccine-preventable diseases: the Kamphaeng Phet-
549 AFRIMS Virology Research Unit (KAVRU). *Vaccine* 31: 4487-4500.
- 550 12. Innis BL, Nisalak A, Nimmannitya S, Kusalerdchariya S, Chongswasdi V, et al. (1989) An enzyme-linked
551 immunosorbent assay to characterize dengue infections where dengue and Japanese encephalitis co-
552 circulate. *Am J Trop Med Hyg* 40: 418-427.
- 553 13. Anonymous (1997) *Dengue haemorrhagic fever: diagnosis, treatment, prevention and control*, 2nd edition.
554 Geneva: World Health Organization. 84 p.
- 555 14. Klungthong C, Gibbons RV, Thaisomboonsuk B, Nisalak A, Kalayanarooj S, et al. (2007) Dengue virus detection
556 using whole blood for reverse transcriptase PCR and virus isolation. *J Clin Microbiol* 45: 2480-2485.
- 557 15. Paul A, Harrington LC, Zhang L, Scott JG (2005) Insecticide resistance in *Culex pipiens* from New York. *J Am*
558 *Mosq Control Assoc* 21: 305-309.
- 559 16. Yoon IK, Getis A, Aldstadt J, Rothman AL, Tannitisupawong D, et al. (2012) Fine scale spatiotemporal clustering
560 of dengue virus transmission in children and *Aedes aegypti* in rural Thai villages. *PLoS Negl Trop Dis* 6:
561 e1730.
- 562 17. Jarman RG, Holmes EC, Rodpradit P, Klungthong C, Gibbons RV, et al. (2008) Microevolution of Dengue viruses
563 circulating among primary school children in Kamphaeng Phet, Thailand. *J Virol* 82: 5494-5500.
- 564 18. Cummings DA, Iamsirithaworn S, Lessler JT, McDermott A, Prasanthong R, et al. (2009) The impact of the
565 demographic transition on dengue in Thailand: insights from a statistical analysis and mathematical
566 modeling. *PLoS medicine* 6: e1000139.
- 567 19. Vaughn DW, Green S, Kalayanarooj S, Innis BL, Nimmannitya S, et al. The dengue hemorrhagic fever project:
568 viremia and antibody responses early in dengue; 1997 14 -17 April; Papeete, Tahiti.

569 20. Vaughn DW, Green S, Kalayanarooj S, Innis BL, Nimmannitya S, et al. (1997) Dengue in the early febrile phase:
570 viremia and antibody responses. *J Infect Dis* 176: 322-330.

571 21. Vaughn DW, Green S, Kalayanarooj S, Innis BL, Nimmannitya S, et al. Antibody response patterns and disease
572 severity by dengue virus serotype; 1997 7-11 December; Orlando, FL.

573 22. Vaughn DW, Green S, Kalayanarooj S, Innis BL, Nimmannitya S, et al. (2000) Dengue viremia titer, antibody
574 response pattern, and virus serotype correlate with disease severity. *J Infect Dis* 181: 2-9.

575 23. Nimmannitya S (1987) Clinical spectrum and management of dengue haemorrhagic fever. *Southeast Asian*
576 *J Trop Med public Health* 18: 392-397.

577 24. Nimmannitya S (1987) Dengue haemorrhagic fever in Thailand. *Southeast Asian J Trop Med public Health* 18:
578 291-294.

579 25. Yoon IK, Rothman AL, Tannitisupawong D, Srikiatkachorn A, Jarman RG, et al. (2012) Underrecognized mildly
580 symptomatic viremic dengue virus infections in rural Thai schools and villages. *J Infect Dis* 206: 389-398.

581 26. Rabaa MA, Klungthong C, Yoon IK, Holmes EC, Chinnawirotpisan P, et al. (2013) Frequent in-migration and
582 highly focal transmission of dengue viruses among children in Kamphaeng Phet, Thailand. *PLoS Negl Trop*
583 *Dis* 7: e1990.

584 27. Burke DS, Nisalak A, Johnson DE, Scott RM (1988) A prospective study of dengue infections in Bangkok.
585 *Am J Trop Med Hyg* 38: 172-180.

586 28. Balmaseda A, Standish K, Mercado JC, Matute JC, Tellez Y, et al. (2010) Trends in patterns of dengue
587 transmission over 4 years in a pediatric cohort study in Nicaragua. *J Infect Dis* 201: 5-14.

588 29. Porter KR, Beckett CG, Kosasih H, Tan RI, Alisjahbana B, et al. (2005) Epidemiology of dengue and dengue
589 hemorrhagic fever in a cohort of adults living in Bandung, West Java, Indonesia. *Am J Trop Med Hyg* 72:
590 60-66.

591 30. Endy TP, Yoon IK, Mammen MP (2010) Prospective cohort studies of dengue viral transmission and severity of
592 disease. *Curr Top Microbiol Immunol* 338: 1-13.

593 31. Harrington LC, Scott TW, Lerdthusnee K, Coleman RC, Costero A, et al. (2005) Dispersal of the dengue vector
594 *Aedes aegypti* within and between rural communities. *Am J Trop Med Hyg* 72: 209-220.

- 595 32. Stoddard ST, Morrison AC, Vazquez-Prokopec GM, Paz Soldan V, Kochel TJ, et al. (2009) The role of human
596 movement in the transmission of vector-borne pathogens. *PLoS neglected tropical diseases* 3: e481.
- 597 33. Stoddard ST, Forshey BM, Morrison AC, Paz-Soldan VA, Vazquez-Prokopec GM, et al. (2013) House-to-house
598 human movement drives dengue virus transmission. *Proc Natl Acad Sci U S A* 110: 994-999.
- 599 34. Nguyet MN, Duong TH, Trung VT, Nguyen TH, Tran CN, et al. (2013) Host and viral features of human dengue
600 cases shape the population of infected and infectious *Aedes aegypti* mosquitoes. *Proc Natl Acad Sci U S A*
601 110: 9072-9077.
- 602