

Corynebacterium rouxii sp. nov., a novel member of the diphtheriae species complex

Edgar Badell, Melanie Hennart, Carla Rodrigues, Virginie Passet, Mélody Dazas, Valerie Bouchez, Leonardo Panunzi, Annick Carmi-Leroy, Julie Toubiana, Sylvain Brisse

► **To cite this version:**

Edgar Badell, Melanie Hennart, Carla Rodrigues, Virginie Passet, Mélody Dazas, et al.. *Corynebacterium rouxii* sp. nov., a novel member of the diphtheriae species complex. *Research in Microbiology*, Elsevier, 2020, 171 (3-4), pp.122-127. 10.1016/j.resmic.2020.02.003 . pasteur-02862963v2

HAL Id: pasteur-02862963

<https://hal-pasteur.archives-ouvertes.fr/pasteur-02862963v2>

Submitted on 18 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





Original Article

Corynebacterium rouxii sp. nov., a novel member of the *diphtheriae* species complex

Edgar Badell ^{a, b, 1}, Mélanie Hennart ^{a, 1}, Carla Rodrigues ^{a, 1}, Virginie Passet ^a,
Melody Dazas ^a, Leonardo Panunzi ^a, Valérie Bouchez ^a, Annick Carmi–Leroy ^{a, b},
Julie Toubiana ^{a, b, c}, Sylvain Brisse ^{a, b, *}

^a Institut Pasteur, Biodiversity and Epidemiology of Bacterial Pathogens, Paris, France

^b National Reference Center for the *Corynebacteria* of the *diphtheriae* complex, Paris, France

^c Université de Paris, Department of General Paediatrics and Infectious Diseases, Necker-Enfants malades University Hospital, AP-HP, Paris, France

ARTICLE INFO

Article history:

Received 1 January 2020

Accepted 19 February 2020

Available online 28 February 2020

Keywords:

Diphtheria

Taxonomy

Phylogeny

Genome sequencing

MALDI-TOF

ABSTRACT

A group of six clinical isolates previously identified as *Corynebacterium diphtheriae* biovar Belfantii, isolated from human cutaneous or peritoneum infections and from one dog, were characterized by genomic sequencing, biochemical analysis and MALDI-TOF mass spectrometry. The six isolates were negative for the diphtheria toxin gene. Phylogenetic analyses showed that the six isolates (including FRC0190^T) are clearly demarcated from *C. diphtheriae*, *Corynebacterium belfantii*, *Corynebacterium ulcerans* and *Corynebacterium pseudotuberculosis*. The average nucleotide identity of FRC0190^T with *C. diphtheriae* NCTC11397^T was 92.6%, and was 91.8% with *C. belfantii* FRC0043^T. *C. diphtheriae* subsp. *lausannense* strain CHUV2995^T appeared to be a later heterotypic synonym of *C. belfantii* (ANI, 99.3%). Phenotyping data revealed an atypical negative or heterogeneous intermediate maltose fermentation reaction for the six isolates. MALDI-TOF mass spectrometry differentiated the new group from the other *Corynebacterium* taxa by the presence of specific spectral peaks. *rpoB* sequences showed identity to atypical, maltose-negative *C. diphtheriae* biovar Belfantii isolates previously described from two cats in the USA. We propose the name *Corynebacterium rouxii* sp. nov. for the novel group, with FRC0190^T (= CIP 111752^T = DSM 110354^T) as type strain.

© 2020 The Authors. Published by Elsevier Masson SAS on behalf of Institut Pasteur. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

The genus *Corynebacterium* currently includes approximately 111 species [1–3]. The most important human pathogen of the genus is *Corynebacterium diphtheriae*, which causes diphtheria [2,4]. *C. diphtheriae* is genetically heterogeneous [5–8] and four biovars were defined: Gravis, Mitis, Belfantii and Intermedius [9–11], the latter being almost never reported in recent literature. In 2010, maltose-non fermenting strains of *C. diphtheriae* biovar Belfantii were reported from two cats in the USA, and were shown to have a divergent *rpoB* sequence [12]. In 2018, some biovar

Belfantii isolates were classified as a novel species, *C. belfantii* [3], with 94.85% average nucleotide identity (ANI) with *C. diphtheriae*. Almost simultaneously, *C. diphtheriae* subsp. *lausannense* was also proposed for strains of biovar Belfantii [13]. The *tox* gene, which codes for diphtheria toxin, is carried on a coryneophage that can lysogenize strains of *C. diphtheriae*. However, the *tox* gene was rarely reported in isolates of biovar Belfantii [5,14,15] and no strain of *Corynebacterium belfantii* or *C. diphtheriae* subsp. *lausannense* was described as *tox*-positive [3,12]. The *tox* gene can also be harboured by strains of *Corynebacterium ulcerans* and *Corynebacterium pseudotuberculosis*, two species that are phylogenetically close to *C. diphtheriae* and *C. belfantii* [16]. Together, the above-mentioned species constitute a single phylogenetic clade nested within the *Corynebacterium* genus. We refer to this clade as the *C. diphtheriae* complex.

Here, we define the taxonomic status of six isolates initially identified as *C. diphtheriae* biovar Belfantii, isolated from five human infections and one dog in France.

Abbreviations: ANI, Average nucleotide identity; MALDI-TOF, Matrix-assisted laser desorption/ionisation time-of-flight.

* Corresponding author. National Reference Center for the *Corynebacteria* of the *diphtheriae* complex, Paris, France.

E-mail address: sbrisse@pasteur.fr (S. Brisse).

¹ These authors contributed equally to this work

<https://doi.org/10.1016/j.resmic.2020.02.003>

0923-2508/© 2020 The Authors. Published by Elsevier Masson SAS on behalf of Institut Pasteur. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

2. Material and methods

We compared the six atypical clinical isolates, among which is strain FRC0190^T, with 13 *C. diphtheriae* strains of biovars Gravis or Mitis (including *C. diphtheriae* type strain NCTC 11397^T) and 8 strains previously [3] identified as *C. belfantii* (including the type strain FRC0043^T; Table 1; Table S1). Type strains of *C. ulcerans* (CIP 106504^T = NCTC 7910^T) and of *C. pseudotuberculosis* (CIP 102968^T = ATCC 19410^T) were also included for comparison.

Clinical samples or isolates were received at the French National Reference Centre for Corynebacteria of the *diphtheriae* complex for isolation and/or characterization, respectively. Oxoid's Tinsdale agar with supplement medium (Thermo Fisher Diagnostics, Dardilly, France) was used to isolate *C. diphtheriae* from clinical samples. Isolates were frozen in Brain-Heart-Infusion (BHI) medium containing 30% of glycerol and stored at –80 °C prior to this study. After thawing, isolates were grown at 37 °C on tryptose-casein soy agar plates during 24 h. DNA was extracted from a few colonies with the DNeasy Blood and Tissue kit (Qiagen, Hilden, Germany). The six isolates were identified as *C. diphtheriae* by multiplex polymerase chain reaction (PCR) combining a *dtxR* gene fragment specific for *C. diphtheriae* [15] and a multiplex PCR [17,18] that targets a fragment of the *pld* gene specific for *C. pseudotuberculosis*, the gene *rpoB* (amplified in all species of the *C. diphtheriae* complex) and a fragment of 16S rRNA gene specific for *C. pseudotuberculosis* and *C. ulcerans*. The *tox* gene was also detected by PCR [19]. These PCR results were confirmed using a more recent four-plex qPCR [20].

For biochemical identification, standard methods were used [14,21,22]. More specifically, strains were characterized for pyrazinamidase, urease, nitrate reductase and for utilization of maltose and trehalose using API Coryne strips (BioMérieux, Marcy l'Etoile, France) and the Rosco Diagnostica reagents (Eurobio, Les Ulis, France) following provider's recommendations. The Hiss serum water test was used for glycogen fermentation. Briefly, this test was performed as follows. Solution A was obtained by dissolving 500 mg of bacteriological peptone (Oxoid, Hampshire, UK; ref. LP0037) in 100 mL of distilled water, adding 100 mg of Na₂HPO₄ (Sigma–Aldrich, Saint-Louis, Missouri, USA; ref: S7907), and homogenizing and heating the mixture until boiling. After cooling to room temperature, 18 mL of sterile horse serum were added and mixed. In parallel, solution B was prepared by adding 430 mg of acid fuchsin (Sigma–Aldrich, Saint-Louis, Missouri, USA, ref: F8129) into 86 mL of distilled water, after which 14 mL of 30% NaOH (ThermoFisher Scientific, Waltham, Massachusetts; ref. S/4950/PB15) were added. Solution B was stored up to 15 days in the dark. To prepare the complete Hiss serum water sugar medium, 780 µL of solution B were added to the total volume of solution A. The pH was adjusted to 7.7 using HCl 5N (Sigma–Aldrich, Saint-Louis, Missouri, USA; ref. H1758). 100 mg of glycogen (Acros Organics, Geel, Belgium, ref: 422,950,050) were then added. The solution was mixed and distributed in 3.5 mL aliquots in 5 mL glass tubes, and sterilized at 108 °C during 30 min. This medium was conserved up to 6 months at 5 °C + 3 °C. To perform the glycogen test, a loopful (10 µL) of a bacterial culture from Columbia blood agar or Tryptose-Casein-Soy agar was introduced into a tube containing 3.5 mL of sterile Hiss serum water sugar medium. Results were read manually after homogenization of the suspension and incubation at 37 °C ± 2 °C during 24 h. Strains NCTC 12077 and NCTC 764 were used as positive and negative controls, respectively (expected results: dark pink and light pink, respectively).

The biovar of isolates was determined based on the combination of nitrate reductase (positive in Mitis and Gravis, negative in Belfantii) and glycogen fermentation (positive in Gravis only).

Antimicrobial susceptibility was characterized by the disk diffusion method using impregnated paper disks (Bio-Rad, Marnes-la-Coquette, France) and minimum inhibitory concentrations were determined using ETEST strips (BioMérieux, Marcy l'Etoile, France). The sensitivity was interpreted using CA-SFM/EUCAST V.1.0 (Jan 2019) criteria for *Corynebacterium* (https://www.sfm-microbiologie.org/wp-content/uploads/2019/02/CASFM2019_V1.0.pdf). Susceptibility was tested for the following antimicrobial agents: fosfomicin, vancomycin, kanamycin, gentamycin, penicillin G, oxacillin, amoxicillin, imipenem, cefotaxime, clindamycin, azithromycin, spiramycin, clarithromycin, erythromycin, ciprofloxacin, trimethoprim-sulfamethoxazole, trimethoprim, sulfonamide, pristinamycin, rifampicin and tetracycline.

MALDI-TOF mass spectrometry was used for identification confirmation. For this purpose, an overnight culture on Tryptone-Casein-Soy Agar (TSA) (37 °C) was used to prepare the samples accordingly to the ethanol/formic acid extraction procedure proposed by the manufacturer (Bruker Daltonics, Bremen, Germany). The cell extracts were then spotted onto an MBT Biotarget 96 target plate, air dried and overlaid with 1 µL of a saturated α -cyano-4-hydroxycinnamic acid (HCCA). 24 mass spectra per strain were acquired on a Microflex LT mass spectrometer (Bruker Daltonics, Bremen, Germany). Re-analysis of the spectra was performed for the purpose of this work. Spectra were first preprocessed by applying smoothing and baseline subtraction with FlexAnalysis software using default parameters, exported as text files from the Bruker system and then imported and analyzed in a dedicated BioNumerics v7.6.3 (Applied-Maths, Belgium) database following the protocol described by Rodrigues et al. [23]. To allocate proteins to the specific peaks detected, we extracted all the molecular weights from the genomes of the type strains (NCTC11397^T, FRC0043^T and FRC0190^T) using a Biopython script (<https://biopython.org/DIST/docs/api/Bio.SeqUtils-module.html>) and performed sequence alignments with ClustalW for the candidate proteins.

Genomic sequencing was performed from Nextera XT libraries using a NextSeq-500 instrument (Illumina, San Diego, USA) with a 2 × 150 nt paired-end protocol. Contig sequences were assembled using SPAdes v3.12.0 [24] (Table S1). JSpeciesWS [25] was used to calculate the BLAST-based average nucleotide identity (ANI_b). BLASTN was used to extract 16S rRNA and *rpoB* sequences from genome assemblies and to determine the presence or absence of the *narI/HJK* nitrate reduction gene cluster using as query the cluster of strain NCTC 13129 (RefSeq accession number: DIP_RS13820 to DIP_RS13845) [26]. *rpoB* and 16S rRNA gene sequences of atypical *C. belfantii* strains from cats [12] were included for comparison. For genome-based phylogenetic analysis, the pairwise *p*-distance (*i.e.*, proportion of aligned nucleotide differences) between each pair of genomes was estimated based on Mash [27] using a multiple hit correction [28] with JolyTree (<https://gitlab.pasteur.fr/GIPhy/JolyTree>). For 16S rRNA and *rpoB* gene sequences, sequences were aligned with MAFFT v7.407 [29] and the resulting alignment was used for phylogenetic tree inference with IQ-TREE v1.6.7.2 [30] using the GTR + I + G4 model. Branch support was obtained after 1000 bootstrap replicates.

3. Results and discussion

Six isolates were isolated from five cutaneous lesions and one ascitic fluid sample (Table 1). Strikingly, human cutaneous lesions were all ulcerations due to underlying chronic arteritis. Ascitic fluid was sampled on a patient with a suspicion of spontaneous peritonitis. The dog was investigated in the context of purulent orbital cellulitis.

Table 1
Strains used in this study and their characteristics.

Isolate ^b	Species	biovar ^a	Isolation year	Country	Geographic origin ^c	tox gene	Isolation source	Disease	Reference
FRC0190 ^T	<i>C. rouxii</i>	Belfanti	2013	France	Lot, Cahors	Negative	Cutaneous	Foot ulceration, chronic arteritis	This study
FRC0071	<i>C. rouxii</i>	Belfanti	2011	France	Haute-Garonne, Toulouse	Negative	Cutaneous	Leg ulceration on chronic arteritis - diabetes	This study
FRC0284	<i>C. rouxii</i>	Belfanti	2015	France	Rhone, Lyon	Negative	Cutaneous	Limb amputation - vasculitis	This study
FRC0297	<i>C. rouxii</i>	Belfanti	2015	France	Herault, Beziers	Negative	Ascitic fluid	Spontaneous peritonitis	This study
FRC0412	<i>C. rouxii</i>	Belfanti	2016	France	Lot, Cahors	Negative	Cutaneous	Purulent orbital cellulitis (dog)	This study
FRC0527	<i>C. rouxii</i>	Belfanti	2017	France	Savoie, Chambéry	Negative	Cutaneous	Foot ulceration on chronic arteritis	This study
FRC0043 ^T	<i>C. belfantii</i>	Belfanti	2009	France	Corrèze, Brives	Negative	Pharyngeal membrane	Laryngitis	Dazas et al., 2018 IJSEM
06–4305	<i>C. belfantii</i>	Belfanti	2006	France	Rhone, Lyon	Negative	Expectoration	Bronchopathy	Dazas et al., 2018 IJSEM
00–0744	<i>C. belfantii</i>	Belfanti	2000	France	Calvados, Caen	Negative	Expectoration	Cystic fibrosis	Dazas et al., 2018 IJSEM
FRC0074	<i>C. belfantii</i>	Belfanti	2011	France	Cote d'Or, Dijon	Negative	Expectoration	Cystic fibrosis	Dazas et al., 2018 IJSEM
FRC0223	<i>C. belfantii</i>	Belfanti	2014	France	Pas-de-Calais, Coquelles	Negative	Sinusitis swab	Sinusitis	Dazas et al., 2018 IJSEM
05–3187	<i>C. belfantii</i>	Belfanti	2005	France	Seine-Maritime, Rouen	Negative	Nasal swab	Rhinitis	Dazas et al., 2018 IJSEM
FRC0250	<i>C. belfantii</i>	Belfanti	2014	France	Bas-Rhin, Strasbourg	Negative	Bronchoalveolar wash	Pneumonia	Dazas et al., 2018 IJSEM
FRC0301	<i>C. belfantii</i>	Belfanti	2015	France	Calvados, Lisieux	Negative	Expectoration	n.a.	Dazas et al., 2018 IJSEM
NCTC 11397 ^T	<i>C. diphtheriae</i>	Gravis	1969	USA	New York, USA	Negative	n.a.	n.a.	Dazas et al., 2018 IJSEM
NCTC 13129	<i>C. diphtheriae</i>	Gravis	1997	United Kingdom	Unknown	Positive	Pharyngeal membrane	Diphtheria	Dazas et al., 2018 IJSEM
FRC0336	<i>C. diphtheriae</i>	Gravis	2015	France	Ille-et-Vilaine, Rennes	Positive	Cutaneous	Leishmaniasis	Dazas et al., 2018 IJSEM
FRC0304	<i>C. diphtheriae</i>	Gravis	2015	France	La Reunion, St Denis	Negative	Cutaneous	Bullous skin lesion	Dazas et al., 2018 IJSEM
FRC0375	<i>C. diphtheriae</i>	Mitis	2015	France	Oise, Creil	Positive	Cutaneous	Ankle ulceration	Dazas et al., 2018 IJSEM
FRC0432	<i>C. diphtheriae</i>	Mitis	2016	France	Seine-et-Marne, Vaires sur Marne	Negative	Cutaneous	Purulent scalp skin injury	Dazas et al., 2018 IJSEM
FRC0157	<i>C. diphtheriae</i>	Mitis	2013	France	Paris	Negative	Cutaneous	Left ankle wound	Dazas et al., 2018 IJSEM
FRC0132	<i>C. diphtheriae</i>	Mitis	2012	France	Yvelines, Le Chesnay (return from Mali)	Negative	Cutaneous	Necrotic lesions	Dazas et al., 2018 IJSEM
FRC0036	<i>C. diphtheriae</i>	Mitis	2009	France	Mayotte, Mamoudzou	Negative	Cutaneous	Burn wound	Dazas et al., 2018 IJSEM
FRC0154	<i>C. diphtheriae</i>	Mitis	2012	France	Haut-Rhin, Colmar	Positive	Cutaneous	Cutaneous infection	Dazas et al., 2018 IJSEM
FRC0049	<i>C. diphtheriae</i>	Mitis	2009	France	Mayotte, Mamoudzou	Positive	Cutaneous	Genital lesion	Dazas et al., 2018 IJSEM
FRC0430	<i>C. diphtheriae</i>	Mitis	2016	France	Rhone, Bron	Positive	Cutaneous	Leg ulcerations	Dazas et al., 2018 IJSEM
FRC0436	<i>C. diphtheriae</i>	Mitis	2016	France	Ille-et-Vilaine, Rennes	Positive	Cutaneous	Cutaneous infection	Dazas et al., 2018 IJSEM
ATCC 19410 ^T	<i>C. pseudotuberculosis</i>	not applicable	1931	n.a.	South America	Negative	Infected gland (sheep)	n.a.	PMID: 13882624 (Cummins, 1962)
NCTC 7910 ^T	<i>C. ulcerans</i>	not applicable	1948	United Kingdom	n.a.	Negative	Throat	n.a.	PMID: 7,729,671 (Riegel et al., 1995)

n.a.: not available.

^a Biovar of *C. diphtheriae* as classically defined.

^b FRC: collection of the French National Reference Center for the Corynebacteria of the *C. diphtheriae* complex; NCTC: National Collection of Type Cultures (Public Health England); ATCC: American Type Culture Collection.

^c Geographic origin for French isolates is given as "French Department, city".

The six isolates were *tox* negative (Table S1); more specifically, they were negative for amplification of the expected 910-bp PCR product encompassing fragments A and B of the toxin gene [19] and also negative for the amplification of a 117-bp region of diphtheria toxin fragment A [31] by multiplex qPCR [20]. We also confirmed by BLASTN that the *tox* gene sequence (query: *tox* gene sequence from strain NCTC 13129, RefSeq accession number: DIP_RS12515) was absent from the genomic assemblies. After species identification by multiplex PCR, the isolates were positive for *dtxR* and *rpoB* and negative for *C. ulcerans/C. pseudotuberculosis* 16S rDNA and *pld*, leading to initial identification as *C. diphtheriae*. Concordant with this identification, the six isolates were pyrazinamidase, urease and trehalose negative. Upon biotyping, the isolates were nitrate and glycogen negative, a pattern that corresponds to biovar Belfanti. Consistently, the *narKGHIJ* nitrate reduction gene cluster was not detected from the genomic assemblies of these isolates and those of *C. belfantii* (Table S2). The phenotypic aspect of colonies on Tinsdale or blood agar medium was undistinctive from *C. diphtheriae* Mitis and Gravis and *C. belfantii*. However, we noted that similar to the Gravis isolates, the colonies of the six atypical isolates looked dry and were friable on TCS medium. Distinctively, the maltose test was negative for the six isolates using API Coryne (Table S1). The same test was atypical using the Rosco Diagnostic method: results showed heterogeneous coloration that was neither as yellow as the typically positive strains, nor as purple as the negative strains (Fig. S3). This atypical maltose result was not observed using API Coryne strips, with which the maltose test was clearly negative for the six isolates. We noted that the four genes of the maltose

utilization pathway [32] are present and undisrupted in the six isolates, as in other members of the *C. diphtheriae* complex. Further work is required to elucidate the mechanisms of maltose utilization and its regulation, and why the two tests give different results.

Regarding their antimicrobial susceptibility (Table S3), the six isolates were resistant to fosfomycin, as is typical of *Corynebacteria* [33], and were susceptible to all other tested antimicrobial agents with the following exceptions: FRC0284 and FRC0527 were resistant to penicillin (minimum inhibitory concentration: 0.19 mg/L), and FRC0412 was resistant to penicillin and cefotaxime (0.19 mg/L and 1.0 mg/L, respectively).

Genomic sequencing results showed that the six isolates had a genome size of 2.4 Mb on average (Table S1), similar to *C. diphtheriae* biovars Mitis and Gravis isolates (average size: 2.45 Mb), but smaller than *C. belfantii* (average size: 2.7 Mb). A genome sequence-based phylogenetic tree (Fig. 1) revealed three main clades. The first one contained all *C. diphtheriae* Mitis and Gravis isolates, whereas the second comprised all *C. belfantii* isolates, and the third comprised the six maltose-atypical isolates. The mean ANiB value of atypical isolates was 92.4% with the *C. diphtheriae* clade and was 91.4% with *C. belfantii* (Table 2). These data indicate that the six isolates forming the atypical clade correspond to a distinct genomic cluster, separated by a level of nucleotide divergence that is well above the currently accepted genomic species threshold of ~94–96% [34,35]. The atypical clade was genetically homogeneous, with ANiB values among the six isolates ranging from 99.21% to 99.94% (Table 2). Phylogenetic analysis of *rpoB* and 16S rRNA coding sequences was consistent




Fig. 1. Phylogenetic relationships derived from the analysis of genomic sequences. The phylogenetic tree and branch supports were inferred using JolyTree [28] (<https://gitlab.pasteur.fr/GIPhy/JolyTree>). Strains *C. ulcerans* NCTC 7910^T and *C. pseudotuberculosis* ATCC 19410^T were used as outgroup as they are the closest phylogenetic neighbors to *C. diphtheriae*, *C. belfantii* and *C. rouxii*. Branch support is indicated using grey circles (see key; only values > 50 are shown). Each taxonomic type strain is shown in bold; note that *C. diphtheriae* subsp. *lausannense* type strain falls within *C. belfantii*. The scale bar corresponds to an estimated evolutionary distance of 0.01.

Table 2
Average nucleotide identity values.

<i>Corynebacterium</i> species	Strain identifier ^a	FRC0071	FRC0190T	FRC0284	FRC0297	FRC0412	FRC0527	NCTC11397T	FRC0043T	NCTC7910T	ATCC19410T
<i>C. rouxii</i> sp. nov.	FRC0071	100	99.60	99.89	99.21	99.37	99.94	92.30	91.22	71.34	70.94
<i>C. rouxii</i> sp. nov.	FRC0190T	99.68	100	99.75	99.24	99.34	99.68	92.41	91.36	71.40	70.93
<i>C. rouxii</i> sp. nov.	FRC0284	99.94	99.71	100	99.22	99.35	99.92	92.30	91.28	71.26	70.85
<i>C. rouxii</i> sp. nov.	FRC0297	99.26	99.28	99.27	100	99.29	99.26	92.44	91.43	71.16	70.67
<i>C. rouxii</i> sp. nov.	FRC0412	99.45	99.26	99.40	99.23	100	99.44	92.39	91.37	71.04	70.79
<i>C. rouxii</i> sp. nov.	FRC0527	99.95	99.55	99.85	99.21	99.40	100	92.30	91.27	71.27	70.99
<i>C. diphtheriae</i>	NCTC11397T	92.33	92.45	92.30	92.32	92.34	92.32	100	95.07	71.29	70.86
<i>C. belfantii</i>	FRC0043T	90.92	91.06	90.93	90.99	90.99	90.92	94.77	100	71.12	70.76
<i>C. ulcerans</i>	NCTC7910T	71.42	71.51	71.43	71.32	71.30	71.42	71.40	71.31	100	84.33
<i>C. pseudotuberculosis</i>	ATCC19410T	71.31	71.30	71.31	71.25	71.25	71.31	71.19	71.06	84.29	100

^a A trailing T after the strain identifier denotes that the strain is the type strain of its corresponding taxon.

with the distinction of the atypical isolates from *C. diphtheriae* and *C. belfantii* (Figs. S1 and S2). However, the 16S rRNA gene sequence alignment showed only 3 insertions and 4 nucleotide substitutions shared among the six atypical isolates as compared to *C. diphtheriae*, resulting in low resolution of phylogenetic relationships (Fig. S1). We noted that *rpoB* and 16S rRNA sequences of previously reported atypical biovar Belfanti isolates from cats in the USA [12] were indistinguishable from those of the atypical isolates from France, suggesting that the cat isolates from the USA belong to the same novel group. Supporting this observation, the USA cat isolates were also reported as maltose negative [12].

Recently, it was proposed that the *C. diphtheriae* taxon should be subdivided into two subspecies, *C. diphtheriae* subsp. *diphtheriae* and *C. diphtheriae* subsp. *lausannense* [13]. Here, we observed that the ANI value between the type strains of *C. diphtheriae* subsp. *lausannense* and *C. belfantii* was 99.3%. Besides, the former was positioned within the phylogenetic branch of *C. belfantii* (Fig. 1, Figs. S1 and S2), and the descriptions of both taxa are very similar [3,13]. Given that *C. belfantii* was validly published in October 2018, a few months before the taxonomic proposal *C. diphtheriae* subsp. *lausannense* was validated (<https://www.microbiologyresearch.org/content/journal/ijsem/10.1099/ijsem.0.003174>; January 2019), the latter subspecies appears to be a later heterotypic synonym of *C. belfantii*.

Based on the MALDI Biotyper Compass database version 4.1.90 (Bruker Daltonics, Bremen, Germany), the six isolates were identified as *C. diphtheriae*. However, detailed analysis of their spectra led to the identification of six pairs of biomarkers (12 peaks corresponding to the same proteins, with either single and double-charged ion forms) corresponding to three different proteins within the range 3255–9495 *m/z*, which were associated either with the group of six isolates or with *C. diphtheriae* and *C. belfantii* (Fig. S4, Table S4). We presumptively identified the specific biomarkers as two ribosomal proteins, L30 and S20, and one putative stress response protein (CsbD). Consistently, their amino-acid sequences differed between the *C. rouxii* on the one hand, and *C. diphtheriae*/*C. belfantii* on the other hand (Fig. S5). Based on the current dataset, the specificity and sensitivity of peak distribution among the three species ranged between 95–100% and 76–100%, respectively (Table S4). MALDI-TOF MS thus allows the discrimination between *C. rouxii* and *C. diphtheriae*/*C. belfantii*. These results warrant future updates of reference MALDI-TOF databases to incorporate the novel taxon.

Based on the above results, the isolates of the novel clade represent a novel species, which we propose to name *Corynebacterium rouxii*.

Description of *C. rouxii* sp. nov. (rou'. xi.i. N.L. gen. n. *rouxii*, of Roux, a French scientist and former director of Institut Pasteur who made critical contributions to diphtheria toxin discovery and antitoxin treatment).

C. rouxii conforms biochemically to the description of *C. diphtheriae* strains belonging to biovar Belfanti [2,21], except that strains are negative for maltose fermentation (API Coryne), being nearly negative or weakly positive with the Rosco Diagnostica maltose test. Key characteristics that distinguish *C. rouxii* from other members of the *C. diphtheriae* complex are specific MALDI-TOF MS biomarkers as described herein. The G + C content of *C. rouxii* genomes ranges from 53.2% to 53.3%, with a value of 53.3% for the type strain. So far, strains were isolated from 5 humans and a dog in France, as well as from two related cats in the USA.

The type strain is FRC0190^T (= CIP 111752^T = DSM 110354^T), isolated in 2013 from a foot ulceration reported in Cahors, France. The genome accession number of strain FRC0190^T is ERS3795540.

Author contributions

Conceived the study: SB. Performed the experiments: EB, CR, VP, MD, ACL. Analyzed the data: MH, CR, LP, VB, SB. Curated data: VB, MD, EB, JT, SB. Wrote the initial draft of the manuscript: SB, CR, JT. Commented on working versions of the manuscript and agreed on the final version of the manuscript: all.

Funding

This work was supported financially by Institut Pasteur (Paris, France) and by Public Health France (Santé publique France, Saint-Maurice, France).

Declaration of Competing Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Acknowledgments

We thank the “Plateforme de Microbiologie Mutualisée” from Institut Pasteur for genomic sequencing, and Prof. Alain Le Coustumier (Centre hospitalier de Cahors, France) for sending the original culture of strain FRC0190^T.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.resmic.2020.02.003>.

Data availability

Sequence data generated in this study were deposited in the European Nucleotide Archive database and are accessible under

project number PRJEB22103. The EMBL (GenBank/DDBJ) accession numbers of the genomic sequences released in this study are ERS3795539 to ERS3795544. The annotated genomic sequence of strain FRC0190^T was deposited in the European Nucleotide Archive and is available under accession number ERZ1195831. *rpoB* and 16S rRNA gene sequences were also submitted individually under accession numbers MN542347 to MN542352 and MN535982 to MN535987, respectively.

References

- [1] Yu Q-L, Yan Z-F, He X, Tian F-H, Jia C-W, Li C-T. *Corynebacterium defluvii* sp. nov., isolated from Sewage. J Microbiol 2017;55:435–9. <https://doi.org/10.1007/s12275-017-6592-3>.
- [2] Bernard KA, Funke G. Family I. Corynebacteriaceae. Bergey's manual of systematic bacteriology: Volume 5: the actinobacteria. Springer Science & Business Media; 2012.
- [3] Dazas M, Badell E, Carmi-Leroy A, Criscuolo A, Brisse S. Taxonomic status of *Corynebacterium diphtheriae* biovar Belfanti and proposal of *Corynebacterium belfantii* sp. nov. Int J Syst Evol Microbiol 2018;68:3826–31. <https://doi.org/10.1099/ijsem.0.003069>.
- [4] Burkovski A. Diphtheria and its etiological agents. In: Burkovski Andreas, editor. *Corynebacterium diphtheriae* and related toxigenic species. Springer; 2014. p. 1–10 [Chapter 1].
- [5] Bolt F, Cassidy P, Tondella ML, Dezoysa A, Efstratiou A, Sing A, et al. Multi-locus sequence typing identifies evidence for recombination and two distinct lineages of *Corynebacterium diphtheriae*. J Clin Microbiol 2010;48:4177–85. <https://doi.org/10.1128/JCM.00274-10>.
- [6] Trost E, Blom J, de Castro Soares S, Huang I-H, Al-Dilaimi A, Schröder J, et al. Pangenomic study of *Corynebacterium diphtheriae* that provides insights into the genomic diversity of pathogenic isolates from cases of classical diphtheria, endocarditis, and pneumonia. J Bacteriol 2012;194:3199–215. <https://doi.org/10.1128/JB.00183-12>.
- [7] Farfour E, Badell E, Dinu S, Guillot S, Guiso N. Microbiological changes and diversity in autochthonous non-toxicogenic *Corynebacterium diphtheriae* isolated in France. Clin Microbiol Infect 2013;19:980–7. <https://doi.org/10.1111/1469-0691.12103>.
- [8] Sangal V, Hoskisson PA. Evolution, epidemiology and diversity of *Corynebacterium diphtheriae*: new perspectives on an old foe. Infect Genet Evol 2016;43:364–70. <https://doi.org/10.1016/j.meegid.2016.06.024>.
- [9] Thompson JS, Gates-Davis DR, Yong DC. Rapid microbiological identification of *Corynebacterium diphtheriae* and other medically important corynebacteria. J Clin Microbiol 1983;18:926–9.
- [10] Anderson JS, Happold FC, McLeod JW, Thomson JG. On the existence of two forms of diphtheria bacillus—*B. Diphtheriae* Gravis and *B. Diphtheriae* Mitis—and a new medium for their differentiation and for the bacteriological diagnosis of diphtheria. J Pathol 1931;34:667–81. <https://doi.org/10.1002/path.1700340506>.
- [11] McLeod JW. The types mitis, Intermedius and Gravis of *Corynebacterium diphtheriae*: a review of observations during the past ten years. Bacteriol Rev 1943;7:1–41.
- [12] Hall AJ, Cassidy PK, Bernard KA, Bolt F, Steigerwalt AG, Bixler D, et al. Novel *Corynebacterium diphtheriae* in domestic cats. Emerging Infect Dis 2010;16:688–91. <https://doi.org/10.3201/eid1604.091107>.
- [13] Tagini F, Pillonel T, Croxatto A, Bertelli C, Koutsokera A, Lovis A, et al. Distinct genomic features characterize two clades of *Corynebacterium diphtheriae*: proposal of *Corynebacterium diphtheriae* subsp. *diphtheriae* subsp. nov. and *Corynebacterium diphtheriae* subsp. *lausannense* subsp. nov. Front Microbiol 2018;9:1743. <https://doi.org/10.3389/fmicb.2018.01743>.
- [14] Efstratiou A, George RC. Laboratory guidelines for the diagnosis of infections caused by *Corynebacterium diphtheriae* and *C. ulcerans*. World Health Organization. Comm Dis Publ Health 1999;2:250–7.
- [15] Pimenta FP, Matias GAM, Pereira GA, Camello TCF, Alves GB, Rosa ACP, et al. A PCR for *dtxR* gene: application to diagnosis of non-toxicogenic and toxicogenic *Corynebacterium diphtheriae*. Mol Cell Probes 2008;22:189–92. <https://doi.org/10.1016/j.mcp.2008.01.001>.
- [16] Riegel P, Ruimy R, de Briel D, Prévost G, Jehl F, Christen R, et al. Taxonomy of *Corynebacterium diphtheriae* and related taxa, with recognition of *Corynebacterium ulcerans* sp. nov. nom. rev. FEMS Microbiol Letters 1995;126:271–6. <https://doi.org/10.1111/j.1574-6968.1995.tb07429.x>.
- [17] Pacheco LGC, Pena RR, Castro TLP, Dorella FA, Bahia RC, Carminati R, et al. Multiplex PCR assay for identification of *Corynebacterium pseudotuberculosis* from pure cultures and for rapid detection of this pathogen in clinical samples. J Med Microbiol 2007;56:480–6. <https://doi.org/10.1099/jmm.0.46997-0>.
- [18] Khamis A, Raoult D, La Scola B. *rpoB* gene sequencing for identification of *Corynebacterium* species. J Clin Microbiol 2004;42:3925–31. <https://doi.org/10.1128/JCM.42.9.3925-3931.2004>.
- [19] Hauser D, Popoff MR, Kiredjian M, Boquet P, Bimet F. Polymerase chain reaction assay for diagnosis of potentially toxigenic *Corynebacterium diphtheriae* strains: correlation with ADP-ribosylation activity assay. J Clin Microbiol 1993;31:2720–3.
- [20] Badell E, Guillot S, Tulliez M, Pascal M, Panunzi LG, Rose S, et al. Improved quadruplex real-time PCR assay for the diagnosis of diphtheria. J Med Microbiol 2019;68:1455–65. <https://doi.org/10.1099/jmm.0.001070>.
- [21] Berger A, Hogardt M, Konrad R, Sing A. Detection Methods for Laboratory Diagnosis of Diphtheria. *Corynebacterium Diphtheriae* and Related Toxigenic Species: Genomics, Pathogenicity and Applications. Andreas Burkovski Ed. Springer; 2014. Springer.
- [22] Efstratiou A, Maple CPA. Laboratory Diagnosis of Diphtheria. WHO; 1994. <https://apps.who.int/iris/handle/10665/108108>.
- [23] Rodrigues C, Passet V, Rakotondraso A, Brisse S. Identification of *Klebsiella pneumoniae*, *Klebsiella quasipneumoniae*, *Klebsiella variicola* and related phylogroups by MALDI-TOF mass spectrometry. Front Microbiol 2018;9:3000. <https://doi.org/10.3389/fmicb.2018.03000>.
- [24] Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. J Comput Biol 2012;19:455–77. <https://doi.org/10.1089/cmb.2012.0021>.
- [25] Richter M, Rosselló-Móra R, Oliver Glöckner F, Peplis J. JSpeciesWS: a web server for prokaryotic species circumscription based on pairwise genome comparison. Bioinformatics 2016;32:929–31. <https://doi.org/10.1093/bioinformatics/btv681>.
- [26] Almeida S, Sousa C, Abreu V, Diniz C, Dorneles EMS, Lage AP, et al. Exploration of nitrate reductase metabolic pathway in *Corynebacterium pseudotuberculosis*. Int J Genomics 2017;9481756. <https://doi.org/10.1155/2017/9481756>.
- [27] Ondov BD, Treangen TJ, Melsted P, Mallonee AB, Bergman NH, Koren S, et al. Mash: fast genome and metagenome distance estimation using MinHash. Genome Biol 2016;17.
- [28] Criscuolo A. A fast alignment-free bioinformatics procedure to infer accurate distance-based phylogenetic trees from genome assemblies. Research Ideas and Outcomes 2019;5:e36178. <https://doi.org/10.3897/rio.5.e36178>.
- [29] Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. Mol Biol Evol 2013;30:772–80. <https://doi.org/10.1093/molbev/mst010>.
- [30] Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol 2015;32:268–74. <https://doi.org/10.1093/molbev/mst030>.
- [31] De Zoysa A, Efstratiou A, Mann G, Harrison TG, Fry NK. Development, validation and implementation of a quadruplex real-time PCR assay for identification of potentially toxigenic corynebacteria. J Med Microbiol 2016;65:1521–7. <https://doi.org/10.1099/jmm.0.000382>.
- [32] Santos AS, Ramos RT, Silva A, Hirata R, Mattos-Guaraldi AL, Meyer R, et al. Searching whole genome sequences for biochemical identification features of emerging and reemerging pathogenic *Corynebacterium* species. Funct Integr Genom 2018;18:593–610. <https://doi.org/10.1007/s10142-018-0610-3>.
- [33] Soriano F, Zapardiel J, Nieto E. Antimicrobial susceptibilities of *Corynebacterium* species and other non-spore-forming gram-positive bacilli to 18 antimicrobial agents. Antimicrob Agents Chemother 1995;39:208–14. <https://doi.org/10.1128/aac.39.1.208>.
- [34] Konstantinidis KT, Ramette A, Tiedje JM. The bacterial species definition in the genomic era. Philos Trans R Soc Lond B Biol Sci 2006;361:1929–40. <https://doi.org/10.1098/rstb.2006.1920>.
- [35] Chun J, Oren A, Ventosa A, Christensen H, Arahall DR, da Costa MS, et al. Proposed minimal standards for the use of genome data for the taxonomy of prokaryotes. Int J Syst Evol Microbiol 2018;68:461–6. <https://doi.org/10.1099/ijsem.0.002516>.