ICTV Virus Taxonomy Profile: Pseudoviridae

Carlos Llorens, Beatriz Soriano, Mart Krupovic, Ictv Report Consortium

► To cite this version:

Carlos Llorens, Beatriz Soriano, Mart Krupovic, Ictv Report Consortium. ICTV Virus Taxonomy Profile: Pseudoviridae. Journal of General Virology, 2021, 102 (3), pp.001563. 10.1099/jgv.0.001563 .
pasteur-03188402

HAL Id: pasteur-03188402

<https://hal-pasteur.archives-ouvertes.fr/pasteur-03188402>

Submitted on 2 Apr 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution| 4.0 International License

ICTV Virus Taxonomy Profile: *Pseudoviridae*

Carlos Llorens^{1,*}, Beatriz Soriano¹, Mart Krupovic^{2,*} and ICTV Report Consortium

Abstract

Pseudoviridae is a family of reverse-transcribing viruses with long terminal repeats (LTRs) belonging to the order *Ortervirales*. Pseudoviruses are commonly found integrated in the genomes of diverse plants, fungi and animals and are broadly known as Ty1/Copia LTR retrotransposons. Inside the cell, they form icosahedral virus particles, but unlike most other viruses, do not have an extracellular phase. This is a summary of the ICTV Report on the family *Pseudoviridae*, which is available at [ictv.global/report/pseudoviridae](https://www.ictv.global/report/pseudoviridae).

Table 1. Characteristics of members of the family *Pseudoviridae*

Example:	<i>Saccharomyces cerevisiae</i> Ty1 virus (M18706), species <i>Saccharomyces cerevisiae</i> Ty1 virus, genus <i>Pseudovirus</i>
Virion	Virions are icosahedral ($T=3$ or 4) and might be enveloped
Genome	Two identical copies of linear single-stranded RNA
Replication	Replication by reverse transcription primed with a host-encoded tRNA
Translation	Genomic RNA is translated into one or more polypeptides
Host range	Fungi, plants and animals
Taxonomy	Realm <i>Riboviria</i> , kingdom <i>Pararnavirae</i> , phylum <i>Artverviricota</i> , class <i>Revtraviricetes</i> , order <i>Ortervirales</i> , family <i>Pseudoviridae</i> ; the genera <i>Pseudovirus</i> , <i>Hemivirus</i> and <i>Sirevirus</i> include >30 species

VIRION

Pseudoviruses form intracellular, somewhat irregularly shaped virus-like particles (VLPs) 30–40 nm in diameter, which do not display infectivity and remain intracellular (Table 1). VLPs are formed by self-assembly of proteins encoded by the *gag* gene, namely the capsid (CP) and nucleocapsid (NC) proteins, which are homologous to the equivalent proteins of retroviruses and other members of the order *Ortervirales* [1]. Expression of truncated Gag protein variants yields icosahedral VLPs of different diameters but with a mean radius of 20 nm built on the $T=3$ or $T=4$ lattice (Fig. 1) [2, 3].

GENOME

The genome of pseudovirids ranges from 4 kb to >9 kb and has an internal region flanked by two identical non-coding

sequences called long terminal repeats (LTRs) (Fig. 2). LTRs are variable in size and contain three regions, named U3-R-U5 in analogy to retroviral LTRs. U3 contains promoters, R is repeated on each end of the transcript, and U5 constitutes the first portion of the reverse-transcribed genome. The internal region is delimited by two short motifs: the primer binding site (PBS), which is located downstream of the 5'-LTR and is usually complementary to the initiator tRNA^{Met}, and the polypurine tract (PPT), which is upstream of the 3'-LTR. The internal region may contain one (*gag-pol*), two (*gag* and *pol*) or three (*gag*, *pol* and *env*) ORFs. The Gag polypeptide includes domains for the CP and NC proteins, while Pol includes domains for the protease (PR), integrase (INT) and reverse transcriptase-ribonuclease H (RT-RH). Members of the genus *Sirevirus* carry a third ORF downstream of *gag-pol* encoding a putative envelope protein [4].

Received 23 December 2020; Accepted 15 January 2021; Published 02 February 2021

Author affiliations: ¹Biotechvana, Scientific Park University of Valencia, 46980, Paterna, Valencia, Spain; ²Archaeal Virology Unit, Institut Pasteur, 75015 Paris, France.

***Correspondence:** Carlos Llorens, carlos.llorens@biotechvana.com; Mart Krupovic, mart.krupovic@pasteur.fr

Keywords: *Pseudoviridae*; ICTV Report; taxonomy.

Abbreviations: CP, capsid; INT, integrase; LTR, long terminal repeat; NC, nucleocapsid; PBS, primer binding site; PPT, polypurine tract; PR, protease; RH, ribonuclease H; RT, reverse transcriptase; VLP, virus-like particles.

001563 © 2021

This is an open-access article distributed under the terms of the Creative Commons Attribution License.

Fig. 1. *Saccharomyces cerevisiae* Ty1 virus particles formed from truncated capsid protein (aa 1–381). The surface structures of two forms ($T=3$, left; $T=4$, right) of around 30–40 nm determined by cryo-electron microscopy, are flanked by the corresponding schematic models. (Courtesy of H. Saibil, adapted from [3] with permission from American Society for Microbiology.)

REPLICATION

The replication of pseudovirids resembles that of retroviruses and occurs via reverse transcription in the VLP. The cellular tRNA molecule, typically initiator tRNA^{Met}, which is packaged into the VLP, anneals to the viral RNA genome at the PBS complementary to the 3'-end of that tRNA and is used by RT as a primer to start reverse transcription. Once the full-length proviral cDNA is synthesized, it is imported into the nucleus, where it is integrated into a chromosomal target site by INT. The integrated form (equivalent to the retroviral provirus) is transcribed by the host RNA polymerase II to generate a new viral RNA molecule, which is translated to produce the Gag and Pol polyproteins and is packaged into the VLPs to reinitiate the replication cycle.

TAXONOMY

Current taxonomy: ictv.global/taxonomy. The family *Pseudoviridae* belongs to the order *Ortervirales* [5] and includes the genera *Pseudovirus*, *Hemivirus* and *Sirevirus*. Current classification is based on host tropism, gene content and the length of the tail of the tRNA used as a primer to initiate reverse transcription. Hemiviruses differ from members of the genus *Pseudovirus* in that they use only a short segment of the tRNA. By contrast, sireviruses are restricted to plants and mostly encode a protein equivalent to retroviral Env downstream of the *gag-pol* gene [4]. Further updates and revisions in the genus demarcation of pseudovirids are expected to be based on phylogenetic criteria as described in the full ICTV Report (see Resources).

RESOURCES

Full ICTV Report on the family *Pseudoviridae*: ictv.global/report/pseudoviridae.

Fig. 2. Pseudovirid genome organization. LTRs are white and show labels for the U3, R and U5 regions. Other labels are: capsid (CP); integrase (INT); long terminal repeat (LTR); nucleocapsid (NC); polypurine tract (PPT); primer binding site (PBS); protease (PR); reverse transcriptase (RT); ribonuclease H (RH); surface (SU); transmembrane (TM).

Gypsy Database (GyDB) devoted to viruses and mobile genetic elements: <http://gydb.org>.

Funding information

B.S. was supported by the pre-doctoral research fellowship from Industrial Doctorates of MINECO (Grant 659 DI-17-09134). Production of this Profile, the ICTV Report, and associated resources was funded by a grant from the Wellcome Trust (WT108418AIA).

Acknowledgements

Members of the ICTV Report Consortium are Stuart G. Siddell, Andrew J. Davison, Elliot J. Lefkowitz, Sead Sabanadzovic, Peter Simmonds, Donald B. Smith, Richard J. Orton and Balázs Harrach.

Conflicts of interest

The authors declare that there are no conflicts of interest.

References

1. Krupovic M, Koonin EV. Homologous capsid proteins testify to the common ancestry of retroviruses, caulimoviruses, pseudoviruses, and metaviruses. *J Virol* 2017;91:e00210–17.
2. Burns NR, Saibil HR, White NS, Pardon JF, Timmins PA *et al*. Symmetry, flexibility and permeability in the structure of yeast retrotransposon virus-like particles. *EMBO J* 1992;11:1155–1164.
3. Palmer KJ, Tichelaar W, Myers N, Burns NR, Butcher SJ *et al*. Cryo-electron microscopy structure of yeast Ty retrotransposon virus-like particles. *J Virol* 1997;71:6863–6868.
4. Peterson-Burch BD, Wright DA, Laten HM, Voytas DF. Retroviruses in plants? *Trends Genet* 2000;16:151–152.
5. Krupovic M, Blomberg J, Coffin JM, Dasgupta I, Fan H *et al*. *Ortervirales*: new virus order unifying five families of reverse-transcribing viruses. *J Virol* 2018;92:e00515–18.