



Bordetella parapertussis Bacteremia: Clinical Expression and Bacterial Genomics

Julie Toubiana, Saba Azarnoush, Valérie Bouchez, Annie Landier, Sophie Guillot, Soraya Matczak, Stéphane Bonacorsi, Sylvain Brisse

► To cite this version:

Julie Toubiana, Saba Azarnoush, Valérie Bouchez, Annie Landier, Sophie Guillot, et al.. Bordetella parapertussis Bacteremia: Clinical Expression and Bacterial Genomics. Open Forum Infectious Diseases, 2019, 6 (4), pp.ofz122. 10.1093/ofid/ofz122 . pasteur-03206244

HAL Id: pasteur-03206244

<https://pasteur.hal.science/pasteur-03206244>

Submitted on 23 Apr 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Bordetella parapertussis Bacteremia: Clinical Expression and Bacterial Genomics

Julie Toubiana,^{1,2,3} Saba Azarnoush,⁴ Valérie Bouchez,¹ Annie Landier,¹ Sophie Guillot,¹ Soraya Matczak,³ Stéphane Bonacorsi,⁵ and Sylvain Brisse^{1,2}

¹ Institut Pasteur, Biodiversity and Epidemiology of Bacterial Pathogens, Paris, France; ² National Reference Center for Whooping Cough and Other *Bordetella* Infections, Paris, France; ³ Department of General Pediatrics and Pediatric Infectious Diseases, Necker-Enfants Malades Hospital, APHP, Paris, France; ⁴ Department of Hematology, Robert Debré Hospital, APHP, Paris, France; ⁵ Microbiology Department, Robert Debré Hospital, APHP, Paris, France

Whooping cough's primary etiological agent is *Bordetella pertussis*. The closely related *Bordetella parapertussis* rarely causes severe disease. Here we report an unusual case of bacteremia caused by *B. parapertussis*, review the literature, and characterize the genomic sequence of the bacterial isolate in comparison with *B. parapertussis* isolates from respiratory infections.

Keywords. bacteremia; *Bordetella parapertussis*; children; genomics.

Bordetella pertussis and *Bordetella parapertussis* are the main causative agents of whooping cough, a disease that is still endemic worldwide despite high pertussis vaccine coverage of young children. *B. parapertussis* is much less often involved in the disease than *B. pertussis*, as it is responsible for only 2%–20% of cases and generally causes a less severe respiratory illness [1–3].

Invasive illnesses with *Bordetella* spp. are generally caused by *B. holmesii* and *B. bronchiseptica* in immunocompromised hosts, such as patients with asplenia, chronic obstructive pulmonary disease, HIV infections, hematological disorders, or transplantation [4–6]. Reports of systemic infection with *B. parapertussis* are scarce, with only 3 cases reported in the literature [2, 7]; 2 of them had a known underlying disease. This systemic manifestation of *B. parapertussis* infection might thus be at least in part determined by host susceptibility.

Phylogenetic analyses of *B. pertussis* and *B. parapertussis* indicated that these 2 closely related human-adapted species have evolved independently from *B. bronchiseptica*-like

ancestors [8]. Further, the population of *B. parapertussis* isolates from humans is highly homogeneous genetically [9, 10]. Unlike *B. pertussis*, *B. parapertussis* does not express the pertussis toxin because of many mutations, particularly within the toxin gene promoter region [11]. In addition, *B. parapertussis* has an O-antigen that confers protection from complement-mediated immunity, and can therefore be considered a virulent factor [12]. The possibility that specific virulence determinants of *B. parapertussis* isolates are implicated in invasive infections remains to be investigated, as genomic sequences from *B. parapertussis* causing systemic disease have not been reported so far.

In this study, we report a clinical case of *B. parapertussis* bacteremia. Further, we determined the genomic sequence of the isolate and compared it with all publicly available genomic sequences of *B. parapertussis* from respiratory infections.

CASE REPORT

The patient was a 3-year-old female without notable medical history. She had received 3 injections for primary vaccination against pertussis (acellular pertussis vaccine) before 1 year of age. She was living in Algeria when an acute leukemia was suspected by the presence of fever, lethargy, and hepatosplenomegaly at physical examination. The diagnosis of acute lymphoblastic leukemia was confirmed in France in April 2017 by hemogram, with an abnormal white blood cell count $42 \times 10^9/L$ with $33 \times 10^9/L$ blasts, anemia, and thrombocytopenia, and by the bone marrow diffuse infiltration of undifferentiated blasts (88%) characterized on immunophenotyping analysis. A central venous catheter was implanted, and chemotherapy was started according to the French Acute Lymphoblastic Leukemia Study Group guidelines. One day before chemotherapy started, the patient presented with fever (38.1°C) associated with a moderate cough, which resolved within the next 24 hours. Inflammatory parameters were slightly elevated, with plasma levels of C-reactive protein at 33 mg/L, then 53 mg/L at day 2. Intravenous piperacillin–tazobactam treatment was started 5 days later, when gram-negative bacilli were observed on the first blood culture, and was switched to oral azithromycin when *Bordetella parapertussis* was identified by Matrix-Assisted Laser Desorption Ionization Time of Flight (MALDI-TOF) mass spectrometry. Real-time polymerase chain reaction targeting the IS1001 insertion sequence was performed on a nasopharyngeal aspirate [13]. Its positivity suggested the presence of *B. parapertussis*. The patient did not present any complications, and the antileukemia chemotherapy was continued. The patient was discharged on day 35 of hospitalization and followed the French guidelines for lymphoblastic acute leukemia chemotherapy without any other infectious complication.

Received 12 December 2018; editorial decision 22 February 2019; accepted 4 March 2019.

Correspondence: J. Toubiana, MD, PhD, Institut Pasteur, Biodiversity and Epidemiology of Bacterial Pathogens, 25 rue du Docteur Roux, Paris, France (julie.toubiana@pasteur.fr).

Open Forum Infectious Diseases®

© The Author(s) 2019. Published by Oxford University Press on behalf of Infectious Diseases Society of America. This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial reproduction and distribution of the work, in any medium, provided the original work is not altered or transformed in any way, and that the work is properly cited. For commercial re-use, please contact journals.permissions@oup.com
DOI: 10.1093/ofid/ofz122

The organism collected from blood culture was sent to the National Reference Center for further investigations (Supplementary Data). The isolate (named FR6242) displayed a brown pigment and a visible hemolysis on Bordet-Gengou agar medium, as expected for *B. parapertussis*. Western blots revealed that it was positive for filamentous hemagglutinin (FHA) production and negative for pertussis toxin, but also negative for pertactin (PRN) production, as observed for most human *B. parapertussis* isolates since 2007 in France [14]. There are no clinical breakpoints for determination of susceptibility or resistance of *Bordetellae* from European Committee on Antimicrobial Susceptibility Testing (EUCAST). No resistance to erythromycin and azithromycin has been reported for *B. parapertussis*, but previous studies have found that *B. pertussis* isolates resistant to erythromycin and azithromycin have, in most cases, a minimum inhibitory concentration (MIC) >256 mg/L [15, 16]. The isolate was considered susceptible to azithromycin, clarithromycin, and erythromycin (MIC, 0.12, 0.19, and 0.125 mg/L, respectively) and trimethoprim-sulfamethoxazole (MIC of trimethoprim, 0.012 mg/L). The genomic sequence was assembled into 78 contigs, with a total size of 4 720 641 base pairs, and was compared with 11 human *B. parapertussis* genomes (mainly collected in the United States) available in public sequence repositories. According to the 7-gene multilocus sequence typing (MLST) scheme [10], FR6242 belonged to ST19, as did the reference *B. parapertussis* strain 12822 (isolated in 1993 from an infant with cough illness) [17] and the 10 other isolates. Genotyping of

virulence-related genes revealed the sequence identity between isolate FR6242 and the other isolates for the main targets including pertussis toxin subunit-encoding genes (Supplementary Table 1). Only the gene *brkB*, involved in *Bordetella* serum resistance, showed variation, with allele 2 in reference strain 12822 and allele 6 in all isolates, including FR6242. This change corresponds to a nonsynonymous SNP in position 742, leading to the modification of AA₂₄₈ of the BrkB protein (threonine vs alanine). Analysis of the *prn* gene coding for pertactin revealed that FR6242 had 2 deletions within this gene. A total of 54 whole-genome single nucleotide polymorphisms (SNP) were identified between FR6242 and reference strain 12822. Among them, 23 nonsynonymous SNPs were shared with the other 10 isolates (Supplementary Table 2). Of these, FR6242 shared 19 SNPs with the most recent isolates (collected after 2004), and more particularly with strain I440 collected in 2012 (unspecified clinical origin). Phylogenetic relationships among the 12 *B. parapertussis* genomes (Figure 1) showed that strain FR6242 was related to I440 and more broadly to a clade comprising the recent isolates, which were clearly grouped into a clade separated from more ancient isolates. Only 2 nonsynonymous substitutions and 1 intergenic nucleotide substitution were specific to the invasive strain FR6242: (1) within gene BPP_RS12865/BPP2546, encoding a putative short chain dehydrogenase; (2) within gene BPP_RS18460/BPP3661, encoding a putative oxidoreductase; and (3) at an intergenic position located between 2 conserved hypothetical proteins (Supplementary Table 2).




Figure 1. Single nucleotide polymorphism–based phylogenetic tree of *Bordetella parapertussis* isolates rooted on ovine Bpp5. The analysis focused on 12 human *B. parapertussis* isolates from the period 1935–2017, including FR6242 (bottom box). The analysis was performed by a mapping approach using the genome sequence of *B. parapertussis* strain 12822 (NC_002928.3) as the reference. The scale bar indicates the number of nucleotide substitutions per site.

DISCUSSION

B. parapertussis is circulating worldwide [1] and can cause outbreaks [18]. In France, this species represents 5%–6% of whooping cough cases [19]. It is associated with mild forms of respiratory disease. Invasive infections by *Bordetella* species are typically observed in immunocompromised hosts, and in most described cases they have been associated with initial respiratory diseases. Most reported bacteremia cases have involved *B. holmesii* and *B. bronchiseptica*. Before this study, only 3 *B. parapertussis* bacteremia cases had been reported, to our knowledge [2, 7]. These cases occurred (1) in a patient with severe asthma needing steroid treatment, which is known to cause immunological dysfunction; (2) in a patient with T-cell acute lymphoblastic leukemia; and (3) in an infant without known medical history who died from respiratory arrest. All 3 cases had concomitant lower respiratory tract infection. Here, we report a fourth case, who had acute leukemia and could be considered immunocompromised with associated susceptibility to infection, even if the patient had not started chemotherapy yet and was not neutropenic at the time of infection. As serology for HIV was negative, no immune functional tests were performed. However, the patient did not display severe symptoms of infection, and blood culture had been performed systematically due to fever occurring in an immunocompromised background.

The *B. parapertussis* isolate FR6242 from the present case was not distinguishable from respiratory *B. parapertussis* isolates based on its antigens and virulence factor gene sequences. Genome-wide analysis revealed 2 amino acid changes and 1 intergenic nucleotide substitution unique to this isolate, as well as a number of other changes shared with subsets of *B. parapertussis* isolates. However, it is not immediately clear how the genomic features disclosed here might be associated with invasive disease, and they may as well represent natural variation with no effect of the capacity of the bacterium to cause invasive infection. Knowing whether these changes are associated with *B. parapertussis* bacteremia will require the availability of genomic sequences of multiple other isolates and experimental demonstration of possible effects on invasiveness. In *B. holmesii*, no differences were found between isolates of respiratory and bacteremia origins, using either genomic analysis or cellular/animal models [20, 21]. Therefore, the bacterial determinants of *Bordetella* bacteremia remain elusive.

Clearly, *B. parapertussis* bacteremia might also be related to the host immune response, rather than primarily to bacterial determinants, as found for *B. holmesii* and *B. bronchiseptica* [4–6]. In addition, *B. parapertussis* might induce a different immune response from *B. pertussis*, as it differs in the structure of lipopolysaccharide (LPS) [22]. In vitro, purified *B. parapertussis* LPS is a stronger activator of the innate immune response than purified *B. pertussis* lipooligosaccharide in terms of maturation of human dendritic cells and cytokine production [23], but the

TLR4 response induced by *B. parapertussis* LPS seems less efficient, allowing the organism to escape from a robust inflammatory response, efficient bacterial clearance [24], and subsequent bacterial dissemination in an immune-compromised host.

Possible cross-immunity against *B. parapertussis*, induced by acellular vaccines that target *B. pertussis* virulence, was underlined in several studies [25]. Such a cross-protection could be due to greater anti-FHA antibody response induced by acellular FHA-containing vaccines, as compared with whole-cell vaccines. In our study, the patient was immunized against *B. pertussis*; however, vaccine-induced immunity was not sufficient to protect against *B. parapertussis*.

In conclusion, we report the fourth case of *B. parapertussis* bacteremia in the literature, and the first for which the genomic sequence of the infectious isolate was determined. This new case underlines that *B. parapertussis* can cause invasive infection, mostly in immunocompromised hosts. Investigation of bacterial genomic determinants potentially associated with invasive infection will require the genomic study of additional isolates.

Supplementary Data

Supplementary materials are available at *Open Forum Infectious Diseases* online. Consisting of data provided by the authors to benefit the reader, the posted materials are not copyedited and are the sole responsibility of the authors, so questions or comments should be addressed to the corresponding author.

Acknowledgments

Author contributions. J.T. and S.M. wrote the report. A.S. managed the patient. S.Bo. carried out laboratory testing and identified *B. parapertussis*. A.L. and S.G. carried out the culture of the strain. V.B. and S.Br. carried out genome sequencing and interpreted the data. All authors were involved in the revision of the manuscript. Written consent to publication was obtained.

Consent for publication. Written consent from the patient's parents was obtained before writing of the manuscript. All potentially identifying information was removed from the text.

Sequence availability. Raw sequence data (fastq files) were deposited in the European Nucleotide Archive and are available under study accession number PRJEB29316.

Financial support. This work was supported by Santé Publique France (Saint-Maurice, France).

Potential conflicts of interest. All authors: no reported conflicts of interest. All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Conflicts that the editors consider relevant to the content of the manuscript have been disclosed.

References

1. Watanabe M, Nagai M. Whooping cough due to *Bordetella parapertussis*: an unresolved problem. *Expert Rev Anti Infect Ther* 2004; 2:447–54.
2. Wallihan R, Selvarangan R, Marcon M, et al. *Bordetella parapertussis* bacteremia: two case reports. *Pediatr Infect Dis J* 2013; 32:796–8.
3. Cherry JD, Seaton BL. Patterns of *Bordetella parapertussis* respiratory illnesses: 2008–2010. *Clin Infect Dis* 2012; 54:534–7.
4. Matic NA, Bunce PE. Isolation of *Bordetella bronchiseptica* from blood and a pancreatic abscess. *J Clin Microbiol* 2015; 53:1778–80.
5. Pittet LF, Emonet S, Schrenzel J, et al. *Bordetella holmesii*: an under-recognised *Bordetella* species. *Lancet Infect Dis* 2014; 14:510–9.
6. Powers HR, Shah K. *Bordetella bronchiseptica* bloodstream infection in a renal transplant patient. *Transpl Infect Dis* 2017; 19(6).
7. Correa-Londono A, Ellner PD. Case report. *Clin Microbiol Newsletter* 1980; 2:4.
8. Linz B, Ivanov YV, Preston A, et al. Acquisition and loss of virulence-associated factors during genome evolution and speciation in three clades of *Bordetella* species. *BMC Genomics* 2016; 17:767.

9. Bouchez V, Brun D, Dore G, et al. *Bordetella parapertussis* isolates not expressing pertactin circulating in France. Clin Microbiol Infect **2011**; 17:675–82.
10. Diavatopoulos DA, Cummings CA, Schouls LM, et al. *Bordetella pertussis*, the causative agent of whooping cough, evolved from a distinct, human-associated lineage of *B. bronchiseptica*. PLoS Pathog **2005**; 1:e45.
11. Aricò B, Rappuoli R. *Bordetella parapertussis* and *Bordetella bronchiseptica* contain transcriptionally silent pertussis toxin genes. J Bacteriol **1987**; 169:2847–53.
12. Zhang X, Rodríguez ME, Harvill ET. O antigen allows *B. parapertussis* to evade *B. pertussis* vaccine-induced immunity by blocking binding and functions of cross-reactive antibodies. PLoS One **2009**; 4:e6989.
13. Roorda L, Buitenwerf J, Ossewaarde JM, van der Zee A. A real-time PCR assay with improved specificity for detection and discrimination of all clinically relevant *Bordetella* species by the presence and distribution of three insertion sequence elements. BMC Res Notes **2011**; 4:11.
14. Bouchez V, Guiso N. *Bordetella pertussis*, *B. parapertussis*, vaccines and cycles of whooping cough. Pathog Dis **2015**; 73.
15. Guillot S, Descours G, Gillet Y, et al. Macrolide-resistant *Bordetella pertussis* infection in newborn girl, France. Emerg Infect Dis **2012**; 18:966–8.
16. Wang Z, Cui Z, Li Y, et al. High prevalence of erythromycin-resistant *Bordetella pertussis* in Xi'an, China. Clin Microbiol Infect **2014**; 20:O825–30.
17. Parkhill J, Sebaihia M, Preston A, et al. Comparative analysis of the genome sequences of *Bordetella pertussis*, *Bordetella parapertussis* and *Bordetella bronchiseptica*. Nat Genet **2003**; 35:32–40.
18. Koepke R, Bartholomew ML, Eickhoff JC, et al. Widespread *Bordetella parapertussis* infections—Wisconsin, 2011–2012: clinical and epidemiologic features and antibiotic use for treatment and prevention. Clin Infect Dis **2015**; 61:1421–31.
19. Tubiana S, Belchior E, Guillot S, et al; Renacoq Participants. Monitoring the impact of vaccination on pertussis in infants using an active hospital-based pediatric surveillance network: results from 17 years' experience, 1996–2012, France. Pediatr Infect Dis J **2015**; 34:814–20.
20. Bouchez V, AlBitar-Nehme S, Novikov A, Guiso N, Caroff M. *Bordetella holmesii*: lipid A structures and corresponding genomic sequences comparison in three clinical isolates and the reference strain ATCC 51541. Int J Mol Sci **2017**; 18.
21. Harvill ET, Goodfield LL, Ivanov Y, et al. Genome sequences of nine *Bordetella holmesii* strains isolated in the United States. Genome Announc **2014**; 2(3):e00438–14.
22. Caroff M, Aussel L, Zarrouk H, et al. Structural variability and originality of the *Bordetella* endotoxins. J Endotoxin Res **2001**; 7:63–8.
23. Fedele G, Nasso M, Spensieri F, et al. Lipopolysaccharides from *Bordetella pertussis* and *Bordetella parapertussis* differently modulate human dendritic cell functions resulting in divergent prevalence of Th17-polarized responses. J Immunol **2008**; 181:208–16.
24. Wolfe DN, Buboltz AM, Harvill ET. Inefficient Toll-like receptor-4 stimulation enables *Bordetella parapertussis* to avoid host immunity. PLoS One **2009**; 4:e4280.
25. Liko J, Robison SG, Cieslak PR. Do pertussis vaccines protect against *Bordetella parapertussis*? Clin Infect Dis **2017**; 64:1795–7.