Making sense of the biodiversity and virulence of *Listeria monocytogenes*

Olivier Disson^{1,2}, Alexandra Moura^{1,2,3} and Marc Lecuit^{1,2,3,4}

¹ Institut Pasteur, Biology of Infection Unit, Paris-France

² Inserm U1117, Paris-France

³ Institut Pasteur, French National Reference Center and WHO Collaborating Center for *Listeria*, Paris-France

⁴ Université de Paris, Service des Maladies Infectieuses et Tropicales, Hôpital Universitaire Necker-Enfants Malades, Institut Imagine, APHP, Paris-France

Keywords: *Listeria monocytogenes*, listeriosis, biodiversity, virulence, adaptation

Correspondence to:

Olivier Disson, olivier.disson@pasteur.fr & Marc Lecuit, marc.lecuit@pasteur.fr

Biology of Infection Unit, Institut Pasteur, Inserm U1117, 28 rue du Dr Roux, 75015 Paris, France

Phone: +33 1 40 61 37 87

Fax: +33 1 40 61 34 21

Abstract

Listeria monocytogenes is a foodborne pathogen responsible for listeriosis, an infection which can manifest in humans as bacteremia, meningoen­cephalitis in immunocompromised patients and the elderly, and fetal-placental infection in pregnant women. Reference strains from this facultative intracellular bacterium have been instrumental to investigate basic mechanisms in microbiology, immunology and cell biology. The integration of bacterial population genomics with environmental, epidemiological and clinical data allowed the uncovering of new factors involved in the virulence of *L. monocytogenes* and its adaptation to different environments. This review illustrates how these investigations have led to a better understanding of the bacterium's virulence and the driving forces that shaped it.

Highlights

- *Listeria monocytogenes* is a foodborne opportunistic bacterial pathogen
- Classical genetic approaches in reference strains allowed identification of core virulence genes
- Investigation of the species diversity of *L. monocytogenes* allowed uncovering hypo- and hypervirulent strains and new virulence genes
- Hypervirulent strains are host-associated, whereas hypovirulent strains are associated with food-processing environments
- Accumulation of loss-of-function mutations in virulence pathways leads to niche restriction and obligate saprophytism

Introduction

Listeria monocytogenes is the causal agent of listeriosis, a foodborne infection associated with high morbidity and mortality. The virulence of *L. monocytogenes* depends on its ability to colonize the gut and disseminate in the host. This Gram-positive bacterium was first isolated by G. Hülphers in Sweden, from rabbit liver necrosis [1] and from a human meningitis case in 1921 by J. Dumont [2]. It was then isolated and characterized by E.G.D. Murray in rabbits and guinea pigs from a breeding facility in the UK in 1926, and J.H. Pirie in wild gerbils in South Africa in 1927 [3, 4]. Human listeriosis is characterized by septicemia and central nervous system (CNS) infections, which mostly occur in immunocompromised and/or elderly individuals, and maternal-fetal infections, as well as rare anatomically localized infections [5-12]. *L. monocytogenes* can also induce gastroenteritis in otherwise healthy individuals when the ingested inoculum is large [13]. The foodborne origin of human listeriosis was formally proven in the 1980s [14]. Although listeriosis is relatively rare compared to other foodborne infections [15, 16], its high fatality rate (approximately 20% to 30%, despite effective antibiotic treatment [5]) makes it a significant public health concern [17]. Recently, the largest recorded outbreak occurred in South Africa, where 937 individuals were affected, with a 27% mortality rate [18]. *L. monocytogenes* has also been detected in the feces of healthy people, but the prevalence of *L. monocytogenes* fecal carriage in the general population is unknown [19, 20]. *L. monocytogenes* is therefore mostly considered as an opportunistic pathogen.

L. monocytogenes is not only an economical and public health issue, but is also a powerful model to study basic microbiology, host biology including innate and adaptive immunity and cell biology [21-24]. The identification of *L. monocytogenes* virulence factors in reference strains in cell culture models allowed deciphering the underlying mechanisms of its interactions with the host, and also addressing basic cell biology questions [22]. Further studies combining epidemiologic data with comparative genomics analyses of food- and clinically-associated strains allowed the uncovering of new genes involved in *L. monocytogenes* virulence [25]. Current investigations indicate that virulence factors are key for the adaptation of *L. monocytogenes* to its host and its optimal spread in the environment, and may modify our understanding of how this bacterium has acquired and selected these genes.

Uncovering virulence genes of *L. monocytogenes* by mutagenesis of reference strains

It was shown early on, using *in vitro* and *in vivo* approaches, that *L. monocytogenes* can enter macrophages, survive and be transferred to adjacent cells [26]. *In vivo* experiments in guinea pigs indicated that *L. monocytogenes* can be released in the cytoplasm of intestinal epithelial cells, surrounded by electron dense material [27], which would later come to be identified as polymerized actin. Entry of *L. monocytogenes* in non-phagocytic epithelial cells was reported *in vitro* in the human enterocyte-like cell line Caco-2 [28]. Most of the factors involved in key phenotypes of the cellular microbiology of *L. monocytogenes*, namely entry into host cells, escape from the internalization vacuole and spread from cell to cell, were characterized in the 1980s and 1990s and shown to be critical for virulence *in vivo*. They were mostly discovered by a combination of molecular and cell biology approaches, using transposon bacterial mutagenesis, gene knock-out and complementation studies coupled with infection of cultured macrophages and epithelial cell lines. These factors were characterized in the laboratory reference strains EGD, 10403S or EGDe, which all belong to the serovar 1/2a but to two distinct lineage II clonal complexes (CCs; see below). EGD (CC7) derives from a strain isolated in guinea pig by E. G. D. Murray, hence its name [3]. 10403S (CC7) is a streptomycin-resistant strain derived from 10403, a strain isolated from a human listeriosis case [29]. The origin of EGDe (CC9) is more difficult to trace. It was supposed to be an EGD isolate, but genome analyses have revealed that both strains are actually unrelated [30, 31].

LIPI-1 genes

A phenotypic characteristic of *L. monocytogenes* is its ability to completely lyse red blood cells (beta hemolysis), due to the expression of a hemolysin encoded by *hly* and named listeriolysin O (LLO) [32, 33]. Screening and characterization of nonhemolytic mutants obtained by transposon mutagenesis indicated that hemolysin expression is critically needed for *L. monocytogenes* virulence [34]. Further characterization of Δhly mutant in intestinal Caco-2 cells showed that LLO's pore-forming toxin is involved in escape of *L. monocytogenes* of its internalization vacuole [28]. Later analysis involved the characterization of two phospholipases, PlcA and PlcB (activated by the metalloprotease Mpl), which are also involved in vacuolar escape [35, 36]. Once in the cytosol, *L. monocytogenes* can polymerize actin and form so-called “actin comet tails” via its surface protein ActA, which propel *L. monocytogenes* in the cytoplasm, and allow its spread from cell to cell [37, 38]. The study of *actA* and *plcA* transposon and deletion mutants indicated that they are both critical for *L. monocytogenes* virulence [39, 40]. Analysis of an avirulent non-hemolytic variant of the NCTC 7973 guinea pig strain (isolated by E.G.D. Murray) allowed the characterization of a

transcription factor involved in *hly* expression which was named PrfA for "Positive regulatory factor of listeriolysin (*lisA*, the other designation for *hly*)" [41, 42]. PrfA was further shown to be the major regulator of *L. monocytogenes* virulence genes *in vivo*, including *prfA*, *hly*, *actA*, *plcA*, *mpl*, *plcB* and the nucleomoduline *orfX* [43], which are all encoded by the Listeria pathogenicity island 1 (LIPI-1) [44] (Figure 1A). PrfA is also transcriptionally regulated by σ^B , that controls the general stress response in *L. monocytogenes*, is regulated post-transcriptionally and requires a glutathione co-factor for full activation [45, 46].

inlAB

Transposon mutagenesis was also used to screen for mutants that are unable to enter Caco-2 cells. This approach allowed for the identification and characterization of internalin (InlA), a leucine-rich repeats (LRR)-containing protein anchored to the bacterial peptidoglycan by a Leu-Pro-X-Thr-Gly (LPXTG) motif [47]. The receptor of InlA, namely E-cadherin (Ecad), which is expressed on epithelial cells and forms *adherens* junctions, was identified by affinity chromatography [48]. Further analyses indicated that the InlA-Ecad interaction is species-specific, as mouse and rat Ecad are not a receptor for InlA, in contrast to human Ecad. Indeed, InlA interaction with Ecad requires a proline in the 16th amino acid position of Ecad, which is present in permissive species (human, rabbit, guinea pig, gerbil), and is replaced by a glutamic acid in non-permissive murine species (mouse, rat), preventing InlA interaction with Ecad in the latter [49]. In guinea pigs and transgenic mice expressing human Ecad, InlA-Ecad interaction is critical for *L. monocytogenes* to cross the intestinal barrier, at the villus level, by transcytosis across goblet cells [50, 51]. Live imaging in intestinal organoids has now shown that *L. monocytogenes* hijacks E-cadherin recycling for its: (i) entry into; (ii) transepithelial transfer across; and (iii) exocytosis from goblet cells, in a dynamin-, microtubule- and Rab11-dependent manner, respectively [52]. *inlA* is located in an operon comprising a downstream gene called *inlB* [47]. InlB is also an LRR-containing protein; however, it is not covalently linked to the bacterial surface by an LPXTG motif, but loosely attached to it by its Gly-Trp (GW) motif. InlB mediates the entry of *L. monocytogenes* in cultured cells by interacting with c-Met/HGFR, which is ubiquitously expressed [53, 54]. InlB-c-Met interaction is also species-specific: it occurs in humans, mice and gerbils but not in rabbits nor guinea pigs [55]. InlB is not involved in *L. monocytogenes* crossing of the intestinal barrier [55]. In order to study the role of the InlA-Ecad and InlB-c-Met interactions in organs other than the intestine, knock-in mice KIE16P, in which the glutamic acid in 16th position of the Ecad is replaced by a proline, were generated [56]. In contrast to the intestinal barrier where only InlA is

involved, InlA and InlB are both required and act in a conjugated manner in *L. monocytogenes* crossing of the placental barrier in gerbils and in KIE16P mice [56, 57].

Uncovering virulence genes of *L. monocytogenes* by comparative *Listeria* genomics

While classical genetics has identified major virulence factors in *L. monocytogenes*, whole genome sequencing of pathogenic bacterial species, which was initiated with *Mycoplasma genitalium* [58], opened a new avenue for identifying genes involved in virulence, by comparison of the genomic content of pathogenic and phylogenetically closely related non-pathogenic bacterial species. Indeed, comparison of the genome sequences of *L. monocytogenes* EGDe and the nonpathogenic *Listeria innocua* CLIP 11262 reference strains allowed identification of putative virulence factors present in *L. monocytogenes* but absent in *L. innocua* [59] (Figure 1B).

Sequencing of EGDe revealed the presence of 25 genes that code for proteins of the internalin family, characterized by the presence of an LRR domain. Seventeen of them had not been described before, including InlJ, an LPXTG anchored protein involved in virulence, whose exact function remains to be determined [60, 61] and InlP (*lmo2470*), a secreted protein involved in the infection of the placenta, among other organs [62, 63].

Comparative genomics also revealed the presence of GW motif-containing proteins other than InlB in EGDe (and which were absent in *L. innocua* CLIP 11262) that are involved in virulence. As an example, the GW-containing protein Auto is an autolysin involved in cell entry and virulence [64].

Another gene that is absent in *L. innocua*, named *bsh*, has been identified by this strategy. This PrfA-regulated virulence gene encodes for a bile salt hydrolase, which increases resistance of *L. monocytogenes* to bile, and enhances liver infection and bacterial faecal carriage [65], an essential property for a foodborne pathogen.

Intraspecies genomic studies and identification of novel virulence genes

The biodiversity of L. monocytogenes

The biodiversity of *L. monocytogenes* was first investigated via serotyping, based on *Listeria* somatic and flagellar antigens, which allowed identifying the four main serotypes (1/2a, 1/2b, 1/2c and 4b) associated to listeriosis [66]. However, this method does not allow establishing reliable phylogenetic links between strains. Multilocus enzyme electrophoresis differentiated two main lineages, I and II [67], while gene variation studies and multi-locus genotyping assays have led to the delineation of two minor lineages III and IV [68, 69]. In order to

determine the population structure of *L. monocytogenes*, multi-locus sequence typing (MLST), allowing to translate sequence diversity into numerical profiles, called sequence types (STs), was set up with seven housekeeping genes [70]. Using this technique, isolates having at least 6 alleles in common out of these seven genes are grouped into the same clonal complex (CC). With advances in whole genome sequencing, this method has been extended to larger sets of loci (typically thousands) to allow strain differentiation at the level of the whole- or core genome (wgMLST or cgMLST, respectively) [71-75]. A standard nomenclature system has been proposed based on the cgMLST scheme of 1,748 core loci [75]. Isolates sharing less than 150 and 7 allelic mismatches are grouped into sublineages (SL) and cgMLST types (CT), respectively, which allows for efficient inter-laboratory communication, in particular for emerging strain types [76] (Figure 2A). In addition, the compilation of genotypic data in dedicated databases, such as BIGSdb-*Lm* (<http://bigsdb.pasteur.fr/listeria>) or GenomeTrakr (hosted at <https://www.ncbi.nlm.nih.gov/pathogens>) allow for in-depth studies of the global spread and biodiversity of *L. monocytogenes*.

Hypervirulent strains belong to distinct CCs compared to hypovirulent strains

L. monocytogenes can be divided into four evolutionary lineages [77], 13 serotypes [78] grouped into 6 genosero groups [79] and more than 170 clonal complexes and 300 sublineages (<http://bigsdb.pasteur.fr/listeria>). Of these, lineage I (serotypes 4b and 1/2b) and II (serotypes 1/2a and 1/2c) isolates represent the majority of clinical isolates. The most prevalent CCs and SLs are geographically and temporally widespread [70, 80, 81]. However, an uneven prevalence is observed among food vs. clinical cases [25, 75, 80], as previously noted at the lineage and serotype levels [77]. While lineage II, in particularly sublineages SL9 (CC9) and SL121 (CC121) are mostly associated with food, lineage I isolates, and in particular sublineages SL1 (CC1), SL2 (CC2), SL4 (CC4) and SL6 (CC6) are strongly associated with clinical cases in the Western world [25, 75, 82-87] (Figure 2B). In Asia, specifically in China and Taiwan, the most prevalent sublineage in both food and clinical cases is SL87 (CC87, lineage I, serotype 1/2b) [84, 88, 89], suggesting that food habits and/or local specificities are involved in geographic diversity of *L. monocytogenes*.

A large study of 6,663 *L. monocytogenes* isolates of clinical and food origins prospectively collected in France between 2005 and 2013 reported a higher prevalence of CC9 and CC121 (both of lineage II) isolates in food samples, while CC1, CC6, CC2 and CC4 (all of lineage I) were the most prevalent in clinical cases [25]. In addition, CC1 and CC4 were found to be

statistically associated with more invasive forms of listeriosis (maternal-neonatal and CNS infections). Similar results have been observed within farm environments, with CC1 being also associated with CNS infection in ruminants in Switzerland [90]. This could reflect a difference in virulence and/or in exposure of at-risk patients to these strains. Association of CCs to clinical cases is inversely correlated to patients' immunosuppressive comorbidities, suggesting that these are more virulent than CCs most associated with food [25]. *In vivo* studies, which compared CC1, CC4, CC6, CC9 and CC121 isolates and the laboratory reference strains EGDe and 10403S in a mouse model permissive for oral *L. monocytogenes* infection, unambiguously showed that strains from CCs most associated with clinical cases (CC1, CC4 and CC6) are hypervirulent when compared to laboratory reference strains EGDe and 10403S, as they induced a more pronounced body weight decrease and are more neuroinvasive [25]. In contrast, CC9 and CC121 are hypovirulent when compared to these laboratory reference strains. Of note, CC9 and CC121 are not fully avirulent, since they can still infect laboratory animals, although to a lower degree, as well as immunodeficient patients [25]. This heterogeneity in virulence among lineages and CCs led to an investigation of the putative molecular mechanisms involved in both hypo- and hyper-virulence.

Virulence factors found in the core genome

While *inlA* is present in all *L. monocytogenes* strains, some strains, including the reference laboratory strain LO28 (CC9, lineage II), express a truncated and therefore nonfunctional InlA [91]. Epidemiological analysis indicated that clinical strains, and most significantly those from maternal-fetal infections, express full-length InlA more frequently than strains from food origins [70, 92]. InlA is truncated in the majority of hypovirulent CC9 and CC121 isolates as well as in other lineage II clonal complexes (e. g. CC331, CC199 and CC321[75]). Hypovirulence can therefore, at least in part, be accounted for by InlA truncation, leading to decreased translocation across the intestinal barrier and placental infection [50, 56, 93].

Similarly to *inlAB*, LIPI-1 genes are present in all *L. monocytogenes* SLs and are therefore core virulence genes (Figure 1A). As indicated above, non-pathogenic *L. innocua* lacks LIPI-1 and *inlAB* genes, however rare *L. innocua* strains contain both LIPI-1 and *inlA* that are functional *in vitro* and *in vivo* [94-99]. This indicates that *L. monocytogenes* and *L. innocua* species likely evolved independently from a common pathogenic ancestor harboring both the LIPI-1 and *inlAB* virulence loci. Speciation likely occurred upon differential environmental selective pressures, with *L. innocua* evolving in an environment where virulence genes are not needed [100]. In contrast, 0.1% of *L. monocytogenes* strains from both lineage I and II are

non-hemolytic. 93.3% of these non-hemolytic strains harbor at least one loss-of-function mutation in *prfA*, and the remaining 6.7% harbor *hly* mutations leading to LLO inactivation. In both of these cases, this results in a strong decrease in virulence, suggesting that niche restriction leading to sub-speciation may still occur within *L. monocytogenes* species [101].

Virulence factors encoded by the accessory genome

LIPI-3

The LIPI-3 virulence cluster, which is absent in *L. monocytogenes* reference strains but present in lineage I F2365 (SL1/CC1) and H7858 (SL6/CC6) outbreak strains, has also been characterized (Figure 1A). It consists of 8 genes, including a hemolytic and cytotoxic factor, listeriolysin S (LLS), involved in virulence [102]. LLS, encoded by *llsA*, acts as a bacteriocin and alters microbiota composition, resulting in an increase in the luminal amount of *L. monocytogenes*, while *llsB*, another gene of LIPI-3, plays a role in the systemic phase of the infection [103, 104]. Functional LIPI-3 or its remnants can also be present in some *L. innocua* strains [98, 105], where its role is not defined, reinforcing the hypothesis that the two species share a common pathogenic ancestor. The presence of LIPI-3 in *Lm* sublineages SL1 (CC1), SL4 (CC4) and SL6 (CC6), which are the most associated with clinical cases [75, 98], and its absence in lineage II strains, suggests a possible role in their hypervirulence.

LIPI-4

Comparative genomic analyses derived from biodiversity studies allowed the characterization of LIPI-4, a putative phosphotransferase system (PTS) present in SL4, a sublineage of the lineage I, which strongly associated with a clinical origin (Figure 2B), particularly in CNS and placental infections [25]. LIPI-4 is involved in placental and CNS infection in an *in vivo* model of listeriosis, by a so far unknown mechanism (Figures 1A and 2A) [25]. Interestingly, LIPI-4, is also present in the *L. monocytogenes* sublineage that is the most prevalent in Asia, SL87 (CC87) [106], as well as in *L. innocua* [98]. Its actual role in such a non-virulent species is currently unknown.

LIPI-2

LIPI-2 is a genomic region discovered in *Listeria ivanovii*, a *Listeria* species which is pathogenic for mammals, mainly ovines and bovines, but also rarely humans [108, 109]. LIPI-2 was first described as an *L. ivanovii*-specific genomic region that encodes a sphingomyelinase involved in phagosome disruption, as well as 10 genes of the internalin

family [110]. This region is involved in *L. ivanovii* virulence, together with the *L. ivanovii* LIPI-1 orthologous region, both *in vitro* and in an *in vivo* sheep model of infection [110]. *L. monocytogenes* isolates belonging to a new sublineage of the major lineage II (SL626/CC33, serovar 4h) recovered from an ovine listeriosis outbreak in China have been characterized as hypervirulent [107]. These isolates contain a partial LIPI-2 locus, comprising only the sphingomyelinase *smcL* and the internalins *i-inlF* and *i-inlE* genes (Figure 1A), likely by acquisition of exogenous DNA from *L. ivanovii*. The specific role of the partial LIPI-2 in hypervirulence of this newly described *L. monocytogenes* sublineage remains to be elucidated.

Heterogeneity in virulence as a reflection of differential host adaptation?

The distribution of *L. monocytogenes* is wide. It has been isolated in humans and animals in all continents except Antarctica [80]. Linked to its ability to tolerate extreme temperatures, acidity and osmolarity, *L. monocytogenes* can persist and grow in soil, plants, water, food and silage [111-113]. However, the distribution of *L. monocytogenes* lineages and CCs is not homogeneous in these ecosystems, indicating that CCs emergence may be driven by their selective adaptation to the environment in which they are the most abundant [25, 77].

When investigating food samples, hypervirulent CC1, CC4 and CC6 were found to be statistically associated with dairy products, whereas hypovirulent *L. monocytogenes* clones, CC9 and CC121 were found associated with meat products [114], suggesting a differential adaptation of the CCs to their environment. Interestingly, CC9 and CC121 harbor genes involved in tolerance to benzalkonium chloride, a surface disinfectant commonly used in food production environments, and are more efficient than their hypervirulent counterparts to form biofilms in the presence of this disinfectant [114, 115]. Other stress resistance genes such as the SSI-1 and SSI-2 stress survival islets [116, 117] are also typically present in lineage II, and contribute to *L. monocytogenes* survival and growth in the stress conditions encountered in food and food processing environments (e.g. high salt concentrations, oxidative stress). Lineage I isolates typically display higher tolerance to low pH [118], which can be advantageous for the survival of *L. monocytogenes* in the host's gastric acidic environment. Consistent with these observations, strains of hypervirulent CCs colonize the gut more efficiently than those of hypovirulent CCs [114], which may account for their strong association with ruminant feces [119] and dairy products, which may result from milk fecal contamination [114] (Figure 3).

Concluding remarks

L. monocytogenes is a major model pathogen which has allowed scientific breakthroughs in microbiology, immunology, cell biology and host-pathogen interactions. It has also emerged as a powerful model to study how microbes evolve from their natural habitat to anthropic environments. While advances have been made in the discovery of genetic factors involved in virulence and adaptation to host vs. food processing environments, the mechanisms by which *L. monocytogenes* adapts to various environments remains to be fully deciphered. Seminal studies have indicated how the transcriptome in the EGDe reference strain responds to the switch from saprophytism to virulence [120]. Future large-scale transcriptomics and metabolomics studies with representative strains from different clonal complexes will enrich our understanding of how *L. monocytogenes* can adapt to the environments in which it thrives. Further studies will also be needed to decipher the forces driving hypervirulence and its relationship with adaptation to the host, and how this relates to capacity of *L. monocytogenes* to be shed back in the environment and transmitted to new hosts, as suggested by data from ruminants and animal models [114, 119]. The continuous analysis of biodiversity of *L. monocytogenes*, together with environmental, epidemiological, clinical and experimental data obtained from genome wide association studies will shed light into the mechanisms and selective pressures that drive *Listeria* adaptation to diverse environments.

Figure Legends

Figure 1. Virulence genes of *L. monocytogenes*

A) Typical genetic organization of *Listeria* pathogenicity islands (LIPI; colored arrows). Flanking genes (gray arrows) are labeled based on their orthologs in *L. monocytogenes* EGDe, when existing. *Lm*, *L. monocytogenes*; *Lin*, *L. innocua*; *Liv*, *L. ivanovii*; *, LIPI-2 genes present in *Liv* atypical *L. monocytogenes* CC33 isolates are shown in darker yellow.

B) Genome maps of *L. monocytogenes* EGDe and *L. innocua* CLIP11262 [59]. Inner circles represent the GC skew $((G - C)/(G + C))$ and GC content $((G+C)/(G+C+A+T))$ of *L. monocytogenes* EGDe. Gaps in *L. innocua* chromosome (blue) represent genomic regions absent in *L. monocytogenes* genome (red). The genomic positions of known *L. monocytogenes* virulence genes are shown and their common and/or locus names indicated. Internalin genes are highlighted in red. Triangles denote the genomic positions of LIPI regions in either EGDe (black triangle) or other *Listeria* species (open triangles), as detailed in panel A.

Figure 2. Population structure of *L. monocytogenes*

A) Population structure of *L. monocytogenes* based on cgMLST profiles (adapted from [75]; $N=1,696$ isolates collected from food and clinical samples, mostly in North America and Europe). Branches are colored by phylogenetic lineage (I-IV, shown in red, blue, green and gray, respectively) and collapsed by sublineage (SL). The vertical dashed bar indicates the established cgMLST cut-off to define SLs (i.e. 150 allelic differences out of 1748 loci). Main SLs are labeled on the branches and corresponding MLST clonal complexes (CCs) are shown in the first column, PCR serogroups, source types and continents of isolation are colored according the key panels (upper left). Dark blue boxes denote the presence of selected genetic traits: pathogenicity islands (LIPI-1 to -4), genes involved in teichoic acid biosynthesis (*gltAB*, *tagB* and *gtcA*), genes coding for internalins (*inlABCEFGHJK*), autolysins (*aut* and *autIVb*) and bile resistance (*bsh*), genes involved in benzalkonium resistance (*bcrABC* operon, *ermC*, *Tn6188::ermC* and *ermE*), stress resistance islets (SSI-1 and SSI-2) and biofilm formation and virulence (*comK*).

B) Frequency of the most prevalent *L. monocytogenes* MLST clonal complexes in clinical (y -axis) versus food (x -axis) isolates, based on a exhaustively collected isolates dataset obtained in the scope of listeriosis surveillance ($N=6,633$ isolates, France 2005-2013). (reproduced with permission from [25]). Each circle represents a clonal complex, and is colored according to its lineage. The size of each circle is proportional to the number of isolates it represents. The clonal complexes of common reference lab strains (CC7: strains EGD and 10403S; CC9: EGDe and LO28), are also indicated.

Figure 3. Trade-off between host-associated and environmental lifestyles of *L. monocytogenes*

Properties of *L. monocytogenes* strains adapted to a host environment (in red) versus of those adapted to saprophytism (natural and food-processing environments; in blue). Listeriosis results from the interplay between host and microbial factors: the less immunocompromised is the host, the more virulent *L. monocytogenes* strains need to be to cause disease. Thus, immunocompromised hosts tend to be susceptible to even saprophytic (hypovirulent) *L. monocytogenes*, whereas healthy hosts are usually only susceptible to host-associated (hypervirulent) strains, as shown in [25].

Acknowledgments

382 Research in the Biology of Infection Unit is supported financially by Institut Pasteur, Inserm,
383 ANR and ERC. Marc Lecuit is a senior member of Institut Universitaire de France.

384 References

- 385 1. Hülphers, G. (1911) Lefvernekros hos kanin orsakad af en ej fovut beskriefuen bakterie.
386 Soensk Veterinar Tidskrift 16, 265-273.
- 387 2. Dumont, J. and Cotoni, L. (1921) Bacille semblable au bacille du rouget du porc rencontré
388 dans le liquide céphalo-rachidien d'un méningitique. Ann. Inst. Pasteur (Paris) 35 (10), 625-
389 33.
- 390 3. Murray, E.G.D. et al. (1926) A disease of rabbits characterised by a large mononuclear
391 leucocytosis caused by hitherto undescribed bacillus *Bacterium monocytogenes* (n. sp.). J.
392 Pathol. Bacteriol. 29, 407-439.
- 393 4. Pirie, J.H.H. (1927) A new disease of veld rodents, "Tiger river disease.". Pub South Africa
394 Inst Med Res 62, 163-186.
- 395 5. Charlier, C. et al. (2017) Clinical features and prognostic factors of listeriosis: the
396 MONALISA national prospective cohort study. Lancet Infect Dis 17 (5), 510-519.
- 397 6. Danion, F. et al. (2017) *Listeria monocytogenes* isolation from urine: a series of 15 cases
398 and review. Clinical microbiology and infection : the official publication of the European
399 Society of Clinical Microbiology and Infectious Diseases 23 (8), 583-585.
- 400 7. Chersich, M.F. et al. (2018) Diagnosis and Treatment of *Listeria monocytogenes*
401 Endophthalmitis: A Systematic Review. Ocular immunology and inflammation 26 (4), 508-
402 517.
- 403 8. Charlier, C. et al. (2012) *Listeria monocytogenes*-associated joint and bone infections: a
404 study of 43 consecutive cases. Clinical infectious diseases : an official publication of the
405 Infectious Diseases Society of America 54 (2), 240-8.
- 406 9. Charlier, C. et al. (2014) *Listeria monocytogenes*-associated biliary tract infections: a study
407 of 12 consecutive cases and review. Medicine 93 (18), e105.
- 408 10. Shoai-Tehrani, M. et al. (2019) *Listeria monocytogenes*-associated endovascular
409 infections: A study of 71 consecutive cases. The Journal of infection 79 (4), 322-331.
- 410 11. Morgand, M. et al. (2018) *Listeria monocytogenes*-associated respiratory infections: a
411 study of 38 consecutive cases. Clinical microbiology and infection : the official publication of
412 the European Society of Clinical Microbiology and Infectious Diseases 24 (12), 1339 e1-1339
413 e5.
- 414 12. Pilmis, B. et al. (2020) Cutaneous listeriosis, a case series of 16 consecutive patients over
415 25 years. The Journal of infection 80 (2), 232-254.
- 416 13. Dalton, C.B. et al. (1997) An outbreak of gastroenteritis and fever due to *Listeria*
417 *monocytogenes* in milk. N Engl J Med 336 (2), 100-5.
- 418 14. Schlech, W.F., 3rd et al. (1983) Epidemic listeriosis--evidence for transmission by food. N
419 Engl J Med 308 (4), 203-6.
- 420 15. ECDC (2018) Annual epidemiological report for 2016 Listeriosis. European Centre for
421 Disease Prevention and Control.
- 422 16. Pohl, A.M. et al. (2019) Differences Among Incidence Rates of Invasive Listeriosis in the
423 U.S. FoodNet Population by Age, Sex, Race/Ethnicity, and Pregnancy Status, 2008-2016.
424 Foodborne pathogens and disease 16 (4), 290-297.
- 425 17. European Food Safety, A. et al. (2019) The European Union One Health 2018 Zoonoses
426 Report. EFSA J 17 (12), e05926.
- 427 18. Thomas, J. et al. (2020) Outbreak of Listeriosis in South Africa Associated with Processed
428 Meat. N Engl J Med 382 (7), 632-643.
- 429 19. Sauders, B.D. et al. (2005) Low prevalence of *Listeria monocytogenes* in human stool. J
430 Food Prot 68 (1), 178-81.
- 431 20. Grif, K. et al. (2003) Incidence of fecal carriage of *Listeria monocytogenes* in three
432 healthy volunteers: a one-year prospective stool survey. Eur J Clin Microbiol Infect Dis 22
433 (1), 16-20.

21. Radosheovich, L. and Cossart, P. (2018) *Listeria monocytogenes*: towards a complete picture of its physiology and pathogenesis. *Nature reviews. Microbiology* 16 (1), 32-46.
22. Lecuit, M. (2020) *Listeria monocytogenes*, a model in infection biology. *Cellular microbiology* 22 (4), e13186.
23. Pamer, E.G. (2004) Immune responses to *Listeria monocytogenes*. *Nature reviews. Immunology* 4 (10), 812-23.
24. Maudet, C. et al. (2020) Innate immune responses to *Listeria* in vivo. *Current opinion in microbiology* In press.
25. Maury, M.M. et al. (2016) Uncovering *Listeria monocytogenes* hypervirulence by harnessing its biodiversity. *Nat Genet* 48 (3), 308-313.
26. Mackaness, G.B. (1962) Cellular resistance to infection. *J Exp Med* 116, 381-406.
27. Racz, P. et al. (1972) Experimental *Listeria* enteritis. I. An electron microscopic study of the epithelial phase in experimental listeria infection. *Lab Invest* 26 (6), 694-700.
28. Gaillard, J.L. et al. (1987) *In vitro* model of penetration and intracellular growth of *Listeria monocytogenes* in the human enterocyte-like cell line Caco-2. *Infect Immun* 55 (11), 2822-9.
29. Edman, D.C. et al. (1968) *Listeria monocytogenes* L forms. I. Induction maintenance, and biological characteristics. *J Bacteriol* 96 (2), 352-7.
30. Cantinelli, T. et al. (2013) "Epidemic clones" of *Listeria monocytogenes* are widespread and ancient clonal groups. *J Clin Microbiol* 51 (11), 3770-9.
31. Becavin, C. et al. (2014) Comparison of widely used *Listeria monocytogenes* strains EGD, 10403S, and EGD-e highlights genomic variations underlying differences in pathogenicity. *mBio* 5 (2), e00969-14.
32. Geoffroy, C. et al. (1987) Purification, characterization, and toxicity of the sulfhydryl-activated hemolysin listeriolysin O from *Listeria monocytogenes*. *Infect Immun* 55 (7), 1641-6.
33. Harvey, P.C. and Faber, J.E. (1941) Studies on the *Listerella* Group: I. Biochemical and Hemolytic Reactions. *J Bacteriol* 42 (5), 677-87.
34. Gaillard, J.L. et al. (1986) Transposon mutagenesis as a tool to study the role of hemolysin in the virulence of *Listeria monocytogenes*. *Infect Immun* 52 (1), 50-5.
35. Camilli, A. et al. (1993) Dual roles of plcA in *Listeria monocytogenes* pathogenesis. *Mol Microbiol* 8 (1), 143-57.
36. Domann, E. et al. (1991) Molecular cloning, sequencing, and identification of a metalloprotease gene from *Listeria monocytogenes* that is species specific and physically linked to the listeriolysin gene. *Infect Immun* 59 (1), 65-72.
37. Kocks, C. et al. (1992) *L. monocytogenes*-induced actin assembly requires the actA gene product, a surface protein. *Cell* 68 (3), 521-31.
38. Tilney, L.G. and Portnoy, D.A. (1989) Actin filaments and the growth, movement, and spread of the intracellular bacterial parasite, *Listeria monocytogenes*. *J Cell Biol* 109 (4 Pt 1), 1597-608.
39. Camilli, A. et al. (1991) *Listeria monocytogenes* mutants lacking phosphatidylinositol-specific phospholipase C are avirulent. *J Exp Med* 173 (3), 751-4.
40. Portnoy, D.A. et al. (1988) Role of hemolysin for the intracellular growth of *Listeria monocytogenes*. *J Exp Med* 167 (4), 1459-71.
41. Leimeister-Wachter, M. et al. (1990) Identification of a gene that positively regulates expression of listeriolysin, the major virulence factor of *Listeria monocytogenes*. *Proc Natl Acad Sci U S A* 87 (21), 8336-40.
42. Mengaud, J. et al. (1991) Pleiotropic control of *Listeria monocytogenes* virulence factors by a gene that is autoregulated. *Mol Microbiol* 5 (9), 2273-83.

43. Prokop, A. et al. (2017) OrfX, a Nucleomodulin Required for *Listeria monocytogenes* Virulence. *mBio* 8 (5).
44. de las Heras, A. et al. (2011) Regulation of *Listeria* virulence: PrfA master and commander. *Current opinion in microbiology* 14 (2), 118-27.
45. Reniere, M.L. et al. (2015) Glutathione activates virulence gene expression of an intracellular pathogen. *Nature* 517 (7533), 170-3.
46. Tiensuu, T. et al. (2019) Flick of a switch: regulatory mechanisms allowing *Listeria monocytogenes* to transition from a saprophyte to a killer. *Microbiology (Reading)* 165 (8), 819-833.
47. Gaillard, J.L. et al. (1991) Entry of *L. monocytogenes* into cells is mediated by internalin, a repeat protein reminiscent of surface antigens from gram-positive cocci. *Cell* 65 (7), 1127-41.
48. Mengaud, J. et al. (1996) E-cadherin is the receptor for internalin, a surface protein required for entry of *L. monocytogenes* into epithelial cells. *Cell* 84 (6), 923-32.
49. Lecuit, M. et al. (1999) A single amino acid in E-cadherin responsible for host specificity towards the human pathogen *Listeria monocytogenes*. *EMBO J* 18 (14), 3956-63.
50. Lecuit, M. et al. (2001) A transgenic model for listeriosis: role of internalin in crossing the intestinal barrier. *Science* 292 (5522), 1722-5.
51. Nikitas, G. et al. (2011) Transcytosis of *Listeria monocytogenes* across the intestinal barrier upon specific targeting of goblet cell accessible E-cadherin. *J Exp Med* 208 (11), 2263-77.
52. Kim, M. et al. (2020) Live Imaging Reveals *Listeria* Hijacking of E-Cadherin Recycling as It Crosses the Intestinal Barrier. *Curr Biol*.
53. Dramsi, S. et al. (1995) Entry of *Listeria monocytogenes* into hepatocytes requires expression of InlB, a surface protein of the internalin multigene family. *Mol Microbiol* 16 (2), 251-61.
54. Shen, Y. et al. (2000) InlB-dependent internalization of *Listeria* is mediated by the Met receptor tyrosine kinase. *Cell* 103 (3), 501-10.
55. Khelef, N. et al. (2006) Species specificity of the *Listeria monocytogenes* InlB protein. *Cell Microbiol* 8 (3), 457-70.
56. Disson, O. et al. (2008) Conjugated action of two species-specific invasion proteins for fetoplacental listeriosis. *Nature* 455 (7216), 1114-8.
57. Gessain, G. et al. (2015) PI3-kinase activation is critical for host barrier permissiveness to *Listeria monocytogenes*. *J Exp Med* 212 (2), 165-83.
58. Fraser, C.M. et al. (1995) The minimal gene complement of *Mycoplasma genitalium*. *Science* 270 (5235), 397-403.
59. Glaser, P. et al. (2001) Comparative genomics of *Listeria* species. *Science* 294 (5543), 849-52.
60. Sabet, C. et al. (2005) LPXTG protein InlJ, a newly identified internalin involved in *Listeria monocytogenes* virulence. *Infect Immun* 73 (10), 6912-22.
61. Sabet, C. et al. (2008) The *Listeria monocytogenes* virulence factor InlJ is specifically expressed in vivo and behaves as an adhesin. *Infect Immun* 76 (4), 1368-78.
62. Faralla, C. et al. (2018) *Listeria monocytogenes* InlP interacts with afadin and facilitates basement membrane crossing. *PLoS Pathog* 14 (5), e1007094.
63. Faralla, C. et al. (2016) InlP, a New Virulence Factor with Strong Placental Tropism. *Infect Immun* 84 (12), 3584-3596.
64. Cabanes, D. et al. (2004) Auto, a surface associated autolysin of *Listeria monocytogenes* required for entry into eukaryotic cells and virulence. *Mol Microbiol* 51 (6), 1601-14.

65. Dussurget, O. et al. (2002) *Listeria monocytogenes* bile salt hydrolase is a PrfA-regulated virulence factor involved in the intestinal and hepatic phases of listeriosis. *Mol Microbiol* 45 (4), 1095-106.
66. Seeliger, H.P.R. and Höhne, K. (1979) Serotyping of *Listeria monocytogenes* and related species. *Methods in Microbiology* New York: Academic Press, 33-48.
67. Piffaretti, J.C. et al. (1989) Genetic characterization of clones of the bacterium *Listeria monocytogenes* causing epidemic disease. *Proceedings of the National Academy of Sciences of the United States of America* 86 (10), 3818-22.
68. Rasmussen, O.F. et al. (1995) *Listeria monocytogenes* exists in at least three evolutionary lines: evidence from flagellin, invasive associated protein and listeriolysin O genes. *Microbiology* 141 (Pt 9), 2053-61.
69. Ward, T.J. et al. (2008) Multilocus genotyping assays for single nucleotide polymorphism-based subtyping of *Listeria monocytogenes* isolates. *Appl Environ Microbiol* 74 (24), 7629-42.
70. Ragon, M. et al. (2008) A new perspective on *Listeria monocytogenes* evolution. *PLoS Pathog* 4 (9), e1000146.
71. Jensen, A.K. et al. (2016) Molecular Typing and Epidemiology of Human Listeriosis Cases, Denmark, 2002-2012. *Emerging infectious diseases* 22 (4), 625-33.
72. Chen, Y. et al. (2016) Core Genome Multilocus Sequence Typing for Identification of Globally Distributed Clonal Groups and Differentiation of Outbreak Strains of *Listeria monocytogenes*. *Applied and environmental microbiology* 82 (20), 6258-6272.
73. Ruppitsch, W. et al. (2015) Defining and Evaluating a Core Genome Multilocus Sequence Typing Scheme for Whole-Genome Sequence-Based Typing of *Listeria monocytogenes*. *Journal of clinical microbiology* 53 (9), 2869-76.
74. Pightling, A.W. et al. (2015) The *Listeria monocytogenes* Core-Genome Sequence Typer (LmCGST): a bioinformatic pipeline for molecular characterization with next-generation sequence data. *BMC microbiology* 15, 224.
75. Moura, A. et al. (2016) Whole genome-based population biology and epidemiological surveillance of *Listeria monocytogenes*. *Nat Microbiol* 2, 16185.
76. Moura, A. et al. (2017) Real-Time Whole-Genome Sequencing for Surveillance of *Listeria monocytogenes*, France. *Emerg Infect Dis* 23 (9), 1462-1470.
77. Orsi, R.H. et al. (2011) *Listeria monocytogenes* lineages: Genomics, evolution, ecology, and phenotypic characteristics. *Int J Med Microbiol* 301 (2), 79-96.
78. Seeliger, H.P. and Langer, B. (1989) Serological analysis of the genus *Listeria*. Its values and limitations. *Int J Food Microbiol* 8 (3), 245-8.
79. Doumith, M. et al. (2004) Differentiation of the major *Listeria monocytogenes* serovars by multiplex PCR. *J Clin Microbiol* 42 (8), 3819-22.
80. Chenal-Francisque, V. et al. (2011) Worldwide distribution of major clones of *Listeria monocytogenes*. *Emerg Infect Dis* 17 (6), 1110-2.
81. Haase, J.K. et al. (2014) The ubiquitous nature of *Listeria monocytogenes* clones: a large-scale Multilocus Sequence Typing study. *Environ Microbiol* 16 (2), 405-16.
82. Kwong, J.C. et al. (2016) Prospective Whole-Genome Sequencing Enhances National Surveillance of *Listeria monocytogenes*. *J Clin Microbiol* 54 (2), 333-42.
83. Lee, S. et al. (2018) *Listeria monocytogenes* Source Distribution Analysis Indicates Regional Heterogeneity and Ecological Niche Preference among Serotype 4b Clones. *mBio* 9 (2).
84. Wang, Y. et al. (2012) Genetic diversity and molecular typing of *Listeria monocytogenes* in China. *BMC Microbiol* 12, 119.
85. Bertrand, S. et al. (2016) Diversity of *Listeria monocytogenes* Strains of Clinical and Food Chain Origins in Belgium between 1985 and 2014. *PLoS One* 11 (10), e0164283.

- 581 86. Kuch, A. et al. (2018) Molecular diversity and antimicrobial susceptibility of *Listeria*
582 *monocytogenes* isolates from invasive infections in Poland (1997-2013). Sci Rep 8 (1), 14562.
- 583 87. Painset, A. et al. (2019) LiSEQ - whole-genome sequencing of a cross-sectional survey of
584 *Listeria monocytogenes* in ready-to-eat foods and human clinical cases in Europe. Microb
585 Genom 5 (2).
- 586 88. Huang, Y.T. et al. (2015) Disease Burden of Invasive Listeriosis and Molecular
587 Characterization of Clinical Isolates in Taiwan, 2000-2013. PloS one 10 (11), e0141241.
- 588 89. Wang, Y. et al. (2015) Characterization of *Listeria monocytogenes* isolated from human
589 Listeriosis cases in China. Emerg Microbes Infect 4 (8), e50.
- 590 90. Dreyer, M. et al. (2016) *Listeria monocytogenes* sequence type 1 is predominant in
591 ruminant rhombencephalitis. Scientific reports 6, 36419.
- 592 91. Jonquieres, R. et al. (1998) The *inlA* gene of *Listeria monocytogenes* LO28 harbors a
593 nonsense mutation resulting in release of internalin. Infect Immun 66 (7), 3420-2.
- 594 92. Jacquet, C. et al. (2004) A molecular marker for evaluating the pathogenic potential of
595 foodborne *Listeria monocytogenes*. J Infect Dis 189 (11), 2094-100.
- 596 93. Lecuit, M. et al. (2004) Targeting and crossing of the human maternofetal barrier by
597 *Listeria monocytogenes*: role of internalin interaction with trophoblast E-cadherin. Proc Natl
598 Acad Sci U S A 101 (16), 6152-7.
- 599 94. den Bakker, H.C. et al. (2010) Comparative genomics of the bacterial genus *Listeria*:
600 Genome evolution is characterized by limited gene acquisition and limited gene loss. BMC
601 Genomics 11, 688.
- 602 95. Johnson, J. et al. (2004) Natural atypical *Listeria innocua* strains with *Listeria*
603 *monocytogenes* pathogenicity island 1 genes. Appl Environ Microbiol 70 (7), 4256-66.
- 604 96. Milillo, S.R. et al. (2012) *Listeria monocytogenes* and hemolytic *Listeria innocua* in
605 poultry. Poult Sci 91 (9), 2158-63.
- 606 97. Moreno, L.Z. et al. (2012) Characterization of atypical *Listeria innocua* isolated from
607 swine slaughterhouses and meat markets. Res Microbiol 163 (4), 268-71.
- 608 98. Moura, A. et al. (2019) Atypical Hemolytic *Listeria innocua* Isolates Are Virulent, albeit
609 Less than *Listeria monocytogenes*. Infect Immun 87 (4).
- 610 99. Volokhov, D.V. et al. (2007) The presence of the internalin gene in natural atypically
611 hemolytic *Listeria innocua* strains suggests descent from *L. monocytogenes*. Appl Environ
612 Microbiol 73 (6), 1928-39.
- 613 100. Schmid, M.W. et al. (2005) Evolutionary history of the genus *Listeria* and its virulence
614 genes. Syst Appl Microbiol 28 (1), 1-18.
- 615 101. Maury, M.M. et al. (2017) Spontaneous Loss of Virulence in Natural Populations of
616 *Listeria monocytogenes*. Infect Immun 85 (11).
- 617 102. Cotter, P.D. et al. (2008) Listeriolysin S, a novel peptide haemolysin associated with a
618 subset of lineage I *Listeria monocytogenes*. PLoS Pathog 4 (9), e1000144.
- 619 103. Quereda, J.J. et al. (2016) Bacteriocin from epidemic *Listeria* strains alters the host
620 intestinal microbiota to favor infection. Proc Natl Acad Sci U S A 113 (20), 5706-11.
- 621 104. Quereda, J.J. et al. (2017) Listeriolysin S: A bacteriocin from epidemic *Listeria*
622 *monocytogenes* strains that targets the gut microbiota. Gut Microbes 8 (4), 384-391.
- 623 105. Clayton, E.M. et al. (2014) Atypical *Listeria innocua* strains possess an intact LIPI-3.
624 BMC Microbiol 14, 58.
- 625 106. Wang, Y. et al. (2019) Genomic dissection of the most prevalent *Listeria monocytogenes*
626 clone, sequence type ST87, in China. BMC Genomics 20 (1), 1014.
- 627 107. Yin, Y. et al. (2019) A hybrid sub-lineage of *Listeria monocytogenes* comprising
628 hypervirulent isolates. Nat Commun 10 (1), 4283.
- 629 108. Guillet, C. et al. (2010) Human listeriosis caused by *Listeria ivanovii*. Emerg Infect Dis
630 16 (1), 136-8.

109. Sergeant, E.S. et al. (1991) Abortions in sheep due to *Listeria ivanovii*. Australian veterinary journal 68 (1), 39.
110. Dominguez-Bernal, G. et al. (2006) A spontaneous genomic deletion in *Listeria ivanovii* identifies LIPI-2, a species-specific pathogenicity island encoding sphingomyelinase and numerous internalins. Mol Microbiol 59 (2), 415-32.
111. Fenlon, D.R. (1985) Wild birds and silage as reservoirs of *Listeria* in the agricultural environment. J Appl Bacteriol 59 (6), 537-43.
112. Nightingale, K.K. et al. (2004) Ecology and transmission of *Listeria monocytogenes* infecting ruminants and in the farm environment. Appl Environ Microbiol 70 (8), 4458-67.
113. Weis, J. and Seeliger, H.P. (1975) Incidence of *Listeria monocytogenes* in nature. Appl Microbiol 30 (1), 29-32.
114. Maury, M.M. et al. (2019) Hypervirulent *Listeria monocytogenes* clones' adaption to mammalian gut accounts for their association with dairy products. Nat Commun 10 (1), 2488.
115. Ortiz, S. et al. (2016) The Connection between Persistent, Disinfectant-Resistant *Listeria monocytogenes* Strains from Two Geographically Separate Iberian Pork Processing Plants: Evidence from Comparative Genome Analysis. Appl Environ Microbiol 82 (1), 308-17.
116. Harter, E. et al. (2017) Stress Survival Islet 2, Predominantly Present in *Listeria monocytogenes* Strains of Sequence Type 121, Is Involved in the Alkaline and Oxidative Stress Responses. Appl Environ Microbiol 83 (16).
117. Ryan, S. et al. (2010) A five-gene stress survival islet (SSI-1) that contributes to the growth of *Listeria monocytogenes* in suboptimal conditions. J Appl Microbiol 109 (3), 984-95.
118. Hingston, P. et al. (2017) Genotypes Associated with *Listeria monocytogenes* Isolates Displaying Impaired or Enhanced Tolerances to Cold, Salt, Acid, or Desiccation Stress. Front Microbiol 8, 369.
119. Esteban, J.I. et al. (2009) Faecal shedding and strain diversity of *Listeria monocytogenes* in healthy ruminants and swine in Northern Spain. BMC Vet Res 5, 2.
120. Toledo-Arana, A. et al. (2009) The *Listeria* transcriptional landscape from saprophytism to virulence. Nature 459 (7249), 950-6.

Outstanding questions

- What are the natural environments that have led to the current biodiversity of *L. monocytogenes*?
- What is the primary function of LIPI-1 genes, and other so-called “virulence genes”?
- What are the driving forces that have selected the virulence genes of *L. monocytogenes*?
- How does *L. monocytogenes* maintain its virulence since disease is not implicated in inter-host transmission?

(A)

LIPI-1
Lm, *Liv* and
atypical *Lin*

LIPI-2
Liv and
atypical *Lm* CC33*

LIPI-3
mainly *Lm* CC1,
CC4, CC6 and
some *Lin*

LIPI-4
mainly *Lm* CC4
and *Lin*

Scale: 1 Kbp

(B)

Figure 3

Host-association

Virulence

Active PrfA

Active LIPIs

Functional InlA

Fecal shedding

Transmission

Virulence

Immunodeficiency

Listeriosis depends on
microbial and host factors

Saprophytism

Loss of virulence

Inactive PrfA

Loss of LIPI-1 regulation

Truncation of InlA

Biofilm formation

Resistance to antiseptics