



**HAL**  
open science

## **Editorial: Role of the IL-23/IL-17 Pathway in Chronic Immune-Mediated Inflammatory Diseases: Mechanisms and Targeted Therapies**

Elisabetta Bianchi, Matteo Vecellio, Lars Rogge

► **To cite this version:**

Elisabetta Bianchi, Matteo Vecellio, Lars Rogge. Editorial: Role of the IL-23/IL-17 Pathway in Chronic Immune-Mediated Inflammatory Diseases: Mechanisms and Targeted Therapies. *Frontiers in Immunology*, 2021, 12, pp.770275. 10.3389/fimmu.2021.770275 . pasteur-03497758

**HAL Id: pasteur-03497758**

**<https://hal-pasteur.archives-ouvertes.fr/pasteur-03497758>**

Submitted on 20 Dec 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution| 4.0 International License



# Editorial: Role of the IL-23/IL-17 Pathway in Chronic Immune-Mediated Inflammatory Diseases: Mechanisms and Targeted Therapies

Elisabetta Bianchi<sup>1,2\*</sup>, Matteo Vecellio<sup>3,4</sup> and Lars Rogge<sup>1,2</sup>

<sup>1</sup> Immunoregulation Unit, Department of Immunology, Institut Pasteur, Paris, France, <sup>2</sup> Unité Mixte de Recherche, Institut Pasteur/AP-HP Hôpital Cochin, Paris, France, <sup>3</sup> Nuffield Department of Orthopaedics, Rheumatology and Musculoskeletal Sciences, University of Oxford, Oxford, United Kingdom, <sup>4</sup> Division of Rheumatology and Clinical Immunology, Humanitas Clinical and Research Center, IRCCS, Milan, Italy

**Keywords:** chronic inflammatory diseases, spondyloarthritis, interleukin-23, interleukin-17, psoriasis

## Editorial on the Research Topic

### Role of the IL-23/IL-17 Pathway in Chronic Immune-Mediated Inflammatory Diseases: Mechanisms and Targeted Therapies

Chronic inflammatory diseases (CID) are clinically heterogeneous conditions that share common inflammatory pathways and derive from aberrant immune responses. The implication of the interleukin-23/interleukin-17 (IL-23/IL-17) axis in several CID is supported by studies in animal models of autoimmune disease (1, 2) and by the genome-wide association studies (GWAS) finding that several of the non-MHC loci genetically linked to Crohn's disease, psoriasis, and axial spondyloarthritis (axSpA), are associated with genes in this pathway (*IL23R*, *IL12B*, *IL6R*, *IL1R2*, *RORC*, *RUNX3*, *TYK2*, *JAK2*, *CARD9*) (3–5).

The clinical relevance of the IL-23/IL-17 axis has been validated by the successful treatment of psoriasis, psoriatic arthritis (PsA) and axSpA with IL-17A inhibitors (6–8). Furthermore, targeting IL-23 has proven highly effective for the treatment of psoriasis, and beneficial in PsA (9, 10), a disease belonging to the SpA spectrum. However, the clinical studies using these drugs have also given unexpected results, dissociating the effectiveness of IL-17 from IL-23 inhibitors in different diseases (11). A recent phase 2 study testing the IL-23 inhibitor risankizumab did not show any clinical improvement compared to placebo in patients with active axial SpA (12), despite the strong GWAS association of *IL23R* with SpA (4). Conversely, targeting IL-23 has proven effective for the treatment of Crohn's disease, while IL-17 inhibition induced worsening of symptoms in this disease (13).

These findings demonstrate our limited understanding of the pathogenic mechanisms of IL-17 and IL-23 in these CIDs, suggesting the need to reassess the link between IL-23 and IL-17 in these diseases.

This Research Topic offers an overview of the impact of the IL-23/IL-17 pathways in CIDs, in particular SpA, with a focus on the mechanisms driving pathogenesis and response to therapy.

IL-23 is important for the expansion and the functional activity of T helper 17 (Th17) cells, which secrete the pro-inflammatory cytokine IL-17 (14), but it may also act on several populations of innate immune cells that express the IL-23 receptor (IL-23R), including innate lymphoid cells (ILC),  $\gamma\delta$  T lymphocytes, iNKT cells, mucosal-associated invariant T cells (MAIT), and, neutrophils (15–20).

## OPEN ACCESS

### Edited and reviewed by:

Silvano Sozzani,  
Sapienza University of Rome, Italy

### \*Correspondence:

Elisabetta Bianchi  
elisabetta.bianchi@pasteur.fr

### Specialty section:

This article was submitted to  
Cytokines and Soluble  
Mediators in Immunity,  
a section of the journal  
Frontiers in Immunology

**Received:** 03 September 2021

**Accepted:** 08 September 2021

**Published:** 23 September 2021

### Citation:

Bianchi E, Vecellio M and Rogge L  
(2021) Editorial: Role of the IL-23/IL-17  
Pathway in Chronic Immune-Mediated  
Inflammatory Diseases: Mechanisms  
and Targeted Therapies.  
*Front. Immunol.* 12:770275.  
doi: 10.3389/fimmu.2021.770275

Some of these populations have been found to accumulate in the diseased tissues of patients or of model animals (21), suggesting that the inflammatory response in CID may be the result of a complex interplay of different immune cell types whose relative role in the pathogenesis of specific CIDs remains to be defined. Rosine and Miceli-Richard provide a comprehensive overview of IL-17 producing innate cell subsets in the context of SpA pathogenesis, while McGinty et al. propose that the immunoregulatory function of Tr1 cells may be impaired in SpA. IL-10 production by Tr1 was shown to prevent gut inflammation, and IL-23 downregulates IL-10 secretion in these cells (22). Given the therapeutic potential of these cells, the future challenge is the development of appropriate pre-clinical models to explore the role of Tr1 cells in CIDs.

Another cytokine regulated by IL-23 is IL-22, which is produced by Th17 cells, among other cell subsets (23). Lindhal and Olson explore in detail the role of IL-22 in Th1/Th17 cell polarization and in CIDs. Although the different studies are not always consistent, IL-22 seem in most models to reduce Th1 responses and may contribute to resolve inflammation by inducing IL-10 production.

Th17 differentiation is regulated at the transcriptional level by the IRF4 transcription factor (24). Using a T cell transfer model of colitis, Buchele et al. demonstrate that IRF4 also controls Th17 pathogenic function indirectly, by acting in a conventional Dendritic Cell 2 (cDC2) subset. This work highlights the role of IRF4 as a molecular switch that controls Th17 differentiation, as well as the importance of the cDC2 subset in the pathogenesis of colitis.

Adding complexity to the regulation of Th17 function, Peng et al. demonstrate a post-transcriptional mechanism that controls IL-17-mediated inflammation. The authors have shown that Tristetraprolin (TTP), an RNA-binding protein, inhibits IL-23 expression. In the present work, Peng et al. show that TTP conditional KO (CD4CreTTP<sup>fl/fl</sup>) mice displayed increased systemic IL-17A and skin Th17 cells, and increased susceptibility to DSS-induced colitis. These data indicate that TTP is an important regulator of inflammation and a potential new therapeutic target.

Animal models of CID have been crucial to improve our understanding of the molecular processes that drive CIDs, as comprehensively illustrated by Mandour et al., in particular for diseases such as axSpA, for which access to human diseased tissue is difficult. Rodent models for SpA have been useful to study the molecular mechanisms of IL-23 induced pathogenesis, despite the fact that none reflects the whole range of pathologic findings of this disease. The study of these models has highlighted the possible role of IL-23-dependent gut and skin inflammation in triggering joint pathology. Another interesting finding of these studies is the importance of IL-23 in the early phases of SpA pathogenesis, demonstrated by the ability of IL-23 blockers to prevent disease onset when administered before the development of symptoms. The study of early events in these models may help develop predictive tools and identify targets for early therapeutic intervention. Whether IL-23 plays a similar role in the pre-clinical phase of the disease in humans remains to be established.

In human studies, GWAS have proven very useful to indicate potential pathogenic pathways in CIDs.

Disease-associated genetic variants may have the power to discriminate between similar conditions, such as psoriasis, PsA and ankylosing spondylitis (AS). In their article, Vecellio et al. highlighted the contribution of the IL-17/IL-23 axis to PsA, a disorder sharing most of the genetics and molecular mechanisms with other inflammatory diseases, like psoriasis, AS, inflammatory bowel disease and Behçet disease. The association of loci in the IL-17/IL-23 axis is the *usual suspect* that characterizes these disorders, together with the contribution of Th17 lymphocytes. The development of biologics blocking IL-17, such as Secukinumab in AS, or IL-23 such as Ustekinumab in psoriasis/PsA, demonstrates the value of a combination of genetic markers as an approach to identify credible targets for treatment. Wordsworth et al. elegantly summarize the progress made in the last years by the research community to identify candidate genes that contribute to increased AS susceptibility. More than 100 loci have been found to be associated with increased AS risk, but this could be an underestimate: it is crucial to have large cohorts and bigger sample size to increase the power of these studies. The authors point out that still no reliable genetic predictors of disease severity in AS or response to treatment are available, despite the efforts of the scientific community. The identification of credible therapeutic targets and the translation of genome wide association studies (GWAS) findings in AS, is the main message from Zaroor et al.'s contribution. Despite recent progress, several challenges are still present in order to predict which are the causal genes regulated by disease-associated genetic variants and to define the relevant cell-type where these SNPs act. The success of biologics targeting the IL-17/IL-23 axis highlights the value of genetic studies for drug development. However, since it's very unlikely that two patients will have the same genetic makeup, stratification based on genetic predictors remains challenging.

Schinocca et al. summarize recent findings in human and animal models supporting the role of the IL-23/IL-17 pathway in SpA and other rheumatic diseases, including Rheumatoid Arthritis, Sjögren Syndrome and Systemic Lupus Erythematosus, linking molecular pathology to the development of biologic therapies. Several novel biologics targeting the IL-23/IL-17 pathway (including IL-17F and the Janus kinases (JAK) downstream of IL-23R) are being currently tested, as detailed by Ceribelli et al. in their comprehensive review of ongoing clinical trials for SpA treatment.

Hammitzsch et al. focus their attention on the role of JAKs in SpA pathogenesis, and the development of inhibitors for treatment. Given the pleiotropic role of a JAK in multiple signaling pathways, they argue for the use of selective inhibitors, to avoid undesired alterations of bone homeostasis.

In a bedside to bench approach, Fiechter et al. interrogate the effect of IL-23 inhibition on molecular pathways and cellular populations in the synovia of PsA patients. Major pathways modulated by treatment were MAPK/ERK, mTOR and Wnt signaling, while IL-17A production was not significantly affected, supporting a non-linearity in the IL-23/IL-17 pathway, and the

possibility of other pathogenic targets downstream of IL-23. Changes induced by treatment in the Wnt pathway also indicate the importance of better investigating the effects of IL-23 on bone metabolism.

Liu et al. explore the role of IL-23 and the effects of its inhibition in a wide range of inflammatory skin diseases. Their analysis suggests that IL-23 may be important for the development of several skin diseases, including Hidradenitis Suppurativa or Pityriasis Rubra, which show clinical improvement upon IL-23 blockade. Bugaut and Aractingi focus on the pathogenesis and treatment of psoriasis, with an eye to new therapeutics inhibiting selective JAKs and the transcription factor ROR $\gamma$ t, which is essential for the function of Th17 cells. The authors underline how improved understanding of IL-23/IL-17 biology, and of the many cell types involved, may lead to the identification of new therapeutic targets, necessary for severe and refractory cases.

The role of the IL-23/IL-17 axis in inflammatory bowel disease (IBD) is reviewed by Schmitt et al. and by Noviello et al. An interesting concept that emerges from this overview is the importance of considering T cell plasticity and changes in immune profiles during disease progression, which may explain the need for a different biological treatment at different stages of disease. Noviello et al. also suggest the possibility of stratifying patients for treatment according to baseline cytokine levels.

Finally, Baeten and Adamopoulos and McGonagle et al. discuss potential reasons why IL-23-inhibition failed in AS. Revisiting the scientific rationale for conducting trials of IL-23-inhibitors in AS, Baeten and Adamopoulos caution that the evidence supporting a central role of IL-23 in the pathobiology of AxSpA was circumstantial, at best. In particular, they state that systemic IL-23 exposure induced chronic arthritis, severe bone loss and myelopoiesis in the bone marrow and spleen of mice (25), a phenotype which is not compatible with AxSpA. This report clearly contrasts the more publicized observation that

IL-23 overexpression induces a SpA-like phenotype in mice (26). This perspective concludes with the notion that the IL-23/IL-17 axis is not a linear “cascade” and that genetic data are an excellent tool to generate hypotheses, but are not sufficient to prove or disprove them. McGonagle et al. argue that IL-23 blockade can prevent disease onset but not established disease in an experimental SpA model (27). Even if it is currently not clear if these observations can be translated to human disease, they point to a role of IL-23 in disease initiation, while persistent disease may be maintained by IL-23-independent IL-17 production by memory T cells. IL-23-independent IL-17 production by various lymphocyte populations is discussed in several reviews in this topic. McGonagle et al. point out that there is heterogeneity within human  $\gamma\delta$  T cell populations with respect to IL-23 receptor expression. Both  $\delta$ 1 and  $\delta$ 2  $\gamma\delta$  T cell populations express IL-17A following stimulation, however only the  $\delta$ 2 population further upregulated IL-17A production when stimulated in the presence of IL-23 (16). An interesting point also discussed by McGonagle et al. is that patients with active PsA and imaging-confirmed sacroiliitis (axial PsA) benefit from treatment with the anti-IL-12/23 inhibitor ustekinumab (28) and the IL-23 inhibitor guselkumab (29). Thus, a subgroup of SpA patients with axial inflammation may actually benefit from IL-23-blockade, and McGonagle et al. propose that adequate IL-23 dosology may be critical in this condition.

In conclusion, the articles included in this collection reached their “primary endpoint”, that is raising more questions to guide future clinical and fundamental research.

## AUTHOR CONTRIBUTIONS

All authors listed have made a substantial, direct, and intellectual contribution to the work and approved it for publication.

## REFERENCES

- Cua DJ, Sherlock J, Chen Y, Murphy CA, Joyce B, Seymour B, et al. Interleukin-23 Rather Than Interleukin-12 is the Critical Cytokine for Autoimmune Inflammation of the Brain. *Nature* (2003) 421(6924):744–8. doi: 10.1038/nature01355
- Langrish CL, Chen Y, Blumenschein WM, Mattson J, Basham B, Sedgwick JD, et al. IL-23 Drives a Pathogenic T Cell Population That Induces Autoimmune Inflammation. *J Exp Med* (2005) 201(2):233–40. doi: 10.1084/jem.20041257
- Bianchi E, Rogge L. The IL-23/IL-17 Pathway in Human Chronic Inflammatory Diseases—New Insight From Genetics and Targeted Therapies. *Genes Immun* (2019) 20(5):415–25. doi: 10.1038/s41435-019-0067-y
- Brown MA, Kenna T, Wordsworth BP. Genetics of Ankylosing Spondylitis—Insights Into Pathogenesis. *Nat Rev Rheumatol* (2016) 12(2):81–91. doi: 10.1038/nrrheum.2015.133
- Ellinghaus D, Jostins L, Spain SL, Cortes A, Bethune J, Han B, et al. Analysis of Five Chronic Inflammatory Diseases Identifies 27 New Associations and Highlights Disease-Specific Patterns at Shared Loci. *Nat Genet* (2016) 48(5):510–8. doi: 10.1038/ng.3528
- Baeten D, Sieper J, Braun J, Baraliakos X, Dougados M, Emery P, et al. Secukinumab, an Interleukin-17a Inhibitor, in Ankylosing Spondylitis. *N Engl J Med* (2015) 373(26):2534–48. doi: 10.1056/NEJMoa1505066
- Langley RG, Elewski BE, Lebwohl M, Reich K, Griffiths CE, Papp K, et al. Secukinumab in Plaque Psoriasis—Results of Two Phase 3 Trials. *N Engl J Med* (2014) 371(4):326–38. doi: 10.1056/NEJMoa1314258
- Mease PJ, McInnes IB, Kirkham B, Kavanaugh A, Rahman P, van der Heijde D, et al. Secukinumab Inhibition of Interleukin-17A in Patients With Psoriatic Arthritis. *N Engl J Med* (2015) 373(14):1329–39. doi: 10.1056/NEJMoa1412679
- Mease PJ, Rahman P, Gottlieb AB, Kollmeier AP, Hsia EC, Xu XL, et al. Guselkumab in Biologic-Naive Patients With Active Psoriatic Arthritis (DISCOVER-2): A Double-Blind, Randomised, Placebo-Controlled Phase 3 Trial. *Lancet* (2020) 395(10230):1126–36. doi: 10.1016/S0140-6736(20)30263-4
- Reich K, Armstrong AW, Langley RG, Flavin S, Randazzo B, Li S, et al. Guselkumab Versus Secukinumab for the Treatment of Moderate-to-Severe Psoriasis (ECLIPSE): Results From a Phase 3, Randomised Controlled Trial. *Lancet* (2019) 394(10201):831–9. doi: 10.1016/S0140-6736(19)31773-8
- Siebert S, Millar NL, McInnes IB. Why did IL-23p19 Inhibition Fail in AS: A Tale of Tissues, Trials or Translation? *Ann Rheum Dis* (2019) 78(8):1015–8. doi: 10.1136/annrheumdis-2018-213654
- Baeten D, Ostergaard M, Wei JC, Sieper J, Jarvinen P, Tam LS, et al. Risankizumab, an IL-23 Inhibitor, for Ankylosing Spondylitis: Results of a Randomised, Double-Blind, Placebo-Controlled, Proof-of-Concept, Dose-Finding Phase 2 Study. *Ann Rheum Dis* (2018) 77(9):1295–302. doi: 10.1136/annrheumdis-2018-213328

13. Fragoulis GE, Siebert S, McInnes IB. Therapeutic Targeting of IL-17 and IL-23 Cytokines in Immune-Mediated Diseases. *Annu Rev Med* (2016) 67:337–53. doi: 10.1146/annurev-med-051914-021944
14. McGeachy MJ, Chen Y, Tato CM, Laurence A, Joyce-Shaikh B, Blumenschein WM, et al. The Interleukin 23 Receptor is Essential for the Terminal Differentiation of Interleukin 17-Producing Effector T Helper Cells *In Vivo*. *Nat Immunol* (2009) 10(3):314–24. doi: 10.1038/ni.1698
15. Cuthbert RJ, Fragkakis EM, Dunsmuir R, Li Z, Coles M, Marzo-Ortega H, et al. Brief Report: Group 3 Innate Lymphoid Cells in Human Enthesis. *Arthritis Rheumatol* (2017) 69(9):1816–22. doi: 10.1002/art.40150
16. Cuthbert RJ, Watad A, Fragkakis EM, Dunsmuir R, Loughenbury P, Khan A, et al. Evidence That Tissue Resident Human Enthesis gammadeltaT-Cells can Produce IL-17A Independently of IL-23R Transcript Expression. *Ann Rheum Dis* (2019) 78(11):1559–65. doi: 10.1136/annrheumdis-2019-215210
17. Gracey E, Qaiyum Z, Almaghouth I, Lawson D, Karki S, Avvaru N, et al. IL-7 Primes IL-17 in Mucosal-Associated Invariant T (MAIT) Cells, Which Contribute to the Th17-Axis in Ankylosing Spondylitis. *Ann Rheum Dis* (2016) 75(12):2124–32. doi: 10.1136/annrheumdis-2015-208902
18. Taylor PR, Roy S, Leal SM Jr., Sun Y, Howell SJ, Cobb BA, et al. Activation of Neutrophils by Autocrine IL-17a-IL-17RC Interactions During Fungal Infection is Regulated by IL-6, IL-23, RORgammat and Dectin-2. *Nat Immunol* (2014) 15(2):143–51. doi: 10.1038/ni.2797
19. Venken K, Jacques P, Mortier C, Labadia ME, Decruy T, Coudensy J, et al. RORgammat Inhibition Selectively Targets IL-17 Producing iNKT and Gammadelta-T Cells Enriched in Spondyloarthritis Patients. *Nat Commun* (2019) 10(1):9. doi: 10.1038/s41467-018-07911-6
20. Watad A, Rowe H, Russell T, Zhou Q, Anderson LK, Khan A, et al. Normal Human Enthesis Harbours Conventional CD4+ and CD8+ T Cells With Regulatory Features and Inducible IL-17A and TNF Expression. *Ann Rheum Dis* (2020) 79(8):1044–54. doi: 10.1136/annrheumdis-2020-217309
21. Bridgewood C, Sharif K, Sherlock J, Watad A, McGonagle D. Interleukin-23 Pathway at the Enthesis: The Emerging Story of Enthesitis in Spondyloarthropathy. *Immunol Rev* (2020) 294(1):27–47. doi: 10.1111/imr.12840
22. Alfen JS, Larghi P, Facciotti F, Gagliani N, Bosotti R, Paroni M, et al. Intestinal IFN-Gamma-Producing Type 1 Regulatory T Cells Coexpress CCR5 and Programmed Cell Death Protein 1 and Downregulate IL-10 in the Inflamed Guts of Patients With Inflammatory Bowel Disease. *J Allergy Clin Immunol* (2018) 142(5):1537–47.e8. doi: 10.1016/j.jaci.2017.12.984
23. Liang SC, Tan XY, Luxenberg DP, Karim R, Dunussi-Joannopoulos K, Collins M, et al. Interleukin (IL)-22 and IL-17 are Coexpressed by Th17 Cells and Cooperatively Enhance Expression of Antimicrobial Peptides. *J Exp Med* (2006) 203(10):2271–9. doi: 10.1084/jem.20061308
24. Brustle A, Heink S, Huber M, Rosenplanter C, Stadelmann C, Yu P, et al. The Development of Inflammatory T(H)-17 Cells Requires Interferon-Regulatory Factor 4. *Nat Immunol* (2007) 8(9):958–66. doi: 10.1038/ni1500
25. Adamopoulos IE, Tessmer M, Chao CC, Adda S, Gorman D, Petro M, et al. IL-23 is Critical for Induction of Arthritis, Osteoclast Formation, and Maintenance of Bone Mass. *J Immunol* (2011) 187(2):951–9. doi: 10.4049/jimmunol.1003986
26. Sherlock JP, Joyce-Shaikh B, Turner SP, Chao CC, Sathe M, Grein J, et al. IL-23 Induces Spondyloarthropathy by Acting on ROR-Gammat(+) CD3(+) CD4(-)CD8(-) Enthesial Resident T Cells. *Nat Med* (2012) 18(7):1069–76. doi: 10.1038/nm.2817
27. van Tok MN, Na S, Lao CR, Alvi M, Pots D, van de Sande MGH, et al. The Initiation, But Not the Persistence, of Experimental Spondyloarthritis Is Dependent on Interleukin-23 Signaling. *Front Immunol* (2018) 9:1550. doi: 10.3389/fimmu.2018.01550
28. Kavanaugh A, Puig L, Gottlieb AB, Ritchlin C, You Y, Li S, et al. Efficacy and Safety of Ustekinumab in Psoriatic Arthritis Patients With Peripheral Arthritis and Physician-Reported Spondylitis: Post-Hoc Analyses From Two Phase III, Multicentre, Double-Blind, Placebo-Controlled Studies (PSUMMIT-1/PSUMMIT-2). *Ann Rheum Dis* (2016) 75(11):1984–8. doi: 10.1136/annrheumdis-2015-209068
29. Mease PJ, Helliwell PS, Gladman DD, Poddubnyy D, Baraliakos X, Chakravarty SD, et al. Efficacy of Guselkumab on Axial Involvement in Patients With Active Psoriatic Arthritis and Sacroiliitis: A Post-Hoc Analysis of the Phase 3 DISCOVER-1 and DISCOVER-2 Studies. *Lancet Rheumatol* (2021). doi: 10.1016/S2665-9913(21)00105-3

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

**Publisher's Note:** All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Bianchi, Vecellio and Rogge. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.