



HAL
open science

Spindle-shaped archaeal viruses evolved from rod-shaped ancestors to package a larger genome

Fengbin Wang, Virginija Cvirkaite-Krupovic, Matthijn Vos, Leticia Beltran, Mark A.B. Kreutzberger, Jean-Marie Winter, Zhangli Su, Jun Liu, Stefan Schouten, Mart Krupovic, et al.

► To cite this version:

Fengbin Wang, Virginija Cvirkaite-Krupovic, Matthijn Vos, Leticia Beltran, Mark A.B. Kreutzberger, et al.. Spindle-shaped archaeal viruses evolved from rod-shaped ancestors to package a larger genome. *Cell*, 2022, 185 (8), pp.1297-1307.e11. 10.1016/j.cell.2022.02.019 . pasteur-03620791

HAL Id: pasteur-03620791

<https://pasteur.hal.science/pasteur-03620791>

Submitted on 26 Mar 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30

Spindle-shaped archaeal viruses
evolved from rod-shaped ancestors
to package a larger genome

Fengbin Wang^{1,†}, Virginija Cvirkaite-Krupovic^{2,†}, Matthijn Vos³, Leticia C. Beltran¹, Mark A.B. Kreutzberger¹, Jean-Marie Winter³, Zhangli Su¹, Jun Liu⁴, Stefan Schouten⁵, Mart Krupovic^{2,*} and Edward H. Egelman^{1,*,a}

¹Department of Biochemistry and Molecular Genetics
University of Virginia
Charlottesville, VA 22903, U.S.A.

²Institut Pasteur, Université de Paris
CNRS UMR6047, Archaeal Virology Unit, 75015 Paris, France

³NanoImaging Core Facility, Centre de Ressources et Recherches Technologiques (C2RT),
Institut Pasteur, 75015 Paris, France

⁴Department of Microbial Pathogenesis & Microbial Sciences Institute,
Yale University School of Medicine
New Haven, CT 06510, U.S.A.

⁵NIOZ Royal Netherlands Institute for Sea Research
Department of Marine Microbiology and Biogeochemistry
Texel, The Netherlands

† These authors contributed equally to this paper

^a Lead Contact

* Correspondence to mart.krupovic@pasteur.fr, egelman@virginia.edu

31 **Summary**

32 Spindle- or lemon-shaped viruses infect archaea in diverse environments. Due to the highly
33 pleomorphic nature of these virions, which can be found with cylindrical tails emanating from
34 the spindle-shaped body, structural studies of these capsids have been challenging. We have
35 determined the atomic structure of the capsid of Sulfolobus monocaudavirus 1, a virus that
36 infects hosts living in nearly boiling acid. A highly hydrophobic protein, likely integrated into the
37 host membrane before the virions assemble, forms seven strands that slide past each other in
38 both the tails and the spindle body. We observe the discrete steps that occur as the tail tubes
39 expand, and these are due to highly conserved quasi-equivalent interactions with neighboring
40 subunits maintained despite significant diameter changes. Our results show how helical
41 assemblies can vary their diameters, becoming nearly spherical to package a larger genome,
42 and suggest how all spindle-shaped viruses have evolved from archaeal rod-like viruses.

43

44

45 **Introduction**

46 Capsids are a hallmark of viruses, which distinguishes them from all other types of
47 mobile genetic elements (Forterre et al., 2014; Raoult and Forterre, 2008). Over billions of
48 years, viruses have ‘invented’ capsids on multiple independent occasions from non-homologous
49 and structurally unrelated proteins (Krupovic et al., 2019). Nevertheless, the majority of known
50 viruses package their genomes into icosahedral or filamentous helical protein capsids (also
51 called nucleocapsids if surrounded by an additional layer, e.g., lipid membrane) (Krupovic and
52 Koonin, 2017; Sevvana et al., 2021). Viruses infecting archaea notoriously deviate from this
53 general paradigm by producing virions with unique, odd-shaped morphologies, which are not
54 observed among bacterial or eukaryotic viruses (Dellas et al., 2014; Prangishvili et al., 2017).
55 Such archaea-specific virion architectures resemble droplets, champagne bottles or spindles
56 (Baquero et al., 2020). Viruses with spindle-shaped (or lemon-shaped) virions are particularly
57 common in diverse extreme and moderate environments and infect a wide range of archaeal
58 lineages from the phyla Crenarchaeota, Euryarchaeota and Thaumarchaeota as well as
59 Asgardarchaeota, a group of archaea widely considered to represent the closest archaeal
60 relatives of eukaryotes (Medvedeva et al., 2021). Because of the broad distribution in Archaea,
61 it has been suggested that spindle-shaped viruses were associated with the last archaeal
62 common ancestor, and possibly, even the last universal cellular ancestor (LUCA) (Krupovic et al.,
63 2020).

64 Based on virion characteristics and genomic relationships, most spindle-shaped viruses
65 fall into two groups (Krupovic et al., 2014). One of the groups comprises smaller spindle-shaped
66 viruses classified into families *Fuselloviridae* (e.g., *Sulfolobus* spindle-shaped virus 1 [SSV1]),

67 *Halspiviridae* (e.g., haloarchaeal virus His1) and *Thaspiviridae* (e.g., Nitrosopumilus spindle-
68 shaped virus 1 [NSV1]) as well as several unclassified viruses. These viruses encode homologous
69 major capsid proteins (MCP) containing two hydrophobic, potentially membrane-spanning
70 domains and are extruded from the host cell through a budding-like mechanism without
71 causing cell lysis (Kim et al., 2019; Quemin et al., 2016). Attempts to determine the structures
72 of SSV1 (Stedman et al., 2015) and His1 (Hong et al., 2015) by cryo-EM did not yield atomic
73 models and provided little insight into virion organization. The second group includes members
74 of the family *Bicaudaviridae*, which have larger virions and genomes. A characteristic feature of
75 this virus assemblage is the presence of ‘tails’ emanating from one or both pointed ends of the
76 spindle. Notably, *Acidianus* two-tailed virus (ATV) and *Sulfolobus* monocaudavirus 1 (SMV1)
77 have been reported to develop tails extracellularly, with the only requirement for this
78 transformation being the incubation at temperatures close to those of the natural habitats (75-
79 90°C) (Haring et al., 2005; Uldahl et al., 2016). It has been suggested that virus-encoded MoxR-
80 type AAA+ ATPase and a von Willebrand domain A-containing cochaperone play an active role
81 in this process by hydrolyzing ATP stored within the virions (Scheele et al., 2011). However, the
82 exact mechanism underlying this virion morphogenesis outside of the host cells has not been
83 elucidated. Other bicaudaviruses, namely, *Acidianus* tailed spindle virus (ATSV) and *Sulfolobus*
84 *tengchongensis* spindle-shaped viruses 1 and 2 (STSV1 and STSV2, respectively) are released
85 from the cells with tails extending from one of the virion poles and do not undergo further
86 transformation (Erdmann et al., 2014; Hochstein et al., 2016; Xiang et al., 2005). Remarkably,
87 upon infection, STSV2 and SMV1 block normal cell division, transforming the host cell into a
88 giant virion-producing factory, which is up to 20 times larger compared to non-infected cells

89 (Liu et al., 2021). The soluble structural protein repeatedly identified as the major component
90 of bicaudavirus virions (Erdmann *et al.*, 2014; Hochstein *et al.*, 2016; Prangishvili et al., 2006;
91 Xiang *et al.*, 2005) is unrelated to that of the smaller spindle-shaped viruses from the first
92 group, leading to the suggestion that the two groups of viruses are evolutionarily distinct
93 (Krupovic *et al.*, 2014). The protein has been crystallized for ATV and ATSV and shows a four-
94 helix bundle fold (Goulet et al., 2010; Hochstein et al., 2018).

95 Filamentous and spherical capsids (displaying helical and icosahedral symmetries,
96 respectively) embody the simplest virion designs dictated by the principle of ‘genetic economy’
97 inherent to viral genomes, whereby relatively large capsids are built from a small number of
98 distinct protein subunits interacting with each other in a way that is identical throughout the
99 capsid (Crick and Watson, 1956). For icosahedral viruses, Caspar and Klug proposed that
100 interactions between the capsomers are quasi-equivalent (rather than identical) and developed
101 the quasi-equivalence theory which allows predicting the arrangement of capsomers in larger
102 icosahedral capsids (Caspar and Klug, 1962). How complex capsids, such as those of spindle-
103 shaped viruses, are organized and how they have evolved remains unclear. While two
104 theoretical models have been postulated for the geometric basis of the spindle shape in
105 archaeal viruses (Perotti et al., 2016; Perotti et al., 2019), we show that they bear no relation to
106 the actual structure of these viruses. Similarly, we can show that the crystal structures for the
107 putative MCPs in both ATV (Goulet *et al.*, 2010) and ATSV (Hochstein *et al.*, 2018) were not of
108 the MCPs, but unrelated viral proteins. Since spindle-shaped viruses appear to have evolved
109 from archaeal rod-like viruses, our results explain why spindle-shaped viruses have thus far only
110 been observed to infect archaea.

111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132

Results

Cryo-EM images of SMV1 (Figure 1A) show that the spindle-shaped body of the virion is continuous with the tubular tails extending out from the body. As observed for ATSV (Hochstein *et al.*, 2018), the periodicity in the spindle body is the same as that in the tails, strongly suggesting a seamless transition between the tails and the spindles. While this periodicity was described as ~ 40 Å for ATSV (Hochstein *et al.*, 2018), we found a 29 Å periodicity in SMV1 (Figure 1A). The presence or absence of tails can be modulated by prolonged incubation at high temperatures (Haring *et al.*, 2005; Uldahl *et al.*, 2016). However, the heat-treated SMV1 population remained very heterogeneous with tails displaying variable lengths and diameters. We noticed that increase in pH from 6 to 10 results in nearly uniform transformation of spindle-shaped virions into tubular structures and is accompanied by the release of material present inside of the viral particle (Figure S1A). Fractionation of the SMV1 particles on 5-20% sucrose gradient followed by analysis of the different gradient fractions showed that the viral genome was released in the form of amorphous aggregates along with proteins present in the lumen of the viral particles, whereas tubular structures were largely devoid of DNA (Figure S1B). Thus, given that the tubular tails must have helical symmetry, and understanding the tails would reveal the structure of the more complicated spindle body, most of the structural studies in this paper were performed on the capsids which were transformed into a tubular state (see Methods). Nevertheless, extensive analysis of millions of segments revealed a nearly continuous variation in the diameter of the tail tubes. Progress was only made by starting from a dataset containing six million particles extracted from 16,401 micrographs, allowing us to use

133 multiple 2D classification cycles to find nearly homogeneous subsets of constant diameter. This
134 classification approach showed that rather than being continuously variable, there were
135 discrete changes in diameter. We were then able to generate three-dimensional
136 reconstructions at a near-atomic level of resolution for 12 discrete diameters, ranging from 270
137 to 335 Å (Figure 1B), each having a different helical symmetry, but each with seven strands
138 (Table 1; Figure S2A). The α -helices in the map had a clear hand and showed unambiguously
139 that the seven strands were left-handed, as had been determined at low-resolution by cryo-
140 electron tomography and sub-tomogram averaging. The resolution of the best volume (Table 1,
141 symmetry 7, ~ 3.5 Å, Figure 1C) would be more than adequate to build a full-atomic model *ab*
142 *initio* when given the protein sequence.

143 To determine the identity of the MCP, SMV1 proteins were separated by SDS-PAGE and
144 the major protein band of ~ 14 kDa was subjected to trypsin digestion followed by mass
145 spectrometry analysis (Figure S3). The dominant protein in the band, gp11 (W0UU99), was the
146 homolog of the MCPs previously identified in all other bicaudaviruses (Erdmann *et al.*, 2014;
147 Hochstein *et al.*, 2016; Prangishvili *et al.*, 2006; Xiang *et al.*, 2005). However, it was impossible
148 to thread the sequence of this putative MCP (gp11) through the density map, trying both
149 possible N-to-C orientations. The absence of density in the map for a bulky side chain might be
150 explained by disorder of that residue, but no threading could account for the pattern of bulky
151 side chains seen in the map. Further, while the atomic structures of the ATV (PDB ID: 3FAJ) and
152 ATSV (PDB ID: 5EQW) putative MCPs each displayed a compact four-helix bundle with a short
153 fifth helix projecting out, the SMV1 capsid protein was clearly composed of two long helices
154 connected by a short turn. We therefore searched for other possible candidates. A C_α trace of

155 the density map suggested that the MCP is ~90 amino acids long. Assuming that there might be
156 disordered residues at either the N- or C-terminus that were not present in the density map, we
157 focused on SMV1 proteins containing between 90 and 150 residues. There were 17 such
158 candidates (out of 96 proteins), but none could be successfully threaded through the map.

159 We next performed the N-terminal sequencing of the proteins in the major gel band.
160 Two different five-residue N-terminal peptides were identified: one from a minor species,
161 VEDYF, corresponding to the N-terminus of gp11 (the putative MCP that did not fit the map),
162 and one from the major species, VTFGT, that did not correspond to the N-terminus from any of
163 the 96 ORFs in the SMV1 genome. A search of the ORFs, found this pentapeptide uniquely in
164 SMV1 gp03 (W0UUV5), a hypothetical 157-residue sequence annotated as a membrane
165 protein. Removing 59 N-terminal residues would place the pentapeptide at the N-terminus of a
166 processed protein, and the resulting 98-residue protein fit the density map perfectly (Figure
167 1D). The 98-residue protein contains no useful tryptic digestion sites (one at the C-terminus
168 would create a three-residue peptide), explaining why peptides from this protein were never
169 detected after tryptic digestion and mass spectrometry. The processing is likely performed by a
170 cellular protease, because SMV1 and other bicaudaviruses do not encode identifiable
171 proteases. Notably, however, gp03 does not contain a canonical signal sequence or
172 recognizable signal peptidase cleavage site (Figure S4A). The fit revealed that additional density
173 facing the outside of the capsid, near the turn between the two helices, was most likely due to
174 heavy N-linked glycosylation of Asn53. Since the predicted MW for the 98-residue MCP is
175 10.1kDa, we suggest that extensive glycosylation of this single residue can explain why the
176 protein runs in a gel with an apparent MW of ~14kDa. The glycosylation on the outside of the

177 virion may play a role in protecting the capsid against very aggressive conditions, as was shown
178 for extensively glycosylated pili of hyperthermophilic and acidophilic archaea (Wang et al.,
179 2020; Wang et al., 2019). To further validate this, we used trifluoromethanesulfonic acid
180 (TFMS), which removes both N- and O-linked glycans from proteins (Sojar and Bahl, 1987).
181 Conditions were found under which a significant shift to lower mass could be obtained for a
182 silver-stained band that ran at ~13 kDa before acid treatment and at ~11 kDa after acid
183 treatment. These two bands were excised from the gel and analysed by mass spectrometry,
184 which found that the gp03 can only be detected in the lower band. Interestingly, the protein
185 gp11 was detected in both bands (Figure S3B-D). Thus, we designate SMV1 gp03 as the actual
186 MCP. Given the abundance of gp11 in the virions of all known bicaudaviruses and lack of any
187 discernable contribution to capsid formation, we hypothesize that this protein functions as a
188 nucleocapsid protein which compacts and, potentially, protects the viral DNA. Notably, in ATV,
189 gp42, one of the two ATV-encoded homologs of SMV1 gp11, has been shown to act as a strong
190 inhibitor of the host RNA polymerase (Sheppard et al., 2016), consistent with the possibility
191 that this soluble protein is injected into the cell upon viral genome delivery.

192

193 *Kinship among all spindle-shaped viruses*

194 We have imaged ATV by cryo-EM (Figure 2A) and carried out a similar analysis. As with
195 SMV1, ATV is formed from 7-start helical strands, and the subunit contains two α -helices linked
196 by a short turn. The repeat of the 7-start helices in ATV is 32 Å compared with the 29 Å in
197 SMV1. Although the resolution we have obtained for ATV (~6 Å) is much worse than for SMV1,
198 there is no ambiguity in building a homology model for the ATV gp59 (YP_319890) based upon

199 the SMV1 results, and this model fits well into the reconstructed density (Figure 2C). The crystal
200 structure for the putative MCP in ATV, gp62 (PDB id: 3FAJ)(Goulet *et al.*, 2010), cannot be fitted
201 into the density, and clearly is the structure of a different protein. Thus, we designate gp59 as
202 the actual MCP of ATV.

203 Homologs of SMV1 gp03 and ATV gp59 are conserved in all bicaudaviruses, including
204 STSV1 and STSV2 (Figure S4B). Based upon the results for SMV1 and ATV, everything suggests
205 that the actual MCP of ATSV is ATSV_F163, rather than ATSV_D135 for which an x-ray crystal
206 structure was determined (Hochstein *et al.*, 2018). The N-glycosylation sequon (NxS/T, where x
207 is any amino acid) located at the beginning of the second α -helix is conserved in all homologs
208 (Figure S4B), suggesting that glycosylation is important for virion assembly and/or stability.
209 Features of the bicaudavirus MCPs are remarkably similar to those of the MCPs of group I
210 spindle-shaped viruses, such as fusellovirus SSV1, halspivirus His1 and thaspivirus NSV1 (Figure
211 S4C). Notably, the MCP of SSV1 is encoded as a pre-protein which is proteolytically processed at
212 the N-terminus (Iverson *et al.*, 2017; Quemain *et al.*, 2015), as in the case of SMV1. Furthermore,
213 MCPs of fuselloviruses also contain the conserved N-glycosylation sites at the position
214 equivalent to that in bicaudaviruses and SSV1 MCP has been indeed shown to be glycosylated
215 (Quemin *et al.*, 2015). Finally, fuselloviruses, halspiviruses and thaspiviruses were all shown to
216 exist in tubular form, either upon virion egress or following genome ejection (Hong *et al.*, 2015;
217 Kim *et al.*, 2019; Quemain *et al.*, 2016). In light of this evidence, combined with the results of
218 bipartite network analysis showing that bicaudaviruses and group I spindle-shaped viruses
219 share several signature genes (Iranzo *et al.*, 2016), we suggest that all archaeal spindle-shaped
220 viruses have evolved from a common ancestor and use similar principles of virion organization

221 (Figure 3C). However, details in both the protein subunits and the virion morphology differ, as
222 would be expected from evolutionary divergence. For instance, viruses STSV1 and STSV2 have
223 only one, relatively short tail and do not undergo morphological changes outside of the cell
224 (Erdmann *et al.*, 2014; Xiang *et al.*, 2005); ATSV has a very long and narrow tail of uniform
225 length (Hochstein *et al.*, 2018); whereas ATV and SMV1 analyzed in the present paper can
226 “grow” tails from both pointed ends of the spindle (Haring *et al.*, 2005; Uldahl *et al.*, 2016).
227 Strikingly, this emerging virus lineage might extend beyond spindle-shaped viruses to also
228 include rod-shaped viruses of the family *Clavaviridae* (Mochizuki *et al.*, 2010). In particular, a
229 high resolution experimentally-determined structure exists for a rod-like clavavirus APBV1
230 virion (Ptchelkine *et al.*, 2017), which is constructed from highly hydrophobic subunits with a
231 helical-hairpin structure and an N-glycosylation site at the same position as in SMV1 and other
232 bicaudaviruses (Figure 3B). Although MCPs of APBV1 and SMV1 do not display significant
233 sequence similarity, which is not unexpected due to high sequence divergence and sparse
234 sampling of archaeal viruses, the interactions between the MCP subunits in the two groups of
235 viruses are very similar, full of hydrophobic interactions (Figure 3B).

236

237 *How the virions expand*

238 The subunit model from “symmetry 7” of SMV1 (Figure 1D) fits the other 11 volumes
239 with no modifications or perturbations. In addition, changes in the local packing of the subunit
240 were almost infinitesimally small in the diameter range from 270 to 335 Å (Figure 4,
241 Supplemental Movies 1 and 2), showing how the diameter could change significantly while at
242 the same time preserving a highly conserved local environment for each subunit. The RMSD

243 between two adjacent subunits in the same strand comparing the smallest and the largest
244 diameter was only 0.3 Å (Figure 4A), while the RMSD between subunits in adjacent strands was
245 only 0.4 Å (Figure 4B) when comparing the smallest and largest diameter.

246 Helical symmetry can be defined as a rise per subunit, a rotation per subunit, and a
247 possible C_n rotational symmetry. Two of the 12 structures had a C_7 rotational symmetry (Table
248 1, symmetries 1 and 8), so the rise per subunit in each strand is the same as the helical rise. For
249 the other symmetries in Figure 4C we have multiplied the helical rise by seven to yield the rise
250 per subunit within a strand. The determined diameters for these 12 points fall into a small
251 range, considering that the spindle bodies can have a maximum diameter of ~1,200 Å. Because
252 the 29 Å periodicity remains constant in both spindle body and tail, and this comes from the
253 repeat of the 7-start helices which have a fixed 203 Å pitch (7×29 Å) in both the tails and the
254 spindle body, the number of subunits per strand in one turn will be u , which is 203 Å divided by
255 the rise per subunit along a strand. We have found for the 12 symmetries reconstructed that u
256 is incremented by seven for each larger diameter (which is an increment of one per 29 Å),
257 explaining why the changes in diameter, D , are discrete, and not continuous. Let p be the path
258 length along a 7-start strand per 203 Å turn, then:

$$259 \quad p^2 = (\pi D)^2 + 203^2$$

260 We have shown that the local packing does not change as the diameter changes (Figure
261 4, Supplemental Movies 1,2), so the spacing between subunits along a strand, q , can be taken
262 as a constant as the diameter changes, and $p = uq$. From the observed tubes, $q=18.8$ Å. Given
263 this, the rise per subunit and the diameter can be easily calculated for each successive value of

264 *u.* Therefore, we can extrapolate the results from the tubes to the larger diameters that would
265 be found in the spindles (Figure 4C, Supplemental Movie 3).

266 How can the subunit structure explain the remarkable ability of these strands to slide
267 past each other in discrete steps? An analysis of the protein sequence (Figure 5A) shows that
268 most of the residues are quite hydrophobic. A plot of transmembrane probabilities for the
269 sequence indicates a strong prediction for two transmembrane helices (Figure 5B). This can be
270 seen in terms of the actual structure, where there is a hydrophobic region in the center of the
271 subunit that is ~ 35 Å thick, with polar regions below and above it (Figure 5C). Therefore, these
272 capsid proteins would not be soluble as monomers and must be integral to the host membrane
273 before the virions are formed. When packed into virions (Figure 5D), the hydrophobic regions
274 are completely buried and the polar regions are facing either the outside or the lumen. Notably,
275 the interface between neighboring 7-start strands is almost completely hydrophobic, with
276 minimal interactions between the hydrophilic regions. This will allow these strands to slide by
277 each other while still excluding solvent (Supplemental Movies 4,5). Consistent with this,
278 treatment of the virions with detergent, such as N-lauryl sarcosine, leads to rupture of the
279 virions (Figure S5A), much as such treatment would destabilize a membrane.

280 The extensive hydrophobicity of the capsid and likely insertion of the protein subunits
281 into the membrane prior to virion assembly motivated us to look for possible lipids associated
282 with the virions (Figure S5B). The distribution of lipids in the SMV1 virions was not the same as
283 that found in the host, suggesting that these lipids were selectively incorporated rather than
284 being contaminants from cellular debris. In bacterial mating pili, the subunits also have a
285 helical-hairpin structure (Costa et al., 2016; Zheng et al., 2020) and these subunits exist as

286 integral membrane proteins before polymerization and after depolymerization. A single lipid
287 molecule has been observed tightly bound to each subunit in the three published structures.
288 However, we found no ordered lipid molecules in the SMV1 tube reconstructions. We can
289 exclude the possibility that disordered lipid molecules (that we do not visualize) are present
290 between the hydrophobic helices as the space between these helices is too narrow to
291 accommodate a lipid. Since the tube segments used for three-dimensional reconstruction were
292 carefully screened to only include the most ordered and regular ones, we are thus left with the
293 possibility that the lipids may be found in the many dislocations or irregular regions in the
294 virions, in particular, where additional subunits are added to helical turns as the diameter
295 becomes larger. Alternatively, lipids could be associated with the specialized structures that are
296 present at the termini of SMV1 virions.

297

298

299 **Discussion**

300 We have shown how a highly hydrophobic subunit can assemble into tubes with a
301 variable diameter, and how these tubes can seamlessly expand and contract to form a spindle-
302 shaped body for a virion that maintains the same quasi-equivalent contacts between subunits
303 that are present in the tubes. The hydrophobic effect is, by definition, non-specific (Hillyer and
304 Gibb, 2016; Newberry and Raines, 2019), and the hydrophobic surfaces of the helices (Figure 5)
305 allow them to slide past each other (Supplemental Movies 4,5) while still maintaining the
306 integrity of the capsid and excluding very acidic solvent in the environment from the interior of
307 the virion. Our expectation is that the interior of the virions is close to the same near-neutral
308 pH as the cytoplasm of the host (Baker-Austin and Dopson, 2007) and that is why the capsid
309 must be impermeable to acidic solvent. While the hydrophobic effect alone is non-specific,
310 specificity in the intersubunit contacts is provided by the helix-helix contacts that exist between
311 neighboring subunits which dictate that discrete changes in diameter are required to maintain
312 these quasi-equivalent contacts (Supplemental Movies 4,5). The notion of quasi-equivalence
313 was first introduced by Caspar and Klug to explain how icosahedral viruses could be assembled
314 from many copies of a single protein (Caspar and Klug, 1962). If one only had 60 copies of such
315 a capsid protein a perfect icosahedron could be assembled so that every subunit was in an
316 equivalent environment. It was also known that it was impossible to place more than 60 copies
317 of a protein on the surface of a sphere such that all would be in strictly equivalent
318 environments. But a capsid built from only 60 protein subunits would be quite small, and it had
319 been observed that larger icosahedral viruses contained many more than 60 copies of the
320 capsid protein. The solution was found in the possibility of quasi-equivalence (Caspar and Klug,

321 1962), wherein multiples of 60 protein subunits could be assembled into a spherical shell
322 maintaining not identical, but quasi-equivalent contacts. This has now been amply confirmed
323 with high resolution structures for many icosahedral viruses (Damodaran et al., 2002; Johnson
324 and Olson, 2021; Johnson and Speir, 1997), although it has also been shown that certain
325 icosahedral viruses can violate quasi-equivalence (Rayment et al., 1982; Stehle et al., 1994).
326 However, the extension of quasi-equivalence theory to the radial expansion of helical viruses
327 has not been envisioned.

328 Here we show that the quasi-equivalence principle explaining the ability of icosahedral
329 viruses to expand the number of subunits needed to encapsulate larger genomes has a striking
330 parallel in helical viruses. In APBV1 or other rod-like viruses, there will be a stoichiometric ratio
331 between the number of capsid proteins and the number of bases or base pairs encapsidated. As
332 pointed out by Caspar and Klug (Caspar and Klug, 1962), the length of tobacco mosaic virus
333 virions is simply determined by the length of the single-stranded RNA genome given the
334 stoichiometric ratio of three bases per protein subunit. In filamentous bacteriophage, the
335 packaging of DNA involves charged residues in the lumen of the phage, and thus modifications
336 of these residues can change the length of the virions while the length of the genome is
337 unaltered (Hay and Lithgow, 2019). Thus, there is a linear relation between the length of the
338 virion and the size of the genome, even though the actual stoichiometry can be modified. In
339 contrast, for spherical viruses the number of capsid subunits will be proportional to the surface
340 area, which is the radius squared, while the size of the genome encapsidated will scale as the
341 radius cubed. Thus, the ratio of base pairs per capsid subunit will scale as the radius, becoming
342 more efficient in genome packaging the larger the capsid.

343 With hydrophobic subunits having the ability to slide past each other, it is easy to
344 imagine how a rod-like virus such as APBV1 could evolve into a spindle-shaped virus (Figure 3C)
345 while still maintaining all of the local interactions that hold the capsid together. We suggest
346 that the minimum energy conformation of the spindle-shaped capsids is actually a tube, as
347 evidenced by the fact that when the genome is released, the SMV1 spindles relax into tubes
348 (Figure S1). The same spindle-to-tube transformation has been observed upon genome ejection
349 for a group I spindle-shaped virus His1 (Hong *et al.*, 2015). It is therefore the pressure of the
350 genome that expands the capsid that would normally be rod-like into a spindle shape. Thus, the
351 unusual spindle morphology results from the radial expansion of a rod.

352 Archaeal and bacterial gas vesicles (Li and Cannon, 1998; Pfeifer *et al.*, 2002; Strunk *et*
353 *al.*, 2011) have some remarkable similarity in morphology to spindle-shaped viruses. The gas
354 vesicles are made of a single major structural protein of ~8 kDa and have a hydrophobic interior
355 surface. However, whereas the capsid proteins contain two hydrophobic α -helices, those of gas
356 vesicles from both bacteria and archaea have a α - β - β - α structure (Knitsch *et al.*, 2017; Strunk *et*
357 *al.*, 2011), and the β -strands are predicted to mediate subunit-subunit interactions. Finally,
358 unlike the capsid proteins which are predicted to contain two transmembrane domains, the
359 proteins of gas vesicles are predicted to have none. Thus, we conclude that the superficial
360 similarity between spindle-shaped viruses and gas vesicles is a result of convergent evolution.
361 However, given the hydrophobic nature of both structures, we predict that changes in diameter
362 described for both are mediated by these hydrophobic interactions.

363 There has been great interest in understanding the role of BAR domain proteins (Frost
364 *et al.*, 2008), ESCRT-III (Endosomal Sorting Complexes Required for Transport) (Pfitzner *et al.*,

365 2021) and dynamin family proteins (Antonny et al., 2016) in membrane remodeling. All of these
366 proteins form helical tubes of variable diameters that have been shown to constrict as part of
367 their role in processes such as membrane fission and vesiculation. Structural studies of these
368 assemblies have been extremely difficult due to the variability of tube diameters, and we still
369 do not have any models at an atomic level of detail for the elementary steps in constriction.
370 Similarly, keratin filaments, assembled from α -helical coiled-coils, have also been revealed to
371 form variable diameter tubes *in situ* (Weber et al., 2021). While we do not suggest any
372 homology between these proteins and the SMV1/ATV capsid proteins, we suggest that the
373 general principle underlying the sliding of the strands of SMV1 proteins past each other while
374 maintaining the structural integrity of the capsid might extend to other filamentous assemblies
375 with variable diameters. The great advances in cryo-EM that have allowed us to determine in
376 atomic detail the structure of SMV1 capsid will likely allow a similar level of understanding for
377 other assemblies in the near future.

378

379 *Limitations of the Study*

380 We have determined a high resolution structure for only a single spindle-shaped virus, SMV1,
381 and determined a lower resolution structure for a second spindle-shaped virus, ATV. These
382 results have been combined with an existing high resolution structure for the clavavirus APBV1
383 (Ptchelkine *et al.*, 2017) and homology models of the major capsid proteins from other spindle-
384 shaped viruses to show the likely common ancestry and suggest how the spindle-shaped viruses
385 have evolved from rod-like viruses such as APBV1. Future high resolution studies of other rod-
386 like and spindle-shaped viruses will greatly extend our understanding of this evolution. Just as

387 knowing the morphology and genetics of apes and humans suggests common ancestry,
388 determining the intermediates that existed in this evolutionary process gives one a much more
389 profound understanding of biological history.

390
391 **Acknowledgments**

392 We are grateful to Dr. Xu Peng (University of Copenhagen) for the kind gift of SMV1 and its
393 host, *Saccharolobus islandicus* Δ C1C2. Part of the cryo-EM imaging of SMV1 was done at the
394 Molecular Electron Microscopy Core Facility at the University of Virginia, which is supported by
395 the School of Medicine. Part of the Cryo-EM and the Cryo-ET imaging were done at the Nano-
396 Imaging Core Facility at Institut Pasteur, created with the help of a grant from the French
397 Government's Investissements d'Avenir program (EQUIPEX CACSICE - Centre d'analyse de
398 systèmes complexes dans les environnements complexes, ANR-11-EQPX-0008). This work was
399 supported by NIH Grant GM122510 (E.H.E.) and K99GM138756 (F.W.). The work in the M.K.
400 laboratory was supported by grants from l'Agence Nationale de la Recherche (ANR-17-CE15-
401 0005-01, ANR-20-CE20-009-02 and ANR-21-CE11-0001-01) and Ville de Paris (Emergence(s)
402 project MEMREMA). We are grateful to Youssef Ghorbal for the help with data transfer, to
403 Caglar Yildiz for support in the lipid analysis and to the Ultrastructural Biolmaging unit of
404 Institut Pasteur for access to electron microscopes.

405

406

407

408 **Author Contributions**

409 M.K. initiated and developed the project. F.W. performed microscopy, image analysis,
410 reconstructed the SMV1 and ATV tubes, and generated and refined the atomic models. V.C-K.
411 cultured the viruses, developed the conditions for tube formation, and performed sample
412 preparation and fractionation analyses. M.V. and J-M.W. performed microscopy. Z.S. analyzed
413 data and developed graphical representations. L.C.B. did the deglycosylation of the MCP.
414 M.A.B.K., M.V. and J-M.W. collected the tomographic data. J.L. analyzed the tomographic data
415 and generated the three-dimensional reconstruction. S.S. analyzed the lipids. E.H.E. obtained
416 funding, analyzed data and supervised the research. F.W., M.K. and E.H.E. wrote the manuscript
417 with input from all authors.

418
419 **Declaration of Interests**

420
421 The authors declare no competing interests.
422
423

424 **Figure Legends**

425 **Figure 1. Cryo-EM of bicaudavirus SMV1**

426 **A**, Representative cryo-EM of the *Sulfolobus monocaudavirus 1* (SMV1). A 29 Å periodicity can
427 be seen in both spindle-shaped bodies and tails. Scale bar, 50 nm.

428 **B**, Cryo-EM reconstructions of the SMV1 tails made from 12 different diameters. The top view
429 models of the largest (symmetry 12) and smallest (symmetry 1) diameters are shown.

430 **C**, A surface of the 3.5 Å resolution reconstruction of symmetry 7. The left-handed 7-start
431 protofilaments and 29 Å spacing between them are labeled.

432 **D**, Ribbon model of a single MCP protein (left). The map quality of the subunit is shown on the
433 right. The arrow points to the N-linked glycosylation associated with Asn53.

434

435 **Figure 2. Cryo-EM of the *Acidianus two-tailed virus* (ATV)**

436 **A**, Representative cryo-EM of ATV. A 32 Å periodicity can be seen in both spindle-shaped bodies
437 and tails, indicated by black arrows. Scale bar, 50 nm.

438 **B**, Surface of the cryo-EM reconstruction of the ATV tails for one diameter, with the 7-start
439 protofilaments labeled.

440 **C**, The homology model of ATV fit into the cryo-EM map in side and top views.

441

442 **Figure 3. Spindle-shaped viruses evolved to carry larger genomes**

443 **A**, Single subunit structures of seven MCPs. The structures of the APBV1 and SMV1 MCPs have
444 been determined by high-resolution cryo-EM (PDB 5OXE and 7ROC). The other five MCP
445 structures are homology models based on the SMV1 structure (see Methods).

446 **B**, Comparison of hydrophobic interactions of APBV1 and SMV1.

447 **C**, The estimated internal volumes of seven viruses plotted against genome size.

448

449 **Figure 4. The local contacts between capsid subunits are conserved as the diameter changes**

450 **A**, Interactions along the 7-start protofilament. One protofilament from the tube is highlighted
451 on the left. Two subunits in one protofilament from symmetry 1 and 12 are aligned (center) and
452 compared (right).

453 **B**, Interactions between adjacent 7-start protofilaments. Two protofilaments are highlighted on
454 the left. The contacts between three subunits, two on the bottom strand (S_a and S_b) and one on
455 the top (S'_a) are compared in two views (center and right).

456 **C**, The rise per subunit along a strand for 12 different symmetries is plotted versus the diameter
457 of the tubes (orange spheres), showing a nearly linear relationship. The blue dashed line shows
458 the extrapolation to both smaller and larger diameters.

459

460 **Figure 5. The SMV1 MCP has clustered lipophilicity**

461 **A**, Amino acid composition of the SMV1 MCP.

462 **B**, TMHMM prediction of transmembrane probabilities for the SMV1 MCP sequence.

463 **C**, The spatial distribution of hydrophobic and hydrophilic residues in a single MCP.

464 **D**, Different views of the SMV1 tubes colored by lipophilicity.

465

466
467

Table 1. Helical symmetry and model statistics for SMV₁ and ATV

Symmetry ID	SMV ₁												ATV
	1	2	3	4	5	6	7	8	9	10	11	12	---
Helical symmetry													
Point group	C7	C1	C1	C1	C1	C1	C1	C7	C1	C1	C1	C1	C1
Helical rise (Å)	5.15	0.72	0.70	0.68	0.67	0.66	0.64	4.38	0.61	0.60	0.59	0.58	1.08
Helical twist (°)	-9.0	50.2	-155.5	-104.1	101.7	153.1	-52.6	-7.7	50.4	-155.3	-103.9	101.8	-156.0
Map resolution (Å)													
Model:map FSC (Å, 0.5)	5.1	4.8	4.3	4.1	4.1	3.9	3.7	3.8	3.7	3.8	4.0	4.3	---
Map:map FSC (Å, 0.143)	4.4	4.3	3.8	3.8	3.8	3.8	3.7	3.7	3.4	3.5	3.6	4.0	6.3
Refinement and Model validation													
Bond lengths RMSD (Å)	0.004	0.004	0.006	0.005	0.006	0.004	0.003	0.004	0.006	0.007	0.005	0.004	---
Bond angles RMSD (°)	0.658	0.640	0.770	0.841	0.671	0.699	0.635	0.703	0.681	0.735	0.651	0.658	---
Clashscore	12.4	9.9	12.0	9.3	10.0	8.4	9.0	10.9	9.2	11.0	8.7	7.4	---
Ramachandran Favored (%)	96.7	95.7	94.6	95.7	94.6	95.7	93.4	95.0	93.4	94.6	94.6	95.7	---
Ramachandran Outlier (%)	0	0	0	0	0	0	0	0.2	0	0	0	0	---
RSCC	0.86	0.87	0.87	0.87	0.88	0.87	0.87	0.88	0.87	0.87	0.86	0.87	---
Deposition ID													
PDB (model)	7RO2	7RO3	7RO4	7RO5	7RO6	7ROB	7ROC	7ROD	7ROE	7ROG	7ROH	7ROI	---
EMDB (map)	24585	24586	24587	24588	24589	24590	24591	24592	24593	24594	24595	24597	24596

468
469
470

471
472

473

474 **STAR Methods**

475

476 **RESOURCE AVAILABILITY**

477

478 **Lead Contact**

479 Further information and requests for resources and reagents should be directed to and will be

480 fulfilled by the lead contact, Edward H. Egelman, egelman@virginia.edu

481 **Materials Availability**

482

- 483 • This study did not generate any new unique reagents.

484

485 **Data and Code Availability**

- 486 • The SMV1 maps were deposited in the Electron Microscopy Data Bank (EMDB) with
487 entry codes EMD-24585, EMD-24586, EMD-24587, EMD-24588, EMD-24589, EMD-
488 24590, EMD-24591, EMD-24592, EMD-24593, EMD-24594, EMD-24595, and EMD-24597
489 and the respective atomic models were deposited in the Protein Data Bank (PDB) with
490 entry codes 7RO2, 7RO3, 7RO4, 7RO5, 7RO6, 7ROB, 7ROC, 7ROD, 7ROE, 7ROG, 7ROH
491 and 7ROI. The ATV map was deposited in the EMDB with entry code EMD-24596.

492

- 493 • This paper does not report original code.

494

- 495 • Any additional information required to reanalyze the data reported in this paper is
496 available from the lead contact upon request

497

498

499

500 **EXPERIMENTAL MODEL AND SUBJECT DETAILS**

501

502 **Archaeal strains and growth conditions**

503 *Sulfolobus monocaudavirus 1*, SMV1, was propagated in *Sulfolobus islandicus* CRISPR deletion
504 mutant delta C1C2 (Gudbergdottir et al., 2011). The host cell culture was grown in *Sulfolobus*
505 medium supplemented with 0.2% tryptone, 0.1% yeast extract, 0.2% sucrose and 0.002% uracil
506 (Zillig et al., 1993). The culture was started from -80°C stock. The cells were grown in 50 ml of
507 medium at 76°C with shaking. After 24 h of incubation, the cell culture was diluted 20 times in
508 pre-warmed (76°C) medium and the growth was continued until it reached an OD_{600} of ~ 0.2 .
509 Then the cells were infected with SMV1 stock and incubation continued at 76°C with agitation
510 for 24h. 24 h post infection the cells were removed by centrifugation in Sorval 3000 rotor, 7000
511 rpm, 20 min, 15°C , virus-containing supernatant was collected and virions precipitated with
512 PEG600 (10.5% w/v) and NaCl (5.8% (w/v) for 2h at room temperature. The PEG-precipitate was
513 then pelleted in a Sorval 3000 rotor, 9000 rpm, 30 min, 15°C and the resulting pellet was re-
514 suspended in one tenth of the original volume in 20 mM Tris-acetate (pH6) buffer.

515

516 **METHODS DETAILS**

517 **Virus production and treatments**

518 To purify the virus, solid CsCl was added to the virus suspension to a final concentration of 45%
519 (w/v). The samples were run in a Beckman SW41 rotor, 38,000 rpm, at least 16 h, 15°C . After the
520 run, the opalescent band corresponding to SMV1 virions was collected. The purified SMV1 was
521 stored in CsCl at 4°C until used.

522 For SMV1 virion dissociation, CsCl purified virions were diluted 50 times in 20 mM Tris-acetate
523 (pH6) buffer and pelleted down by centrifugation in a Beckman 45Ti rotor, 35,000 rpm, 2 h, 15°C.
524 The resultant pellet was re-suspended in 20mM Tris-acetate (pH6) buffer and used for
525 dissociation. The SMV1 virions were dissociated by addition of 0.1% (v/v) NaOH and incubation
526 at 37°C for 1 h. Then non-treated (control) and NaOH-treated virions were loaded on top of the
527 linear 5-20 % sucrose gradients and run in a Beckman SW41 rotor, 24,000 rpm, 20 min, 15°C.
528 After the run, 12 fractions from the top of each tube were collected and analyzed in 4-12% Bis-
529 Tris acrylamide gels. After the run the gels were stained for DNA with ethidium bromide and for
530 proteins with InstantBlue. The contents of the collected fractions were also visualized by negative
531 staining TEM. For that, 10 µl of the fraction of interest was adsorbed onto copper grids with
532 carbon-coated Formvar films and negatively stained with 2.0% (w/v) uranyl acetate for 20 s. The
533 samples were observed under a FEI Tecnai BioTwin 120 microscope operated at 120 kV.

534

535 **Cryo-electron microscopy and image processing**

536 The SMV1 sample was applied to glow-discharged lacey carbon grids and vitrified using a
537 Vitrobot Mark IV (Thermo Fisher). Grids were imaged at on a Titan Krios (300 keV, Thermo
538 Fisher) with a K3 camera (Gatan). 16,401 micrographs were collected under electron counting
539 mode at 1.1 Å per pixel, using a defocus range of 1–2 µm with ~50 electrons/Å² distributed into
540 40 fractions. Motion correction and CTF estimation were done in cryoSPARC (Punjani et al.,
541 2017; Rohou and Grigorieff, 2015; Zheng et al., 2017). A total of seven million particles were
542 auto-picked by “Filament Tracer” with a shift of 13 pixel, and non-virion bad particles were
543 removed by 2D classification. About six million SMV1 particles remained having a range of

544 diameters. From this six million particle dataset, 12 relatively homogeneous subsets were
545 sorted out by iterative 2D classifications. The possible helical symmetries were calculated from
546 an averaged power spectrum for each subset, generated from the raw particles. For each
547 subset, the actual helical symmetry was determined in cryoSPARC by trial and error, until the
548 hand of α -helices and amino acid side chains were seen (Egelman, 2000; Punjani et al., 2020).
549 The resolution of each reconstruction was estimated by both Map:Map FSC and Model:Map
550 FSC. The final volumes were then sharpened with a negative B-factor automatically estimated
551 in cryoSPARC, and the statistics are listed in Table 1.
552 For ATV the same approach was used, with 2.8M particle images collected using a shift of 9
553 pixels between adjacent boxes.

554

555 **Model building of SMV1 MCP**

556 The density corresponding to a single SMV1 MCP was segmented from the experimental cryo-
557 EM density using Chimera (Pettersen et al., 2004). Possible MCP sequences were threaded
558 through the density using DeepTracer (Pfab et al., 2021). Only one sequence, WOUUV5, also
559 detected by N-terminal sequencing, could be threaded through the map, matching the bulky
560 side chain densities and the glycosylation site. This model was adjusted manually in Coot
561 (Emsley and Cowtan, 2004) and real-space refined in PHENIX (Afonine et al., 2018). Using the
562 determined helical symmetry, a filament model was generated in Chimera and refined against
563 the full cryo-EM map using PHENIX real-space refinement. MolProbity was used to evaluate the
564 quality of the filament model (Williams et al., 2018). The refinement statistics are shown in
565 Table 1.

566

567 **Cryo-electron tomography**

568 For cryo-electron tomography, CsCl-purified virions were pelleted by ultracentrifugation
569 (Beckman Type 50.2 Ti rotor, 32000 rpm, 2h, 15 °C), resuspended in water and incubated at 90
570 °C for one week. Heat-treated SMV1 particles were mixed with CsCl-purified rudivirus SIRV2 as
571 a control, since the helical hand had been determined previously (DiMaio et al., 2015). Data
572 were collected on a Titan Krios microscope (TFS) operating at 300kV in EFTEM mode using a
573 bioquatum/K3 (Gatan) energy filter and camera with a 20 eV slit. Tilt series were recorded at a
574 magnification of 26kx corresponding to 3.4 Å/pixel. A tilt range of +/-60° was set with a 3° tilt
575 increment and a total dose of 120 e⁻/Å² with a dose symmetric tilting scheme using the TFS
576 tomography software with defocus values between -4 and -6 micron defocus.

577 A total of 935 segments were manually picked along the long tails of SMV1 from 8×
578 binned tomograms by “tomopick” in the tomography package i3 (Winkler, 2007). Sub-
579 tomograms of SMV1 were first extracted from the 8× binned tomograms, and then aligned in
580 the i3 software package. After initial alignment, sub-tomograms were extracted from unbinned
581 tomograms for further refinement. Multivariate statistical analysis (MSA) implemented in the I3
582 package was then used for 3D classification and sub-tomogram averaging. As a control, a total
583 of 200 segments were manually picked from the tubes of SMV2 from the same 8× binned
584 tomograms. The similar procedure was used to generate a 3D averaged structure of the SIRV2
585 tube without applying any helical symmetry. The averaged structure of SMV1 is a left-handed
586 helix with ~29 Å periodicity, while the averaged structure of SIRV2 is a right-handed helix with
587 ~43 Å periodicity.

588

589 **Homology modeling of ATV, SSV1, His1 and ATSV MCP**

590 The MCP homologs in bicaudaviruses were identified by PSI-BLAST (Altschul et al., 1997)
591 searches (E-value cutoff of 0.05) queried with the sequence of SMV1 MCP (YP_009008070)
592 against the NCBI protein database restricted to members of the family *Bicaudaviridae*
593 (taxid:423358). The sequences were aligned using PROMALS3D (Pei and Grishin, 2014). MCPs of
594 SSV1 and His1 were determined experimentally (Pietila et al., 2013; Queminn *et al.*, 2015). The
595 MCP structures of ATV, SSV1, His1 and ATSV were first predicted by AlphaFold2 (Jumper et al.,
596 2021). Then the predicted structures were flexibly aligned to SMV1 by FATCAT (Li et al., 2020).
597 For ATV, the predicted MCP structure was then docked into the 6 Å resolution cryo-EM map
598 and real-space refined in PHENIX (Afonine *et al.*, 2018) with helical symmetry imposed.

599

600 **N-terminal sequencing of SMV1 by Edman Degradation**

601 The analysis was performed on an ABI Procise 494 sequencer. The sample was placed on a
602 PVDF membrane for the Edman degradation with a cyclic procedure where residues were
603 cleaved off one at a time and identified by chromatography. There were three steps in each
604 cycle. In step 1 the PITC reagent was coupled to the N-terminal amino group under alkaline
605 conditions. In step 2 the N-terminal residue was cleaved in acidic media. In step 3, the PITC-
606 coupled residue was transferred to a flask, converted to a PTH-residue and identified by HPLC
607 chromatography. The cycle was then started again for the identification of the next N-terminal
608 residue.

609

610 **SMV1 deglycosylation reaction and tandem mass spectrometry**

611 Briefly, 50 μ l of concentrated SMV1 were lyophilized. Then, 150 μ l of TFMS was added to the
612 tube and the mixture was incubated at 4 °C for 24 h. Following this, 150 μ l of 60% pyridine
613 solution, cooled to ~15 °C with a methanol dry ice bath, was added to the reaction tube,
614 neutralizing the TFMS acid. Using a Slide-A-Lyzer dialysis cassette with a 2,000 Da protein
615 molecular weight cut-off (Thermo Fisher Scientific), the reaction solutions were removed from
616 the sample with overnight dialysis at 4 °C into Tris/HCl (pH 8) buffer. After dialysis, small
617 amounts of aggregates—presumably deglycosylated SMV1—were observed in the cassette.
618 Centrifugation for 15 min at 4 °C and 20,000 $\times g$ was used to pellet the aggregates. The pellet
619 was resuspended with 50 μ l of Tris buffer (pH 8). SDS-PAGE was then performed at a constant
620 voltage of 120 V using 16.5% precast polyacrylamide Mini-PROTEAN Tris-Tricine precast gels
621 (Bio-Rad) and Tris/Tricine/SDS running buffer (Bio-Rad). For each sample, 15 μ l of boiled sample
622 in SDS was added to separate wells in the gel, and 5 μ l of Precision Plus Protein Dual Xtra
623 protein standard (Bio-Rad) was used as a marker. Silver staining was performed using a Pierce
624 Silver Stain for Mass Spectrometry kit (Thermo Fisher Scientific). The detected gel bands were
625 then excised, and mass spectrometry (Alphalyse) was performed to analyze the composition of
626 each band. The protein samples were reduced and alkylated with iodoacetamide, and
627 subsequently digested with chymotrypsin. The resulting peptides were concentrated by Speed
628 Vac lyophilization. The peptides were dissolved in 0.1% formic acid and injected on a Dionex
629 Ultimate 3000 nano-LC system (Thermo Scientific) coupled to a Bruker Maxis Impact QTOF mass
630 spectrometer for MS/MS analysis. Each sample underwent a 30-minute-long gradient run on
631 the instruments. As an external control of the instrument, tryptic BSA was analysed together

632 with the samples. The results were run against the Sulfolobus monocaudavirus SMV1 proteins
633 from the UniProt.

634

635 **Analysis of SMV1 and host cell lipids**

636 For lipid analyses, CsCl-purified virions were pelleted by ultracentrifugation (Beckman Type 50.2
637 Ti rotor, 35000 rpm, 2h, 15 °C), resuspended in 20 mM Tris-acetate (pH6) and run on 5-20 %
638 (wt/vol) sucrose gradient (Beckman SW32 Ti rotor, 24000 rpm, 30 min, 15 °C). The opalescent
639 virus-containing band was collected, pelleted as above and resuspended in 20 mM Tris-acetate
640 (pH6). The cellular and viral lipids were analyzed by UHPLC-MSD at NIOZ using an Agilent 1290
641 Infinity II ultra-high performance LC coupled to a 6230 Agilent MSD in selected ion mode as
642 described (Besseling et al., 2020).

643

644

645

646 **Supplemental Figures**

647

648 **Figure S1. Related to Figure 1. The pH shift to induce SMV1 to form long tails**

649 **A**, Cryo-EM of SMV1 before and after the NaOH (pH=10) treatment. Scale bars are 50 nm. The
650 arrow indicates a tube containing a viral genome, while all of the other tubes in this image are
651 empty.

652 **B**, SDS-PAGE and negative staining analysis of different fractions from the 5-20% sucrose
653 gradient which were loaded with non-treated virus (left) and NaOH-treated virus (right). The
654 non-treated virus migrated to the bottom of the gradient, where most of the protein and DNA
655 were found (fractions 8-12), whereas following NaOH treatment, the viral DNA was found in
656 fraction 1 in the form of amorphous aggregates, and the virion tubes were enriched in fraction
657 3.

658

659 **Figure S2. Related to Figure 1 and Table 1. Cryo-EM volumes and helical symmetries for all 12**
660 **reconstructions of SMV1**

661 **A**, The symmetry ID, helical rise, helical twist and point group symmetry are labeled.

662 **B**, The “gold-standard” half map FSCs were calculated in cryoSPARC using a 0.143 criterion.

663

664 **Figure S3. Related to Figure 1. TFMS deglycosylation analysis of SMV1**

665 **A**, SDS-PAGE with Coomassie blue staining of SMV1. The MW marker lane M is labeled on the
666 left. A single band at ~ 13-14 kDa is observed for SMV1.

667 **B**, SMV1 samples on silver-stained SDS-PAGE before and after deglycosylation using TFMS. M
668 indicates the molecular weight markers lane.

669 **C**, The band indicated by blue arrowhead in (B) was cut and used for tandem mass
670 spectrometry analysis. Protein W0UU99 was detected.

671 **D**, The band indicated by red arrowhead in (B) was cut and used for tandem mass spectrometry
672 analysis. Both W0UU99 and the correct capsid protein, W0UUv5, were detected.

673

674 **Figure S4. Related to Figure 3. Sequence analysis and alignments**

675 **A**, SignalP (Almagro Armenteros et al., 2019) analysis did not reveal any signal peptides or signal
676 peptidase cleavage sites in the cleaved N-terminus.

677 **B**, The N-terminal regions which are processed in the mature proteins are not shown. The start
678 site experimentally determined for SMV1 is indicated with an arrowhead. The secondary
679 structure determined for the SMV1 protein is shown above the alignment, with the red ribbon
680 representing α -helices. The N-glycosylation seqon is boxed, with the target Asn residue
681 indicated with an asterisk. Each sequence is identified with the corresponding GenBank
682 accession number followed by the virus name.

683 **C**, The predictions of transmembrane helices within the MCPs of seven viruses were made using
684 the TMHMM server v2.0.

685

686 **Figure S5. Related to Figure 5. Sensitivity to detergent and presence of lipids in SMV1 virions**

687 **A**, Transmission electron micrographs of negatively stained SMV1 virions treated with 0.1% N-
688 lauryl sarcosine. Bars, 200 nm. **B**, Distribution of the lipid species identified in SMV1 host,

689 *Saccharolobus islandicus*, and SMV1 virions. GDGT, glycerol dibiphytanyl glycerol tetraether
690 lipids; GTGT, glycerol tribiphytanyl glycerol tetraether lipids. Numbers following GDGT or GTGT
691 represent the number of cyclopentane rings present in the lipid species.
692
693

694 **Supplementary Movie 1. Related to Figure 4.**

695

696 An alignment of one subunit (yellow) between two adjacent diameters: symmetry 7 (cyan) and
697 symmetry 8 (magenta).

698

699 **Supplementary Movie 2. Related to Figure 4.**

700

701 An alignment of one subunit (yellow) between two very different diameters: symmetry 2
702 (orange) and symmetry 12 (green).

703

704 **Supplementary Movie 3. Related to Figure 4.**

705

706 The helical symmetry established for the tubes and extrapolated to the spindle-shaped bodies
707 (Fig. 4C) has been used to construct a model for the full capsid. The seven strands of subunits
708 are each shown in a different color. Although discrete subunits are not seen in this visualization,
709 98,000 copies of the capsid protein have been used to generate this model. The basis of the
710 model is that the local packing throughout the structure is fixed with a spacing of 2.7 Å
711 between subunits along a strand, and the helical pitch of each strand is also fixed at 203 Å. As
712 the diameter increases due to internal pressure from the packaged genome, the strands slide
713 past each other.

714

715 **Supplementary Movie 4. Related to Figure 5.**

716

717 The role of hydrophobic (lipophilic) surfaces in the sliding of SMV1 strands past each other. This
718 animation shows the continuous transformation of the same number of total subunits from the
719 narrowest diameter tube to the widest diameter tube of the 12 that we have reconstructed. As
720 the tube becomes wider, more subunits are added per turn to the left-handed 7-start helices.
721 The hydrophobic surfaces (gold) at the interfaces between the strands allow for this sliding to
722 take place, at the same time that solvent is excluded from passing through the capsid.

723

724 **Supplementary Movie 5. Related to Figure 5.**

725 The top view of Supplemental movie 4.

726

727 **References**

728 Afonine, P.V., Poon, B.K., Read, R.J., Sobolev, O.V., Terwilliger, T.C., Urzhumtsev, A., and Adams,
729 P.D. (2018). Real-space refinement in PHENIX for cryo-EM and crystallography. *Acta Crystallogr*
730 *D Struct Biol* 74, 531-544. 10.1107/S2059798318006551.

731 Almagro Armenteros, J.J., Tsirigos, K.D., Sonderby, C.K., Petersen, T.N., Winther, O., Brunak, S.,
732 von Heijne, G., and Nielsen, H. (2019). SignalP 5.0 improves signal peptide predictions using
733 deep neural networks. *Nat Biotechnol* 37, 420-423. 10.1038/s41587-019-0036-z.

734 Altschul, S.F., Madden, T.L., Schaffer, A.A., Zhang, J., Zhang, Z., Miller, W., and Lipman, D.J.
735 (1997). Gapped BLAST and PSI-BLAST: a new generation of protein database search programs.
736 *Nucleic Acids Res* 25, 3389-3402.

737 Antonny, B., Burd, C., De Camilli, P., Chen, E., Daumke, O., Faelber, K., Ford, M., Frolov, V.A.,
738 Frost, A., Hinshaw, J.E., et al. (2016). Membrane fission by dynamin: what we know and what
739 we need to know. *EMBO J* 35, 2270-2284. 10.15252/embj.201694613.

740 Baker-Austin, C., and Dopson, M. (2007). Life in acid: pH homeostasis in acidophiles. *Trends in*
741 *microbiology* 15, 165-171. 10.1016/j.tim.2007.02.005.

742 Baquero, D.P., Liu, Y., Wang, F., Egelman, E.H., Prangishvili, D., and Krupovic, M. (2020).
743 Structure and assembly of archaeal viruses. *Advances in virus research* 108, 127-164.
744 10.1016/bs.aivir.2020.09.004.

745 Besseling, M.A., Hopmans, E.C., Bale, N.J., Schouten, S., Damste, J.S.S., and Villanueva, L. (2020).
746 The absence of intact polar lipid-derived GDGTs in marine waters dominated by Marine Group
747 II: Implications for lipid biosynthesis in Archaea. *Sci Rep* 10, 294. 10.1038/s41598-019-57035-0.

748 Caspar, D.L., and Klug, A. (1962). Physical principles in the construction of regular viruses. *Cold*
749 *Spring Harb. Symp. Quant. Biol.* 27, 1-24.

750 Costa, T.R., Ilangovan, A., Ukleja, M., Redzej, A., Santini, J.M., Smith, T.K., Egelman, E.H., and
751 Waksman, G. (2016). Structure of the Bacterial Sex F Pilus Reveals an Assembly of a
752 Stoichiometric Protein-Phospholipid Complex. *Cell* 166, 1436-1444 e1410.
753 10.1016/j.cell.2016.08.025.

754 Crick, F.H., and Watson, J.D. (1956). Structure of small viruses. *Nature* 177, 473-475.
755 10.1038/177473a0.

756 Damodaran, K.V., Reddy, V.S., Johnson, J.E., and Brooks, C.L., 3rd (2002). A general method to
757 quantify quasi-equivalence in icosahedral viruses. *J. Mol. Biol.* 324, 723-737. 10.1016/s0022-
758 2836(02)01138-5.

759 Dellas, N., Snyder, J.C., Bolduc, B., and Young, M.J. (2014). Archaeal Viruses: Diversity,
760 Replication, and Structure. *Annu Rev Virol* 1, 399-426. 10.1146/annurev-virology-031413-
761 085357.

762 DiMaio, F., Yu, X., Rensen, E., Krupovic, M., Prangishvili, D., and Egelman, E.H. (2015). *Virology*.
763 A virus that infects a hyperthermophile encapsidates A-form DNA. *Science* 348, 914-917.
764 10.1126/science.aaa4181.

765 Egelman, E.H. (2000). A robust algorithm for the reconstruction of helical filaments using single-
766 particle methods. *Ultramicroscopy* 85, 225-234.

767 Emsley, P., and Cowtan, K. (2004). Coot: model-building tools for molecular graphics. *Acta*
768 *Crystallogr.D.Biol.Crystallogr.* 60, 2126-2132.

769 Erdmann, S., Chen, B., Huang, X., Deng, L., Liu, C., Shah, S.A., Le Moine Bauer, S., Sobrino, C.L.,
770 Wang, H., Wei, Y., et al. (2014). A novel single-tailed fusiform *Sulfolobus* virus STSV2 infecting
771 model *Sulfolobus* species. *Extremophiles* 18, 51-60. 10.1007/s00792-013-0591-z.

772 Forterre, P., Krupovic, M., and Prangishvili, D. (2014). Cellular domains and viral lineages.
773 *Trends in microbiology* 22, 554-558. 10.1016/j.tim.2014.07.004.

774 Frost, A., Perera, R., Roux, A., Spasov, K., Destaing, O., Egelman, E.H., De, C.P., and Unger, V.M.
775 (2008). Structural basis of membrane invagination by F-BAR domains. *Cell* 132, 807-817.

776 Goulet, A., Vestergaard, G., Felisberto-Rodrigues, C., Campanacci, V., Garrett, R.A., Cambillau,
777 C., and Ortiz-Lombardia, M. (2010). Getting the best out of long-wavelength X-rays: de novo
778 chlorine/sulfur SAD phasing of a structural protein from ATV. *Acta crystallographica. Section D,*
779 *Biological crystallography* 66, 304-308. 10.1107/S0907444909051798.

780 Gudbergsdottir, S., Deng, L., Chen, Z., Jensen, J.V., Jensen, L.R., She, Q., and Garrett, R.A. (2011).
781 Dynamic properties of the *Sulfolobus* CRISPR/Cas and CRISPR/Cmr systems when challenged
782 with vector-borne viral and plasmid genes and protospacers. *Mol Microbiol* 79, 35-49.
783 10.1111/j.1365-2958.2010.07452.x.

784 Haring, M., Vestergaard, G., Rachel, R., Chen, L., Garrett, R.A., and Prangishvili, D. (2005).
785 Virology: independent virus development outside a host. *Nature* 436, 1101-1102.
786 10.1038/4361101a.

787 Hay, I.D., and Lithgow, T. (2019). Filamentous phages: masters of a microbial sharing economy.
788 *EMBO reports* 20. 10.15252/embr.201847427.

789 Hillyer, M.B., and Gibb, B.C. (2016). Molecular Shape and the Hydrophobic Effect. *Annu Rev*
790 *Phys Chem* 67, 307-329. 10.1146/annurev-physchem-040215-112316.

791 Hochstein, R., Bollschweiler, D., Dharmavaram, S., Lintner, N.G., Plitzko, J.M., Bruinsma, R.,
792 Engelhardt, H., Young, M.J., Klug, W.S., and Lawrence, C.M. (2018). Structural studies of
793 *Acidianus* tailed spindle virus reveal a structural paradigm used in the assembly of spindle-
794 shaped viruses. *Proc. Natl. Acad. Sci. U.S.A.* 115, 2120-2125. 10.1073/pnas.1719180115.

795 Hochstein, R.A., Amenabar, M.J., Munson-McGee, J.H., Boyd, E.S., and Young, M.J. (2016).
796 *Acidianus* Tailed Spindle Virus: a New Archaeal Large Tailed Spindle Virus Discovered by
797 Culture-Independent Methods. *J Virol* 90, 3458-3468. 10.1128/JVI.03098-15.

798 Hong, C., Pietila, M.K., Fu, C.J., Schmid, M.F., Bamford, D.H., and Chiu, W. (2015). Lemon-
799 shaped halo archaeal virus His1 with uniform tail but variable capsid structure. *Proc. Natl. Acad.*
800 *Sci. U.S.A.* 112, 2449-2454. 10.1073/pnas.1425008112.

801 Iranzo, J., Koonin, E.V., Prangishvili, D., and Krupovic, M. (2016). Bipartite Network Analysis of
802 the Archaeal Virosphere: Evolutionary Connections between Viruses and Capsidless Mobile
803 Elements. *J Virol* 90, 11043-11055. 10.1128/JVI.01622-16.

804 Iverson, E.A., Goodman, D.A., Gorchels, M.E., and Stedman, K.M. (2017). Genetic Analysis of the
805 Major Capsid Protein of the Archaeal Fusellovirus SSV1: Mutational Flexibility and
806 Conformational Change. *Genes (Basel)* 8. 10.3390/genes8120373.

807 Johnson, J.E., and Olson, A.J. (2021). Icosahedral virus structures and the protein data bank. *J*
808 *Biol Chem* 296, 100554. 10.1016/j.jbc.2021.100554.

809 Johnson, J.E., and Speir, J.A. (1997). Quasi-equivalent viruses: a paradigm for protein
810 assemblies. *J. Mol. Biol.* 269, 665-675. 10.1006/jmbi.1997.1068.

811 Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K.,
812 Bates, R., Zidek, A., Potapenko, A., et al. (2021). Highly accurate protein structure prediction
813 with AlphaFold. *Nature*. 10.1038/s41586-021-03819-2.

814 Kim, J.G., Kim, S.J., Cvirkaite-Krupovic, V., Yu, W.J., Gwak, J.H., Lopez-Perez, M., Rodriguez-
815 Valera, F., Krupovic, M., Cho, J.C., and Rhee, S.K. (2019). Spindle-shaped viruses infect marine
816 ammonia-oxidizing thaumarchaea. *Proc. Natl. Acad. Sci. U.S.A.* 116, 15645-15650.
817 10.1073/pnas.1905682116.

818 Knitsch, R., Schneefeld, M., Weitzel, K., and Pfeifer, F. (2017). Mutations in the major gas vesicle
819 protein GvpA and impacts on gas vesicle formation in *Haloferax volcanii*. *Mol Microbiol* 106,
820 530-542. 10.1111/mmi.13833.

821 Krupovic, M., Dolja, V.V., and Koonin, E.V. (2019). Origin of viruses: primordial replicators
822 recruiting capsids from hosts. *Nature reviews. Microbiology* 17, 449-458. 10.1038/s41579-019-
823 0205-6.

824 Krupovic, M., Dolja, V.V., and Koonin, E.V. (2020). The LUCA and its complex virome. Nature
825 reviews. Microbiology 18, 661-670. 10.1038/s41579-020-0408-x.

826 Krupovic, M., and Koonin, E.V. (2017). Multiple origins of viral capsid proteins from cellular
827 ancestors. Proc. Natl. Acad. Sci. U.S.A. 114, E2401-E2410. 10.1073/pnas.1621061114.

828 Krupovic, M., Quemin, E.R., Bamford, D.H., Forterre, P., and Prangishvili, D. (2014). Unification
829 of the globally distributed spindle-shaped viruses of the Archaea. J Virol 88, 2354-2358.
830 10.1128/JVI.02941-13.

831 Li, N., and Cannon, M.C. (1998). Gas vesicle genes identified in *Bacillus megaterium* and
832 functional expression in *Escherichia coli*. Journal of bacteriology 180, 2450-2458.

833 Li, Z., Jaroszewski, L., Iyer, M., Sedova, M., and Godzik, A. (2020). FATCAT 2.0: towards a better
834 understanding of the structural diversity of proteins. Nucleic Acids Res 48, W60-W64.
835 10.1093/nar/gkaa443.

836 Liu, J., Cvirkaite-Krupovic, V., Baquero, D.P., Yang, Y., Zhang, Q., Shen, Y., and Krupovic, M.
837 (2021). Virus-induced cell gigantism and asymmetric cell division in archaea. Proc. Natl. Acad.
838 Sci. U.S.A. 118. 10.1073/pnas.2022578118.

839 Medvedeva, S., Sun, J., Yutin, N., Koonin, E.V., Nunoura, T., Rinke, C., and Krupovic, M. (2021).
840 Viruses of Asgard archaea. bioRxiv, 2021.2007.2029.453957. 10.1101/2021.07.29.453957.

841 Mochizuki, T., Yoshida, T., Tanaka, R., Forterre, P., Sako, Y., and Prangishvili, D. (2010). Diversity
842 of viruses of the hyperthermophilic archaeal genus Aeropyrum, and isolation of the Aeropyrum

843 pernix bacilliform virus 1, APBV1, the first representative of the family *Clavaviridae*. *Virology*
844 402, 347-354. 10.1016/j.virol.2010.03.046.

845 Newberry, R.W., and Raines, R.T. (2019). Secondary Forces in Protein Folding. *ACS Chem Biol*
846 14, 1677-1686. 10.1021/acscchembio.9b00339.

847 Pei, J., and Grishin, N.V. (2014). PROMALS3D: multiple protein sequence alignment enhanced
848 with evolutionary and three-dimensional structural information. *Methods Mol Biol* 1079, 263-
849 271. 10.1007/978-1-62703-646-7_17.

850 Perotti, L.E., Dharmavaram, S., Klug, W.S., Marian, J., Rudnick, J., and Bruinsma, R.F. (2016).
851 Useful scars: Physics of the capsids of archaeal viruses. *Phys Rev E* 94, 012404.
852 10.1103/PhysRevE.94.012404.

853 Perotti, L.E., Zhang, K., Rudnick, J., and Bruinsma, R.F. (2019). Kirigami and the Caspar-Klug
854 construction for viral shells with negative Gauss curvature. *Phys Rev E* 99, 022413.
855 10.1103/PhysRevE.99.022413.

856 Pettersen, E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M., Meng, E.C., and
857 Ferrin, T.E. (2004). UCSF Chimera--a visualization system for exploratory research and analysis.
858 *J.Comput.Chem.* 25, 1605-1612.

859 Pfab, J., Phan, N.M., and Si, D. (2021). DeepTracer for fast de novo cryo-EM protein structure
860 modeling and special studies on CoV-related complexes. *Proc. Natl. Acad. Sci. U.S.A.* 118.
861 10.1073/pnas.2017525118.

862 Pfeifer, F., Gregor, D., Hofacker, A., Pl \check{d} sser, P., and Zimmermann, P. (2002). Regulation of gas
863 vesicle formation in halophilic archaea. *J. Mol. Microbiol. Biotechnol.* *4*, 175-181.

864 Pfitzner, A.K., Moser von Filseck, J., and Roux, A. (2021). Principles of membrane remodeling by
865 dynamic ESCRT-III polymers. *Trends in cell biology.* 10.1016/j.tcb.2021.04.005.

866 Pietila, M.K., Atanasova, N.S., Oksanen, H.M., and Bamford, D.H. (2013). Modified coat protein
867 forms the flexible spindle-shaped virion of haloarchaeal virus His1. *Environmental microbiology*
868 *15*, 1674-1686. 10.1111/1462-2920.12030.

869 Prangishvili, D., Bamford, D.H., Forterre, P., Iranzo, J., Koonin, E.V., and Krupovic, M. (2017). The
870 enigmatic archaeal virosphere. *Nature reviews. Microbiology* *15*, 724-739.
871 10.1038/nrmicro.2017.125.

872 Prangishvili, D., Vestergaard, G., Haring, M., Aramayo, R., Basta, T., Rachel, R., and Garrett, R.A.
873 (2006). Structural and genomic properties of the hyperthermophilic archaeal virus ATV with an
874 extracellular stage of the reproductive cycle. *J. Mol. Biol.* *359*, 1203-1216.
875 10.1016/j.jmb.2006.04.027.

876 Ptchelkine, D., Gillum, A., Mochizuki, T., Lucas-Staat, S., Liu, Y., Krupovic, M., Phillips, S.E.V.,
877 Prangishvili, D., and Huiskonen, J.T. (2017). Unique architecture of thermophilic archaeal virus
878 APBV1 and its genome packaging. *Nat. Commun.* *8*, 1436. 10.1038/s41467-017-01668-0.

879 Punjani, A., Rubinstein, J.L., Fleet, D.J., and Brubaker, M.A. (2017). cryoSPARC: algorithms for
880 rapid unsupervised cryo-EM structure determination. *Nat. Methods* *14*, 290-296.
881 10.1038/nmeth.4169.

882 Punjani, A., Zhang, H., and Fleet, D.J. (2020). Non-uniform refinement: adaptive regularization
883 improves single-particle cryo-EM reconstruction. *Nat. Methods* 17, 1214-1221.
884 10.1038/s41592-020-00990-8.

885 Quemin, E.R., Chlanda, P., Sachse, M., Forterre, P., Prangishvili, D., and Krupovic, M. (2016).
886 Eukaryotic-Like Virus Budding in Archaea. *mBio* 7. 10.1128/mBio.01439-16.

887 Quemin, E.R., Pietila, M.K., Oksanen, H.M., Forterre, P., Rijpstra, W.I., Schouten, S., Bamford,
888 D.H., Prangishvili, D., and Krupovic, M. (2015). Sulfolobus Spindle-Shaped Virus 1 Contains
889 Glycosylated Capsid Proteins, a Cellular Chromatin Protein, and Host-Derived Lipids. *J Virol* 89,
890 11681-11691. 10.1128/JVI.02270-15.

891 Raoult, D., and Forterre, P. (2008). Redefining viruses: lessons from Mimivirus. *Nature reviews.*
892 *Microbiology* 6, 315-319. 10.1038/nrmicro1858.

893 Rayment, I., Baker, T.S., Caspar, D.L., and Murakami, W.T. (1982). Polyoma virus capsid
894 structure at 22.5 Å resolution. *Nature* 295, 110-115. 10.1038/295110a0.

895 Rohou, A., and Grigorieff, N. (2015). CTFFIND4: Fast and accurate defocus estimation from
896 electron micrographs. *J. Struct. Biol.* 192, 216-221. 10.1016/j.jsb.2015.08.008.

897 Scheele, U., Erdmann, S., Ungewickell, E.J., Felisberto-Rodrigues, C., Ortiz-Lombardia, M., and
898 Garrett, R.A. (2011). Chaperone role for proteins p618 and p892 in the extracellular tail
899 development of Acidianus two-tailed virus. *J Virol* 85, 4812-4821. 10.1128/JVI.00072-11.

900 Sevana, M., Klose, T., and Rossmann, M.G. (2021). Principles of Virus Structure. Encyclopedia
901 of Virology, 257-277. 10.1016/B978-0-12-814515-9.00033-3.

902 Sheppard, C., Blombach, F., Belsom, A., Schulz, S., Daviter, T., Smollett, K., Mahieu, E., Erdmann,
903 S., Tinnefeld, P., Garrett, R., et al. (2016). Repression of RNA polymerase by the archaeo-viral
904 regulator ORF145/RIP. Nat. Commun. 7, 13595. 10.1038/ncomms13595.

905 Sojar, H.T., and Bahl, O.P. (1987). Chemical deglycosylation of glycoproteins. Meth. Enzymol.
906 138, 341-350.

907 Stedman, K.M., DeYoung, M., Saha, M., Sherman, M.B., and Morais, M.C. (2015). Structural
908 insights into the architecture of the hyperthermophilic Fusellovirus SSV1. Virology 474, 105-
909 109. 10.1016/j.virol.2014.10.014.

910 Stehle, T., Yan, Y., Benjamin, T.L., and Harrison, S.C. (1994). Structure of murine polyomavirus
911 complexed with an oligosaccharide receptor fragment. Nature 369, 160-163.
912 10.1038/369160a0.

913 Strunk, T., Hamacher, K., Hoffgaard, F., Engelhardt, H., Zillig, M.D., Faist, K., Wenzel, W., and
914 Pfeifer, F. (2011). Structural model of the gas vesicle protein GvpA and analysis of GvpA
915 mutants in vivo. Mol Microbiol 81, 56-68. 10.1111/j.1365-2958.2011.07669.x.

916 Uldahl, K.B., Jensen, S.B., Bhoobalan-Chitty, Y., Martinez-Alvarez, L., Papathanasiou, P., and
917 Peng, X. (2016). Life Cycle Characterization of Sulfolobus Monocaudavirus 1, an Extremophilic
918 Spindle-Shaped Virus with Extracellular Tail Development. J Virol 90, 5693-5699.
919 10.1128/JVI.00075-16.

920 Wang, F., Baquero, D.P., Su, Z., Beltran, L.C., Prangishvili, D., Krupovic, M., and Egelman, E.H.
921 (2020). The structures of two archaeal type IV pili illuminate evolutionary relationships. *Nat.*
922 *Commun.* *11*, 3424. 10.1038/s41467-020-17268-4.

923 Wang, F., Cvirkaite-Krupovic, V., Kreutzberger, M.A.B., Su, Z., de Oliveira, G.A.P., Osinski, T.,
924 Sherman, N., DiMaio, F., Wall, J.S., Prangishvili, D., et al. (2019). An extensively glycosylated
925 archaeal pilus survives extreme conditions. *Nat Microbiol* *4*, 1401-1410. 10.1038/s41564-019-
926 0458-x.

927 Weber, M.S., Eibauer, M., Sivagurunathan, S., Magin, T.M., Goldman, R.D., and Medalia, O.
928 (2021). Structural heterogeneity of cellular K5/K14 filaments as revealed by cryo-electron
929 microscopy. *eLife* *10*. 10.7554/eLife.70307.

930 Williams, C.J., Headd, J.J., Moriarty, N.W., Prisant, M.G., Videau, L.L., Deis, L.N., Verma, V.,
931 Keedy, D.A., Hintze, B.J., Chen, V.B., et al. (2018). MolProbity: More and better reference data
932 for improved all-atom structure validation. *Protein Science* *27*, 293-315. 10.1002/pro.3330.

933 Winkler, H. (2007). 3D reconstruction and processing of volumetric data in cryo-electron
934 tomography. *J. Struct. Biol.* *157*, 126-137. <https://doi.org/10.1016/j.jsb.2006.07.014>.

935 Xiang, X., Chen, L., Huang, X., Luo, Y., She, Q., and Huang, L. (2005). *Sulfolobus tengchongensis*
936 spindle-shaped virus STSV1: virus-host interactions and genomic features. *J Virol* *79*, 8677-8686.
937 10.1128/JVI.79.14.8677-8686.2005.

938 Zheng, S.Q., Palovcak, E., Armache, J.P., Verba, K.A., Cheng, Y., and Agard, D.A. (2017).
939 MotionCor2: anisotropic correction of beam-induced motion for improved cryo-electron
940 microscopy. *Nat. Methods* *14*, 331-332. 10.1038/nmeth.4193.

941 Zheng, W., Pena, A., Low, W.W., Wong, J.L.C., Frankel, G., and Egelman, E.H. (2020).
942 Cryoelectron-Microscopic Structure of the pKpQIL Conjugative Pili from Carbapenem-Resistant
943 *Klebsiella pneumoniae*. *Structure* *28*, 1321-1328 e1322. 10.1016/j.str.2020.08.010.

944 Zillig, W., Kletzin, A., Schleper, C., Holz, I., Janekovic, D., Hain, J., Lanzendörfer, M., and
945 Kristjansson, J.K. (1993). Screening for Sulfolobales, their plasmids and their viruses in *Icelandic*
946 *sofataras*. *Systematic and Applied Microbiology* *16*, 609-628.

947


TABLE FOR AUTHOR TO COMPLETE



Please upload the completed table as a separate document. **Please do not add subheadings to the key resources table.** If you wish to make an entry that does not fall into one of the subheadings below, please contact your handling editor. **Any subheadings not relevant to your study can be skipped.** (NOTE: For authors publishing in *Cell Genomics*, *Cell Reports Medicine*, *Current Biology*, and *Med*, please note that references within the KRT should be in numbered style rather than Harvard.)




Key resources table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Archaeal and virus strains		
SMV1	Uldahl et al., 2016	N/A
ATV	Prangishvili et al., 2006	N/A
<i>Saccharolobus islandicus</i> ΔC1C2	Gudbergsdottir et al., 2011	N/A
<i>Acidianus covivator</i> AA9	Prangishvili et al., 2006	N/A
Chemicals, peptides, and recombinant proteins		
Bacto Tryptone	Gibco	Cat #: 211705
Bacto yeast extract	Gibco	Cat #: 212750
Sucrose	Sigma-Aldrich	Cat #: S0389
PEG6000	Sigma-Aldrich	Cat #: 81260
NaCl	Sigma-Aldrich	Cat #: S9888
Trisma base	Sigma-Aldrich	Cat #: T1503
Acetic acid	Sigma-Aldrich	Cat #: A6283
CsCl	Euromedex	Cat #: EU0770
Ethidium bromide	Eurobio Scientific	Cat #: GEPBET02-AF
InstantBlue	Abcam	Cat #: ab119211
Deposited data		


SMV1 Cryo-EM maps	This paper	EMDB: EMD-24585, EMD-24586, EMD-24587, EMD-24588, EMD-24589, EMD-24590, EMD-24591, EMD-24592, EMD-24593, EMD-24594, EMD-24595, EMD-24597
SMV1 Cryo-EM atomic models	This paper	PDB: 7RO2, 7RO3, 7RO4, 7RO5, 7RO6, 7ROB, 7ROC, 7ROD, 7ROE, 7ROG, 7ROH, 7ROI
ATV Cryo-EM map	This paper	EMDB: EMD-24596
Software and algorithms		
cryoSPARC	Punjani et al., 2017	https://cryosparc.com
Chimera	Pettersen et al., 2004	https://www.cgl.ucsf.edu/chimera
DeepTracer	Pfab et al., 2021	https://deeptramer.uw.edu
Coot	Emsley and Cowtan, 2004	https://www2.mrc-lmb.cam.ac.uk/personal/pemsley/coot
Phenix	Afonine et al., 2018	https://phenix-online.org
MolProbity	Williams et al., 2018	http://molprobity.biochem.duke.edu
Psi-Blast	Altschul et al., 1997	https://www.ebi.ac.uk/Tools/sss/psiblast/
PROMALS3D	Pei and Grishin, 2014	http://prodata.swmed.edu/promals3d/promals3d.php
AlphaFold2	Jumper et al., 2021	https://alphafold.ebi.ac.uk
FATCAT	Li et al., 2020	http://fatcat.godziklab.org/









A




Interactions along the 7-start




Aligned by two subunits




B





Interactions between 7-start protofilaments






C










TMHMM transmembrane probabilities









A**B**